ISSN 0300-9246

JOURNAL OF THE CHEMICAL SOCIETY

Dalton Transactions

A journal of inorganic chemistry

1987

ROYAL SOCIETY OF CHEMISTRY

© The Royal Society of Chemistry, 1987. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form, or by any means, electronic, mechanical, photographic, recording, or otherwise, without the prior permission of the publishers.

Printed in Great Britain by
Richard Clay Ltd.,
Bungay, Suffolk

AUTHOR INDEX

Abad, J. Antonio, 41
Abel, Edward W., 417, 757
Abou-El-Wafa, Moustafa H. M., 1059, 2311
Abou-Hamdan, Amira, 489
Abrahams, Ian L., 1577
Abrams, Gillian, 1581
Adam, Kenneth R., 2537
Adams, Harry, 207, 341, 479, 1947, 2459
Adams, Jeanette S., 3049
Addison, Anthony W., 2621
Ahsen, Vefa, 1827
Aime, Silvio, 253, 349
Ainscough, Eric W., 493, 1583
Airoldi, Claudio, 2955
Al-Daffaee, Hafid K., 2615
Alami, Mahmoud Kalam, 1983
Albano, Vincenzo G., 21, 1133
Albertin, Gabriele, 1813
Alcock, Nathaniel W., 193, 545, 889, 1107, 1659, 1805, 2273, 2643, 2793, 3107
Alexander, Rikki P., 885, 2013
Allan, James, 2853
Allen, Geoffrey C., 3009
Allen, Stephen R., 591
Alonzo, Giuseppe, 789, 2337, 2451
Alvarez, Celia, 2045
Amadelli, Rossano, 349
Amer, Saiid A., 2009
Anderson, David G., 3029, 3035, 3061
Anderson, David M., 2843
Ando, Katsunori, 1977
Andreu, Ana María, 953
Antiñolo, Antonio, 975, 1463
Antolini, Luciano, 1921
Antoniutti, Stefano, 1813
Aoki, Katsuyuki, 1009, 2017
Apreda, Maria C., 981
Arcas, Aurelia, 1655
Arce, Alejo J., 1457, 1861, 2935
Arif, Atta M., 1665, 2159
Arif, Muhammad, 2397
Armstrong, David R., 1071
Armstrong, Janet A., 3029, 3061
Arnaiz, Francisco J., 819
Arriortua, Maria I., 285
Arthurs, Michael, 2615
Asaba, Yasuyuki, 1977
Asato, Eiji, 1305
Ascenzi, Paolo, 369
Ashcroft, Barry R., 2059, 2067
Ashmawy, Fathy M., 2009
Astapov, Boris, 1593
Attfield, Michael J., 2219
Atwood, Jerry L., 747
Auffret, Joëlle, 1687
Baccolini, Graziano, 21
Backer-Dirks, J. D. Julius, 1665
Bailey, Neil A., 207, 341, 479, 1865, 1947, 2519, 2815
Balakrishnan, Karappulli P., 545, 889
Baldas, John, 1721
Ballester, Loreto, 281
Bandoli, Giuliano, 867, 1693
Bandy, Judith A., 391
Banister, Arthur J., 915, 1565
Bannister, Colin, 341
Barbieri, Renato, 789
Bard, Allen J., 249
Barker, James, 2687
Barr, Donald, 1071, 2141
Barratt, David S., 2497, 2683
Barron, Andrew R., 837, 2947
Barron, Peter F., 1099
Barthelat, Jean-Claude, 2373
Bashall, Alan, 2543
Basu, Amitabha, 1797
Bataille, Michel, 661
Bates, Paul A., 551, 1529, 2397, 2935
Battaglia, Luigi P., 1333, 1779
Battle, Peter D., 667, 2359
Baumann, Franz-Erich, 2661, 2917
Bayón, J. Carlos, 3003
Beagley, Brian, 1999
Beardwood, Peter, 939
Beattie, Ian R., 267, 1521
Becke, Sigurd H. F., 1229
Beckett, Michael A., 417
Beevor, Robert G., 591, 1319
Begley, Michael J., 145
Behling, Torsten, 169
Bekâroğlu, Özer, 1827
Bell, Graeme A., 3023
Bellachioma, Gianfranco, 473, 519
Bellitto, Carlo, 35
Beltrán, Daniel, 1847
Beltran-Porter, Daniel, 285
Bentz, Peter O., 2709
Bergamini, Paola, 1659
Beringhelli, Tiziana, 2083
Bermúdez, M. Dolores, 1229, 1235
Berry, Frank J., 1727
Bertani, Roberta, 1381
Bertazzi, Nuccio, 2337, 2451
Berthon, Guy, 1145
Bertolasi, Valerio, 613, 2235
Besançon, Jack, 107
Betterton, Eric A., 761
Beveridge, Andrew D., 373
Bhargava, Suresh K., 757
Bhattacharjee, Manabendra N., 1055
Bhattacharjee, Manish, 477
Bianchi, Antonio, 1779
Bigoli, Francesco, 2407
Bingham, Alistair G., 493, 1583
Birchall, Thomas, 1675
Bismondo, Arturo, 695
Bitcon, Caroline, 3049
Bladon, Peter, 2503
Blagg, Adrian, 221, 769, 1495, 2171
Blake, Alexander J., 2853, 3035
Blakeley, Christine, 1775
Blanco, Severino García, 975
Blinn, Elliott L., 2599
Bögge, Hartmut, 493, 1583
Boas, John F., 1721
Bois, Claudette, 2875
Bombieri, Gabriella, 2337, 2451
Bonamico, Mario, 35
Bonnet, Jean-Jacques, 91
Bonnyman, John, 1721
Booth, Brian L., 1415, 2959
Bor, György, 1707
Bordignon, Emilio, 1813
Borrachero, M. Victoria, 1655
Bott, Simon G., 747
Botta, Mauro, 253
Bourdauducq, Paul, 1897
Bousher, Andrew, 943
Bowmaker, Graham A., 1089
Bown, Mark, 1169, 2781
Boyar, Esther, 551
Boyd, Peter D. W., 307
Boyde, Stephen, 297, 1083, 2267
Bracci, Michela, 1883
Bradley, Donald C., 2059, 2067, 2159
Braga, Dario, 1133
Braunstein, Pierre, 119
Bravo, Jorge, 683
Bray, Joanne M., 2989
Brianese, Nicola, 691, 2039
Brimblecombe, Peter, 943
Bristow, Garry S., 2843
Bristow, Stuart, 303
Brodie, Andrew M., 493, 1583
Brooker, Sally, 2555
Brown, David A., 2531
Brown, Jack R., 3049
Brown, Paul L., 2149
Brown, Scott S. D., 1967, 2333
Bruce, Duncan W., 2941
Bruck, Michael A., 2267
Bruno, Giuseppe, 2337, 2451
Bruno, Jordi, 2431, 2439, 2445
Brunori, Maurizio, 369
Bürger, Hans, 1839
Burman, Sudhir, 2621
Burnett, Michael G., 1059, 2311
Busch, Daryle H., 2505
Busetto, Luigi, 21, 1133
Byrne, Paul G., 1215, 1243
Caballol, Rosa, 2373
Cabeza, Javier A., 2459
Cabral, João de O., 2595
Cabral, M. Fermamda, 2595
Cairns, Colin J., 2505
Calligaris, Mario, 1381
Calvet, Josep, 119
Calvin, Melvin, 2105
Camellini, Marisa Tiripicchio, 1275, 1551
Cameron, Alan D., 2857
Cameron, T. Stanley, 969
Cano, Félix H., 981
Canty, Allan J., 1477
Carassiti, Vittorio, 349
Cardaci, Giuseppe, 473
Cardin, Christine J., 1641
Cardin, David J., 1641
Carlsen, Lars, 481
Carlton, Laurence, 895, 2741
Carmona, Daniel, 639
Carr, Stuart W., 1763, 3067
Carrera, Sagrario Martinez, 975
Carriedo, Gabino A., 1769, 2867
Carthey, Nicholas A., 2329
Carty, Arthur J., 1463
Casals, Isidre, 2391
Casas, José S., 683
Casellato, Umberto, 2039, 2299, 3107
Cassol, Alberto, 657
Castilla, Maria Luz, 3017
Castiñeiras, Alfonso, 683
Cecconi, Franco, 831
Cenini, Sergio, 605, 1583
Cesarotti, Edoardo, 1149
Chadha, Sunehri L., 1253
Chamberlain, Linda R., 155, 163
Chan, Lisa, 513
Chandra, Harish, 729
Charnock, John M., 303
Chatterjee, Chinmay, 2095
Chaudhuri, Mihir K., 477, 1055
Chaudhuri, Nirmalendu Ray, 997
Chaudhury, Muktimoy, 1733, 3043
Chemaly, Susan M., 761
Chicote, María-Teresa, 881
Chiesi-Villa, Angiola, 1367, 1789, 1883, 2099
Chimatadar, Shivamurti A., 573, 2291
Chiu, K.-vok W., 733
Chopra, Suman K., 2927
Ciani, Gianfranco, 1149
Cihon, Irene, 2599
Cipriano, Robert A., 1901
Ciriano, Miguel A., 981
Clague, Diane, 1733
Claire, Preet P. K., 263
Clark, George R., 2067
Clark, Robin J. H., 1535
Clayden, Nigel J., 1877
Clegg, William, 303, 1071, 1083, 1577, 2141
Clemente, Dore A., 1823
Clucas, Jennifer A., 277
Cohen, Abraham, 235
Cole-Hamilton, David J., 443, 2683, 2941
Collison, David, 111, 3049
Collyer, Charles A., 2573
Colmanet, Silvano F., 2305
Connelly, Neil G., 467, 1403, 1769, 2737
Constable, Edwin C., 5, 293, 2281
Contractor, Sorab R., 2023
Cook, Malcolm R., 2757
Corazza, Fabio, 709
Coronado, Eugenio, 1847
Corradi, Anna Bonamartini, 1333, 1779
Countryman, Rachel M., 193
Courtot, Pierre, 1687
Couture, Christiane, 2245, 2253
Cowie, Alan G., 2839
Cowley, Alan H., 249
Cradock, Stephen, 3029, 3035, 3061
Craig, Scott, 2605
Cresswell, Clifford J., 795
Crocker, Mark, 2803
Cross, Ronald J., 139
Cruywagen, Johannes J., 1701
Cuadro, Ana M., 183
Cunningham, Desmond, 2261, 2927
Cunningham, Michael, 3049
Cupertino, Domenico C., 443

Curzon, Eirian, 2615
Díez, Josefina, 1275
D'Alfonso, Giuseppe, 2083
Dagnall, Stephen P., 2897
Dahan, Françoise, 1983
Dahlstrom, Phillip, 529
Darriet, Jacques, 285
Dartmann, Mechtild, 2579
Das, V. G. Kumar, 129
Dauter, Zbigniew, 27
Davidson, Jack L., 895, 2165, 2715
Davidson, Michael F., 139
Davies, David L., 1201
Davies, Simon J., 2177
Davis, William M., 2505
Dawes, Helen, 2159
Démarcq, Michel C., 1897
De, Goutam, 997
de Ilarduya, Jesús Martinez, 975
de Souza, Antonio G., 2955
De Sanctis, Ysaura, 1457, 2935
DeArmond, M. Keith, 2467
Deeming, Antony J., 551, 1457, 1529, 1861, 1869, 2473, 2935
Delgado, Esther, 41, 1191, 1235
Dell'Amico, Daniela Belli, 1329
Demartin, Francesco, 21, 605, 1583
Dengel, Andrew C., 991
Denholm, Stella, 2789
Dent-Glasser, Lesley S., 2323
Deplano, Paolo, 2407
der Merwe, Izak F. J. van, 1701
Desseyn, Herman O., 1921
Dessy, Giulia, 35
Devi, Meenakshi, 1055
Dhubhghaill, Orla Ni., 699
Di Bernardo, Plinio, 657, 695
Dickson, Mark K., 2769
Dillon, Keith B., 315
Dilworth, Jonathan R., 529
Ditzel, Evert J., 1285, 1289, 1293, 1623
Dommisse, Roger, 1921
Donovan-Mtunzi, Susan, 1457, 1869
Drake, Simon R., 1051
Drew, Michael G. B., 123, 263, 873, 1027, 1163, 1569, 1605, 2563, 2975
Dromzée, Yves, 819
Duatti, Adriano, 613, 867, 2299, 3107
Duffy, John A., 1485, 2323
Dyason, Jeffrey C., 1089, 1099
Eaborn, Colin, 747
Earnshaw, Alan, 97
Ebsworth, E. A. V., 2831, 3035
Edwards, Howell G. M., 681
Edwards, Peter G., 169
Egert, Ernst, 847
Eggleston, Drake S., 677
Egsgaard, Helge, 481
Ekstrom, Alfred, 2537
Elbel, Susanne, 481
Elliott, Gregory P., 1235, 2177, 2189
Elrington, Margaret, 451
Emsley, John, 1271, 2397
Enemark, John H., 297, 619, 2267
Engelhardt, Lutz M., 457, 1089, 1099, 1463, 1647, 2347
Ercolani, Claudio, 369, 369
Errington, R. John, 2059, 2067
Escrivá, Emilio, 1847
Espinet, Pablo, 2077
Ettorre, Renato, 815
Evans, David G., 1889
Evans, Dennis F., 1279, 3099
Fabbrizzi, Luigi, 2631
Fabretti, Antonio C., 1921
Facchin, Giacomo, 1381
Faers, Malcolm A., 341
Faggiani, Romolo, 1675
Fantucci, Piercarlo, 355, 355
Faraglia, Giuseppina, 1823
Fares, Vincenzo, 35, 1035
Farrar, David H., 1249, 2245, 2253
Farrugia, Louis J., 51, 1191
Fava, Giovanna Gasparri, 227
Fawcett, John, 567, 733
Fedorko, Peter, 341
Fenton, David E., 207, 219, 1865, 2519, 2543
Ferguson, George, 699
Ferrari, Marisa Belicchi, 227
Ferretti, Valeria, 613, 2235
Ferri, Diego, 2439
Fiallo, Marina, 1367
Field, John S., 1933
Figgis, Brian N., 1737, 1747
Fittschen, Christa, 881, 3017
Fitzgerald, Joan, 2261
Flamini, Alberto, 35
Flint, Colin D., 939, 1929
Floriani, Carlo, 709, 1367, 1789, 1883, 2099
Foces-Foces, Concepción, 981
Folting, Kirsten, 155
Fompeyrine, Patricia, 91
Font-Bardía, Mercé, 1619, 2391
Fontaine, Xavier L. R., 777, 1169, 1431, 1501, 1573, 2385, 2401, 2417, 2781, 3067
Fontanille, Michel, 1041
Forniés, Juan, 2077
Forsyth, George A., 1027
Fortuño, Consuelo, 2077
Foster, Douglas F., 277
Fowkes, Hayat, 1431, 2417
Franchini, Giancarlo, 1921
Freeman, Hans C., 2573
Freeman, Mark J., 379 1001, 1283
Frigo, Dario M., 2159
Fritzer, Harald P., 939
Frlec, Boris, 2379
Fronczek, Frank R., 2769
Froom, Simon F. T., 2803
Fuertes, Amparo, 1847
Fuess, Hartmut, 431
Fujita, Masanori, 1977
Fukuhara, Choki, 1305
Gahan, Brendan, 111
Gamasa, M. Pilar, 1275
Gambarotta, Sandro, 1789, 1883, 2099
Ganapathiappan, Sivapackia, 579, 585
Gans, Peter, 673
Gantar, Darja, 2379
García, José R., 2045
Garcia, Gabriel, 819, 1403, 2737
Garcia, M. Esther, 41, 1209, 1215, 1243, 2201
Garner, C. David, 297, 303, 1083, 1577, 2267
Gayoso, Miguel, 683
Geiger, William E., 467
Gentil, Luis A., 1267
Geraty, Suzanne M., 3073
Gervasio, Giuliana, 1707
Ghilardi, Carlo A., 831
Ghosh, Ashutosh, 997
Gibb, Terence C., 667, 1419, 2359
Gibson, John F., 939
Gilbert, Mark, 1403
Gill, J. Bernard, 673, 2995
Gilli, Gastone, 613, 2235
Gilmour, David I., 335
Gilson, Trevor R., 267
Gimeno, José, 1275
Giordano, Thomas J., 1175
Giuliani, Anna Maria, 1035
Giusti, Jol, 1779
Glass, William K., 2531
Glidewell, Christopher, 501, 509, 2981
Gökçeli, Füsun, 1827
Gómez-Sal, M. Pilar, 1623
Gómez-Sal, Pilar, 975
Gobetto, Roberto, 253
Goggin, Peter L., 2411
Gohshi, Yohichi, 3093
Gold, Victor, 1271
González-Casso, Maria E., 281
González-Duarte, Pilar, 2391
Good, Rosalind J., 207
Goodall, David C., 2995
Goswami, Kisholoy, 2105
Gott, George A., 1785, 2241
Gould, Robert O., 2687, 2831
Govan, Norman, 963
Graham, William A. G., 1181
Grant, Janice L., 2105
Grassi, Maria, 2649
Gray, Colin J., 1665, 2289
Graziani, Rodolfo, 2039, 2299, 3107
Grebenik, Peter D., 9
Green, Jennifer C., 391, 391
Green, Malcolm L. H., 9, 391, 403, 411, 1611, 2031, 2823
Green, Michael, 61, 73, 81, 591, 1313, 1319, 2803
Greenwood, Norman N., 275, 451, 1169, 1431, 2385, 2417, 2781
Gregson, David, 431
Grenthe, Ingmar, 2439, 2445
Griffith, William P., 991
Grosse-Ophoff, Markus J., 41
Grossel, Martin C., 705
Grove, David M., 1359
Grzybkowski, Wacław, 2863
Gündüz, Necla, 925
Gündüz, Turgut, 925
Guastini, Carlo, 1367, 1789, 1883, 2099
Guimon, Claude, 107
Gutowsky, Herbert S., 2317
Hägele, Gerhard, 795
Härter, Peter, 2757
Haasnoot, Jaap G., 1389
Hage, Ronald, 1389
Hague, David N., 2889, 2897
Haiduc, Ionel, 145, 1257
Haines, Raymond J., 1933
Hallam, Malcolm F., 1889
Hambley, Trevor W., 489
Hamor, Thomas A., 1063, 2653
Hanabusa, Kenji, 1801
Hanck, Kenneth W., 2467
Hancock, Robert D., 2911
Hansen, Lars K., 27
Hanson, Brian E., 1285
Hanson, Graeme, 1755
Harbron, Stephen K., 633
Harding, Marjorie M., 277
Harding, P. Alexander, 947, 2537
Harris, Nigel J., 1683
Harris, Robin K., 795
Harrison, Philip G., 807
Harrison, W. David, 2995
Hart, F. Alan, 1665, 2289
Hart, Ian J., 41
Hasan, Masihul, 1027
Hashimoto, Yukio, 2963
Haslop, John, 2615
Hata, Kazuhiko, 1127
Hauptman, Zdenek V., 915
Hausser-Wallis, Renée, 1839
Hay, Robert W., 963, 2605
Healy, Peter C., 1089, 1099
Heath, Graham A., 249
Heckman, Roger A., 2505
Hein, Joachim, 1191, 2211
Hempstead, Michael R., 2511
Henderson, Richard A., 197, 529, 1115
Henderson, William, 733
Henrick, Kim, 2537, 2669
Herber, Rolfe H., 789
Herbst, Regine, 847
Higgins, Simon J., 777, 1501
Hill, William E., 1581
Hiller, Wolfgang, 683, 1329
Hills, Elaine F., 1397, 2293
Hirotsu, Takahiro, 2489
Hitchcock, Peter B., 319, 747, 2635, 2843
Hitchman, Michael A., 457, 825, 1817
Hobson, Richard J., 1027, 1163, 1569
Hodge, Stephen R., 931
Hofmans, Hendrik C., 1921
Holden, John G., 1727
Holloway, John H., 741, 2285, 2379
Holmes-Smith, Rupert D., 969
Holmes, Jeremy M., 5
Holmes, Nigel R., 3009
Horton, Andrew D., 1557
Hough, Edward, 427
Hounslow, Andrea M., 489
Howard, Judith A. K., 61, 73, 81, 259, 1221, 1229, 2177, 2189, 2219, 2661, 2917
Hoyano, James K., 1181
Hudson, Stephen, 1967
Huffman, John C., 155
Hughes, David L., 1115
Hunter, Geoffrey, 1545, 2789
Hursthouse, Michael B., 169, 551, 557, 837, 1529, 1655, 1665, 1861, 1869, 2059, 2067, 2159, 2397, 2473, 2723, 2935, 2947, 2981
Hussain, Bilquis, 557, 2159
Hussain, Wasif, 619
Hynes, Michael J., 563
Iggo, Jonathan A., 2669
Iitaka, Yoichi, 853
Iliopoulos, Peter, 1585
Imoto, Tomosaku, 1801
Imperatori, Patrizia, 1035
Inoue, Hidenari, 2455
Issa, Rafaat M., 2009
Itagaki, Takaharu, 2489
Ito, Kaname, 1853
Ito, Takashi, 1857
Ivan Leban, 2379
Izquierdo, Alejo, 9
Jackman, Martin P., 2573
Jacobsen, Grant B., 151, 1489, 1509, 2005, 2751, 3079
Jageland, Per T., 795
Jalón, Félix A., 953

James, Alun P., 61, 81, 1221
James, Tony D., 2013
Janecki, Tomasz, 2503
Jasim, Hatam A., 2843
Jassal, Varinder S., 2815
Jasztal, Michael J., 2793
Jeannin, Yves, 819, 2875
Jeffery, John C., 41, 51, 1191, 1209, 1215, 1243, 2201, 2211
Jelfs, Alasdair N. de M., 73, 81, 1221, 2219
Jemmis, Eluvathingal D., 271
Jobe, Ian R., 2117
Jochem, Klaus, 969
Johnson, Brian F. G., 327, 931, 1051, 1285, 1289, 1293, 1623, 2839
Johnson, Lawrence H., 673
Johnson, Owen, 73, 1229, 2661, 2917
Johnston, Roy L., 647, 1445
Jones, Christopher J., 619, 1063, 1733, 2653
Jones, Deborah J., 2111
Jones, Peter G., 881, 3017
Jones, Peter J., 267
Jones, Ray, 907
Joyner, David S., 2823
Kabir, Shariff E., 1457, 1529, 1869
Kaden, Thomas A., 2631
Kang, Ding-Xue, 2599
Kato, Kazumasa, 1801
Kato, Noriyuki, 1341
Katoh, Shunsaku, 2489
Katsura, Kosho, 1305
Katti, Kattesh V., 847
Kavanagh, Seamus, 2927
Kemball, Alan C., 2669
Kemmitt, Raymond D. W., 733
Kemp, Terence J., 1659
Kendrick, Aidan G., 915, 1565
Kennedy, Brendan J., 825
Kennedy, John D., 275, 451, 1169, 1431, 1573, 2385, 2401, 2417, 2781
Keong, Yap Chee, 129
Kerridge, David H., 2329
Kida, Sigeo, 1157, 1261, 1957, 3055
Kieboom, Antonius P. G., 2051
Kikabhai, Thakor, 807
Kilduff, Jan E., 249
Kılıc, Esma, 925
Kılıç, Zeynel, 2023
Kilner, Melvyn, 2687
King, Christopher, 2769
Kino, Koji, 1157, 1957
Kinomura, Nobukazu, 609
Kintz, Andy, 2599
Kisenyi, Jonathan M., 1605, 2459
Kiss, Tamas, 1263
Kitaura, Eiji, 3055
Kitazaki, Kenji, 1341
Knox, Selby A. R., 2087, 2703
Kobayashi, Masayuki, 609
Kobayashi, Nagao, 1801
Kohzuki, Masatomo, 3093
Kolowich, J. Bruce, 3003
Komura, Michihiro, 1953
Kowalik, Teresa, 1
Kozłowski, Henryk, 1, 661
Krebs, Bernt, 2579
Krishnamurthy, Setharampattu S., 579, 585
Kristofzski, John G., 2267
Kückelhaus, Wolfgang, 795
Kubal, Gina, 2615
Kuhn, Norbert, 541
Kumada, Nobuhiro, 609
Kumar, N. R. Sreenivas, 271
Kumar, P. N. V. Pavan, 271
Kumar, Sujit Baran, 3043
Kunwar, Ajit C., 2317
Kupče, Ēriks, 1593
Kupryszewski, Godfryd, 1
Kwan, Linda, 457
Laguna, Antonio, 3017
Laguna, Mariano, 3017
Lam, Oi M., 97
Lamata, M. Pilar, 639
Lane, Janette D., 197
Lanfredi, Anna M. Manotti, 953
Lang, Peter F., 1929
Langrick, C. Richard, 777
Lappert, Michael F., 1463, 2635, 2843
Larkworthy, Leslie F., 965
Latour, Jean Marc, 2519
Lavery, Aidan J., 373, 2975
Lavigne, Guy, 91
Lawrance, Geoffrey A., 1425
Laycock, David E., 2857
Lees, Alistair J., 513
Leland, Jonathan K., 249
Leong, Anthony J., 2537, 2543
Leporati, Enrico, 435, 1283, 1409, 1583
Leung, Wing-Por, 2347
Levason, William, 633, 1901
Lever, Alfred B. P., 2511
Lewandos, Glenn S., 2703
Lewis, Jack, 327, 931, 1051, 1285, 1289, 1293, 1623, 2839
Leznoff, Clifford C., 2511
Li, Ke-Si, 2959
Lichon, Michael, 1817
Liepa, Imants, 2537
Liepiņš, Edvards, 1593
Liles, David C., 1631
Limosin, Daniele, 2519
Lincoln, Stephen F., 489, 2885
Lindoy, Leonard F., 2537, 2543
Lindsay, Alan J., 2723
Lindsell, W. Edward, 2741
Little, Mark, 2261
Liu, Wei, 2511
Livera, Cynara E., 661
Llavona, Ricardo, 2045
Lloyd, Brian R., 201
Lloyd, Douglas, 501, 509, 2981
Lock, Colin J. L., 1675
Loizidou, Maria, 1911
Longoni, Giuliano, 685
Loretto, Michael H., 1727
Lortz, Wolfgang, 623
Lowe, Barry M., 1271
Luke, Michael A., 335
Lukevics, Edmunds, 1593
Lumbard, Keith W., 501, 509, 2981
Luna, Silvio Alvarez, 867
Lunniss, Julie A., 1249
Mabbs, Frank E., 111
MacCurtain, Jacqui A., 699
Mackay, Kenneth M., 2365
Mackay, Maureen F., 2305
MacKenzie, Thomas E., 757
MacKinnon, Peter I., 2385, 2781
Macpherson, Ian, 2385
Maeda, Hiromi, 2455, 2963
Maeda, Katsunori, 961
Maeda, Masunobu, 1853
Maeda, Yonezo, 627, 1341
Magon, Luciano, 613, 2299, 3107
Maitlis, Peter M., 639, 1963, 2459, 2709
Mak, Thomas C. W., 129
Makani, Thomas, 2111
Makino, Tadashi, 1127
Malik, K. M. Abdul, 1869
Mangani, Stefano, 2999
Manivannan, Veeragathy, 1675
Mann, Brian E., 1947, 1963, 2649, 2709
Manohar, Hattikudur, 1009
Manojlović-Muir, Ljubica, 201, 2117
Marangoni, Giampaolo, 2235
Marchi, Andrea, 613, 867, 2299, 3107
Marcos, Enrique Sánchez, 2373
Mari, Marina, 1381
Markwell Anthony J., 1349
Marsden, Christine, 51
Marshall, Andrew, 315
Martín, José, 881
Maruta, Katsuhiko, 361
Mason, Mark, 2599
Mathieu, René, 1619, 1983
Matsubayashi, Gen-etsu, 1793
Matsuda, Kiyoshi, 1833
Matsumoto, Keiji, 1305
Maverick, Andrew W., 2881
Mawby, Roger J., 27, 1713, 2989
Maya, Beverley J., 2117
Mayo, Richard A., 2831
Mayor-Real, Fern, 1235
Mays, Martin J., 1557, 2669
Mazzi, Ulderico, 867, 1693
McArdle, James V., 677
McArdle, Patrick, 2927
McAuliffe, Charles A., 1299, 1415, 1785, 1999, 2009, 2241, 2497, 2959
McCleverty, Jon A., 619, 1063, 1733, 2653
McCooey, K. Michelle, 1713
McCool, Brian J., 2537
McCullagh, John F., 1059, 2311
McCullough, Kevin J., 2503, 2741
McDonald, Robbie G., 1817
McGinnis, Joseph, 2573
McKechnie, James S., 2981
McKee, Vickie, 523, 2519, 2555
McLennan, Alistair J., 2761
McMaster, Alexander D., 1181
McPartlin, Mary, 327, 1631, 1967, 2537, 2543, 2669
McQueen, Roderick C. S., 5, 1051
Melník, Milan, 2511
Melnyk, Alexandra C., 2505
Menabue, Ledi, 1333, 1921
Meunier, Philippe, 107
Michalczyk, Michael J., 2635
Micskei, Károly, 255
Midgley, Derek, 943
Midollini, Stefano, 831
Miguel, Daniel, 2867, 2875
Miguel, Jesus A., 2875
Mikuriya, Masahiro, 1261
Millar, John C., 2165
Miller, J. David, 985
Millington, Keith R., 1521
Milone, Luciano, 349
Minchin, Nigel J., 1477
Minelli, Martin, 619
Mingos, D. Michael P., 335, 647, 1445, 1889
Minten, Karl, 1999
Miravitlles, Carlos, 1847
Mise, Takaya, 2189
Mishiro, Junichi, 1341
Missen, Paul H., 1279
Mitchell, Gary F., 1017
Mitra, Samiran, 2089
Mittal, Pramod K., 757
Miwa, Makoto, 961
Miyamura, Kazuo, 1127, 3093
Miyazaki, Toshiaki, 961
Moehring, Gregory A., 715
Moldes, Isabel, 1619
Molinari, Henriette, 2083
Molloy, Kieran C., 101
Monacelli, Fabrizio, 369
Monari, Magda, 1133
Montagna, Laura, 2631
Moodie, W. Thomas, 2605
Moody, Richard, 207, 219, 2519
Moore, Laila S., 2333
Moore, Madeleine H., 2125
Moore, Peter, 545, 889, 1107, 2643
Moran, Grainne, 2927
Moreton, Anthony D., 2889, 2897
Mori, Masayasu, 1305
Morris, Michael J., 2087
Morton-Blake, D. Anthony, 1641
Motevalli, Majid, 169, 557, 837, 2723, 2947
Moulding, Roger P., 705
Mtetwa, Victor S. B., 9
Müller, Achim, 493, 1583
Mu-guang, Liu, 1415
Muir, Kenneth W., 2165
Mukherjee, Alok K., 997
Mukherjee, Monika, 997
Mulvey, Robert E., 1071, 2141
Mumba, Peter P. E. M., 1163, 1569
Muñoz, Maria, 2445
Murase, Ichiro, 1261
Murata, Mitsuo, 2573
Murata, Yutaka, 1853
Murgich, Juan, 1637
Murphy, Brian P., 123, 873, 2543
Murray, Keith S., 825, 1585
Muto, Fumio, 609
Nagle, Katharine R., 2803
Nakai, Hiroshi, 1953
Nakano, Toshiyuki, 1857
Nandana, W. A. Shantha, 1989
Nandibewoor, Sharanappa T., 573, 2291
Nardin, Giorgio, 1381
Narushima, Ryoichi, 3093
Nassimbeni, Luigi R., 2125
Navarro, Rafael, 2077
Neissl, Wolfgang, 939
Nelson, Jane, 123, 873
Nelson, S. Martin, 123, 873, 2975
Nelson, William J. H., 327
Net, Gemma, 3003
Nevin, W. Andrew, 2511
Newberry, Vincent, 1299
Ngwenya, M. Patrick, 2911
Nicholson, David G., 427
Nicholson, John R., 303
Nicholson, Terrence, 529
Nicola, Giuseppe, 349
Nicolini, Marino, 1693
Nicoló, Franco, 2451
Nielson, Alastair J., 307, 2059, 2067
Nishida, Yuzo, 1157, 1957, 3055
Nishide, Hiroyuki, 1917, 2455, 2493, 2963

Niven, Margaret L., 2125
Nomiya, Kenji, 961
Norén, Bertil, 3103
Norman, Nicholas C., 249
Norman, Richard E., 2905
Nowell, Ian W., 101
Nuber, Bernhard, 187
Nunn, Christine M., 61, 81, 1221, 2177, 2189, 2219
Öztaş, S. Gül, 925
Ōkawa, Hisashi, 3055
O'Dowd, Maire, 563
O'Hare, Dermot, 391, 403, 2031
Obaidi, Najat Al, 1063, 1733, 2653
Oberhammer, Heinz, 1839
Ogawa, Nobuhiro, 627
Ogden, J. Steven, 859
Ogura, Kotaro, 1377
Ohgomori, Yuji, 2969
Ohkubo, Katsutoshi, 361
Ohshiba, Masashi, 609
Oja, Tonis, 1637
Ojo, J. Folorunso, 957
Olabe, José A., 1267
Oldfield, Eric, 2317
Olubuyide, Olusegun, 957
Omar, Hadi A. A., 545, 1107, 2643
Oohama, Tadashi, 853
Ooi, Shun'ichiro, 1305
Orioli, Pierluigi, 2999
Orlandini, Annabella, 831
Oro, Luis A., 639, 981
Orpen, A. Guy, 379, 591, 1001, 1283, 1313, 1319, 2703, 2803
Orrell, Keith G., 757
Ortega, Richard B., 297
Osa, Tetsuo, 1801
Osborn, Valerie A., 341, 479
Osella, Domenico, 253, 349
Oshio, Hiroki, 1341
Ossola, Franco, 691, 2039
Otero, Antonio, 953, 975
Otvos, John W., 2105
Oyetunji, Olayinka, 957
Pacchioni, Gianfranco, 355
Paddick, Kathleen E., 591
Padovan, Franco, 253
Pakawatchai, Chaveng, 1089, 1099
Pakkanen, Tapani A., 781
Pakulski, Marek, 249
Palenik, Gus J., 1175
Pannell, Lewis K., 2555
Paoletti, Piero, 1779
Papasergio, Rocco I., 1647, 2347, 3085
Parge, Hans E., 1641
Parish, R. V. ('Dick'), 1605, 2009, 2333
Parkes, Harold G., 2479
Parkin, Gerard, 1611
Parlier, Andrée, 1041
Parrott, Michael J., 1201
Parvez, Masood, 699
Passmore, Jack, 1989
Patrick, Vincent A., 1099
Pauson, Peter L., 2503, 2757
Pawelke, Gottfried, 1839
Paxton, Keith, 1063, 2653
Peacock, Raymond D., 567
Pearsall, Mary Ann, 327
Pearson, Julie C., 1733
Pearson, Matthew D., 2615
Pelizzi, Corrado, 177, 227
Pelizzi, Giancarlo, 177
Pellacani, Gian Carlo, 1921
Pellinghelli, Maria Angela, 2407
Peloso, Arnaldo, 1473
Pennesi, Giovanna, 369
Perly, Bruno, 661
Perotti, Angelo, 2605
Perpiñán, Maria F., 281
Persson, Ingmar, 2411
Peruzzo, Valerio, 815
Peters, Joop A., 2051
Petocz, György, 1717
Pettit, Leslie D., 661
Pfister-Giullouzo, Geneviève, 107
Phadke, Sanjiv Vasant, 2095
Philoche-Levisalles, Michèle, 2875
Pichon, Roger, 1687
Pickup, Barry T., 2649
Pike, Graham A., 889
Pilbrow, John R., 1755, 2341
Pitteri, Bruno, 2235
Pizzotti, Maddalena, 605, 1583
Platt, Andrew W. G., 2273
Platzer, Nicole, 1041
Plazzogna, Gualtiero, 815
Pletcher, Derek, 1901
Poggi, Antonio, 2631
Pogorzelec, Peter, 2941
Pohl, Klaus, 187
Pombeiro, Armando L. J., 319
Porchia, Marina, 691, 2039
Porta, Francesca, 605, 1583
Portanova, Roberto, 657
Powell, Nicholas I., 1529
Powell, Nigel A., 1901
Powell, Richard D., 991
Prabhakaran, C. P., 965
Prati, Laura, 1149
Pratt, John M., 761, 1349
Preston, Peter N., 2741
Pringle, Paul G., 1495, 1659, 2273
Prins, Rob, 1389
Pritchard, Robin G., 1999
Probitts, E. Jane, 1713
Prout, K., 9
Puddephatt, Richard J., 201, 2117
Puebla, M. Pilar, 639
Pujari, Mahesh P., 963, 2605
Pursiainen, Jouni, 781
Quill, Kieran, 101
Radomska, Barbara, 661
Raithby, Paul R., 327, 931, 1557, 1623, 2281, 2669
Rajan, Odatt A., 2621
Raju, Javali R., 573, 2291
Randle, Neil P., 551, 2473
Rankin, David W. H., 785, 3023, 3035, 3061
Rao, D. N. Ramakrishna, 729
Rao, S. P. Sudhakara, 1009
Rasmussen, Paul G., 3003
Raston, Colin L., 1463, 1647, 2347, 3085
Ratermann, Andrew L., 1191
Raven, Stephen J., 467, 1769
Raynor, J. Barrie, 421
Reader, C. James, 2643
Reed, David, 699, 1071, 2761
Reedijk, Jan, 1389, 2579
Refosco, Fiorenzo, 1693
Reichenbach, Gustavo, 519
Reinisch, Paul F., 3023
Rest, Antony J., 1181
Reynolds, Colin D., 27
Reynolds, Philip A., 1737, 1747
Rice, David A., 1027, 1163, 1569
Richards, Raymond L., 319, 1115
Richens, David T., 2605
Rickard, Clifton E. F., 307, 2067
Riera, Victor, 819, 1551, 1769, 2867, 2875
Rizzo, Luigi, 695
Roberts, Patrick B., 1865
Robertson, Heather E., 785, 3035
Robinson, Stephen D., 947
Robson, Richard, 1585, 2171
Rodríguez, Julio, 2045
Rodriguez de Barbarin, Cecilia O., 207, 2519
Roesky, Herbert W., 847
Rojo, Teofilo, 285
Rolka, Krzysztof, 1
Romero, Antonio, 183
Roncarolo, Antonella, 21
Rook, John, 2285
Ros, Josep, 1619
Rosales, Maria J., 327
Rose, Norman J., 2905
Rossell, Oriol, 119
Rossetti, Rosanna, 1707
Rossetto, Gilberto, 691, 2039
Rossi, Roberto, 2299, 3107
Rothwell, Ian P., 155, 163
Roundhill, D. Max, 2769
Rowbottom, Graham L., 825
Rowe, Teresa M., 2621
Rowland, Ian J., 421
Roy, Abhijit, 1641
Royo, Pascual, 953, 975
Rozière, Jacques, 2111
Rudler, Henri, 1041
Ruiz, Javier, 285
Ruiz, Jose, 639, 1963, 2709
Ruiz, Miguel A., 1551
Russell, David R., 567, 733
Šraga, Ján, 2757
Saburi, Masahiko, 1127, 3093
Sakaki, Shigeyoshi, 361
Sakurai, Tosio, 721, 3093
Saladini, Monica, 1333
Salam, Sithy S., 1733
Salaün, Jean-Yves, 1687
Salem, Geoffrey, 1647
Salt, Julian E., 2947
Salt, Nicholas J., 2459
Salter, Ian D., 379, 1001, 1283, 1967, 2333
Sambrani, Mahantesh I., 573, 2291
Sampson, Clifford N., 1933
Sandström, Magnus, 2411, 2439
Sangster, David F., 1425
Santos, Amelia, 183, 281
Saple, Ashok R., 1797
Sapre, Niteen Y., 1797
Saunders, David R., 27
Savage, Paul D., 557
Schaverien, Colin J., 1313
Schmidt, Hans-Georg, 847
Schön, Günter, 623
Schröder, Martin, 373
Schumann, Hans, 541
Sciambarella, I. Nina, 2519
Seco, Miquel, 119
Seddon, Kenneth R., 293, 705
Seega, Jürgen, 795
Segal, Michael G., 2485
Seigneurin, Aline, 2111
Senior, John B., 741
Seno, Manabu, 2489
Seseke, Ulrich, 847
Shah, S. Sadiq, 2881
Shaikjee, M. Salim, 1349
Shariffudin, Rio S., 747
Sharma, Anuradha, 1253
Sharma, Vijay, 1253
Sharp, Christopher, 2293
Shaw, Bernard L., 151, 221, 769, 777, 1489, 1495, 1501, 1509, 1763, 2005, 2171, 2401, 2751, 3067, 3079
Shaw, Leylâ S., (née Gözen), 925
Shaw, Robert A., 925, 2023, 2479
Sheldrick, George M., 847
Shepard, William B., 2555
Sherwood, Paul, 51, 1201, 1209, 1215, 2201, 2211
Shiina, Muneki, 853
Shimizu, Ryuichi, 1793
Shimoji, Toshimi, 1305
Shirai, Hirofusa, 1801
Shirai, Tsuneo, 2455
Shiro, Motoo, 1953
Short, Richard L., 2059, 2067, 2473, 2981
Shortman, Caroline, 1115
Shying, Michael E., 2149
Siegfried, Liselotte C., 2631
Sík, Vladimir, 757
Simmons, Mark S., 41
Simon, Arndt, 431
Sinclair, Geoffrey R., 1755
Sindellari, Livia, 1823
Singh, Anirudh, 1463
Singh, Langonjam Kanhai, 2089
Sinn, Ekkehard, 2621
Sironi, Angelo, 327, 1149
Skapski, Andrzej C., 991
Skelton, Brian W., 1477, 1817, 2843
Small, Ronald W. H., 915
Smith, Anthony K., 277
Smith, Arnold J., 2459
Smith, J. David, 747
Smith, Peter J., 129
Smith, Peter W., 1817
Smith, Thomas D., 1755, 2341
Smith, Vedene H., 2857
Smolander, Kimmo, 781
Snaith, Ronald, 1071, 2141
Sóvágó, Imre, 1, 1717
Sola, Joan, 2391
Sola, Marcó, 1333
Solans, Joaquim, 2391
Solans, Xavier, 1619, 2391, 2867
Sordo, José, 683
Soum, Alain, 1041
Sowerby, D. Bryan, 145, 1257
Spalding, Trevor R., 699
Spek, Anthony L., 1359
Spencer, Catriona M., 1963, 2649, 2709
Spencer, John L., 259
Spreer, Larry O., 2105
Stanghellini, Pier Luigi, 685, 1707
Stavropoulos, Pericles, 169, 557
Steel, Andrew T., 2359
Stella, Sergio, 1789
Stenkamp, Ronald E., 2905
Stephenson, G. Richard, 885, 2013
Stirling, Janet S., 1403
Stobart, Stephen R., 969
Stollmaier, Friederike, 431
Stone, F. Gordon A., 41, 51, 61, 73, 81, 1191, 1201, 1209, 1215, 1221, 1229, 1235, 1243, 2177, 2189, 2201, 2211, 2219, 2661, 2917
Strähle, Joachim, 683

Stranger, Robert, 1817
Straughan, Brian P., 97
Suárez, Marta, 2045
Suber, Lorenza, 1035
Sugasaka, Kazuhiko, 2489
Sullivan, Alice C., 747
Sundberg, Peter, 3103
Sunley, Glenn J., 2459
Suresh, Bettadapura S., 1123
Sykes, A. Geoffrey, 1397, 2293, 2573
Sylva, Ronald N., 2149
Symons, Martyn C. R., 421, 729
Szary, Anne C., 741
Szeto, Wing Tsee Alice, 1271
Tahir, Tahir A., 985
Tait, C. Drew, 2467
Takahashi, Katsuo, 1833
Takashima, Yoshimasa, 627, 1341
Takizawa, Takeo, 853
Tan, Chee-Kai, 1299
Tanaka, Toshio, 1793
Tandon, Santokh S., 965
Tani, Yuichiroh, 1917
Tarasconi, Pieralberto, 227
Tarli, Franco, 1035
Tasker, Peter A., 1631, 2537
Tatehata, Akira, 1977
Tattershall, Bruce W., 1515
Taylor, Brian F., 1947, 1963, 2709
Taylor, John G., 1805, 3107
Taylor, Nicholas J., 1463
Tegen, Marvin, 2599
Terheijden, Jos, 1359
Thatcher, Peter, 2615
Thomas, Dianne M., 2803
Thompson, Arthur R., 2317
Thompson, James Charlton, 1123
Thornton-Pett, Mark, 275, 451, 769, 1169, 1431, 1489, 1501, 1509, 1763, 2171, 2417, 2751, 2781, 3079.
Tirant, Michael, 1755, 2341
Tiripicchio, Antonio, 953, 1275, 1551
Tisato, Francesco, 867, 1693
Tobe, Martin L., 1775
Toerien, James G., 1349
Tolazzi, Marilena, 657
Toma, Hani J., 2531
Tooze, Robert P., 557
Tosi, Gabriella, 227
Tossing, Gudrun, 795
Tourné, Claude M., 1283
Tourné, Gilbert F., 1283
Townsend, Rodney P., 1911
Tran-Huy, N. Hoa, 1229, 1235, 1243, 2201
Traverso, Orazio, 1659
Treichel, Paul M., 2599
Trincia, Lucio, 1823
Trinquier, Georges, 2373
Trogu, Emanuele F., 2407
Tsuboyama, Kaoru, 721, 3093
Tsuboyama, Sei, 721, 3093
Tsuchida, Eishun, 1917, 2455, 2493, 2963
Tüzün, Mürşide, 925
Turner, David G., 259
Turp, Janet E., 293
Turp, Neil, 1163
Twiss, Paul, 2347
Uchida, Hiroaki, 1377
Underhill, Allan E., 1683
Ungaretti, Luciano, 2605
Urch, David S., 1727
Usón, Rafael, 2077, 3017
Valle, Giovanni, 815
van Bekkum, Herman, 2051
van Duin, Martin, 2051
van Koten, Gerard, 1359
van Rijn, Jacobus, 2579
Varnagy, Katalin, 1
Vasapollo, Giuseppe, 2165
Vasudevan, Asha K., 427
Vefghi, Rahmatollah, 969, 2815
Vegas, Angel, 183
Veitch, Paul M., 2853
Venturini, Marilyne, 1145
Vess, Thomas M., 2467
Vettori, Umberto, 815
Vicente, José, 881, 1655
Vila, José M., 2401
Villarroya, B. Eva, 981
Villeneuve, Gerard, 285
Vitali, Francesca, 177
Vos, Johannes G., 1389, 3073
Vrieze, Kees, 1359
Waghorne, W. Earle, 2531
Wade, Kenneth, 1071, 2141
Wahlgren, Curtis G., 2621
Walford, Alison M., 1865
Walker, Fiona J., 705
Walker, Nigel P. C., 1861, 1869
Walkinshaw, Malcolm D., 373, 2831
Wallbridge, Malcolm G. H., 1805, 2793, 3107
Wallis, Julian M., 2823
Walton, Jacqueline R., 1535
Walton, Richard A., 715
Wardle, Robert W. M., 1889
Warrens, Christopher P., 907
Watanabe, Yoshihisa, 2969
Watkins, David A., 2479
Watkinson, Philip J., 2365
Watson, Rhuann R., 391
Weakley, Timothy J. R., 1283, 1545, 2789
Webb, Thomas R., 1299
Webster, Michael, 1901
Wei, Chen, 129
Weiss, Johannes, 187
Weissensteiner, Walter, 1545
Welch, Alan J., 997, 1017, 2761, 3035
Wernberg, Ole, 293
Wertz, Dennis W., 2467
Whitaker, Claire R., 1647
White, Alex, 2885
White, Allan H., 457, 1089, 1099, 1463, 1477, 1647, 1737, 1817, 2347, 3085
White, Colin, 1149, 1947, 2815
White, Peter S., 1989
Whiteley, Mark W., 3049
Whitwell, Ian, 1181
Wickens, Denys A., 1581
Wieghardt, Karl, 187
Wild, David J., 267
Wilkins, Alistair L., 2365
Wilkins, Cuthbert J., 523
Wilkinson, Geoffrey, 169, 557, 837, 2723, 2947
Willey, Gerald R., 263
Williams, David J., 907
Williams, Geoffrey A., 1585, 2305
Williams, Ian D., 591, 1313, 1319
Wilson, Jenny, 1585
Winter, Mark J., 341, 479
Winter, Michael J., 1235
Winterborn, Duncan J. W., 1463
Wong, Chi-Ming, 1989
Wong, Luet-Lok, 411, 2031
Wood, David, 3099
Wood, G. Leslie, 1485
Woollins, J. Derek, 275, 907
Wyatt, Robert S., 859
Wynd, Andrew J., 2761
Yamamoto, Yasuhiro, 851, 853, 1833
Yamase, Toshihiro, 1597
Yamazaki, Hiroshi, 851, 853, 1009, 1833, 2017
Yates, Paul C., 123, 2563
Yavari, Parvin, 1947
Yhome, Khriesavilie, 1055
Yichang, Wu, 1501
Yoshida, Shin-ichi, 2969
Yoshikawa, Sadao, 1127
Young, Charles G., 619
Yuasa, Makoto, 1917, 2455, 2493
Zanazzi, Pier Francesco, 473, 473
Zanella, Pierino, 691, 2039
Zanello, Piero, 831
Zangen, Mendel, 235
Zanonato, Pier Luigi, 657, 695
Zanotti, Valerio, 1133
Zao, Xizhang, 1801
Zaworotko, Michael J., 969
Zehnder, Margareta, 709
Zerga, Hugo O., 1267
Zuber, Gary E., 677
Zubieta, Jon A., 529

SUBJECT INDEX, 1987

ACETATE
Cyclopalladation to form planar tridentate [N–C–N]$^-$ intramolecular co-ordination systems involving pyridine donor groups, including ligand synthesis and *X*-ray structural studies., 1477–84
Substitution reactions of the osmium(I) dinuclear compound $[Os_2(MeCO_2)_2(CO)_6]$. *X*-Ray crystal structure of $[N(PPh_3)_2][Os_2(MeCO_2)_2Cl(CO)_5]$, 2473–8

ACETIMIDIC ACID
Preparation, nuclear magnetic resonance spectra, and *X*-ray diffraction study of bis(*N*-benzylacetimidic acid ethyl ester)dichloropalladium(II), 2503–4

ACETOIN
Photo-oxidations of acetoin and biacetyl catalysed by tetrabutylammonium decatungstate(4−) in acetonitrile under excess of oxygen, 961–2

ACETONITRILE
A reversible metal framework rearrangement in high-nuclearity osmium-platinum cluster compounds. *X*-Ray crystal structures of $[Os_6Pt(CO)_{17}(\mu_3\text{-NCMe})(C_8H_{12})]$, $[Os_6Pt(CO)_{17}(\mu_4\text{-NCMe})(C_8H_{12})]$ (C_8H_{12} = cyclo-octa-1,5-diene), and $[Os_6(CO)_{19}(\mu_3\text{-NCMe})]$: compounds with novel metal–acetonitrile bonding modes, 2245–52
Electrolytic behaviour of transition metal bromides in acetonitrile solution, 2863–6

ACETYL
Conversion of uncharged methyl complexes of ruthenium(II) into a cationic acyl complex: a mechanistic study, 1713–6
Reactivity of a cluster-co-ordinated acyl. Reduction of a μ-acetyl ligand to acetaldehyde, 2839–42

ACETYLIDE
Bimetallic systems. Part 18. Nickel(II)–mercury(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$ ligands, 777–80

ACETYLPYRIDINE
Complexes of cobalt(II) and nickel(II) with 3- and 4-acetylpyridine, 2595–8

ACTINIDE
Reaction of monoborane with tris(cyclopentadienyl)uranium(IV) derivatives, 691–4
Uranyl(VI) complexes of 3-aminopropanoic acid and 4-aminobutanoic acid in aqueous solution: a potentiometric and calorimetric study, 695–8
Insertion of 2,6-dimethylphenyl isocyanide into the U–N bond of $[U(\eta\text{-}C_5H_5)_3(NEt_2)]$. Crystal structure of $[U(\eta\text{-}C_5H_5)_3\{C(NEt_2)=NC_6H_3Me_2\text{-}2,6\}]$, 2039–44
Mixed-metal phosphinito complexes of platinum(II) and palladium(II) with lanthanide and actinide elements. The single-crystal *X*-ray structure determination of $[UO_2(OH_2)\{(OPPh_2)_2Pd(S_2CNEt_2)\}_2]$, 2853–6

ADENOSINE
The configuration and lattice dynamics of complexes of dialkyltin(IV) with adenosine 5′-monophosphate and phenyl phosphates, 789–94

ADSORPTION
Infrared spectroscopic studies of uranyl(VI) species adsorbed from aqueous $[UO_2(CO_3)_3]^{4-}$ solutions on to a polymer bearing amidoxime groups, 2489–92

ALANINATE
Amino-acid complexation to a cobalt(III) complex of optically-active (2*R*,5*R*,8*R*,11*R*)-2,5,8,11-tetraethyl-1,4,7,10-tetra-azacyclododecane (L). Preparation, characterization, and structure determinations of $[Co^{III}\{(R)\text{- or }(S)\text{-alaO}\}L]Br\text{-}[ClO_4]\cdot H_2O$, 721–8

ALCOHOL
Synthesis of the alkoxo(hydrido)-clusters $[Ru_3(\mu\text{-H})(CO)_{10}(\mu\text{-OR})]$ [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4

ALKALI METAL
A quantitative approach to host–guest interactions for matrix-isolated alkali-metal salts of hexafluoroanions and perchlorates, 1521–8
Molten lithium carbonate–sodium carbonate–potassium carbonate eutectic: electronic spectroscopy of first-row transition metal compounds, 2329–32

ALKANE
Activation of alkanes involving rhenium atoms: synthesis and electronic structure of binuclear μ-alkylidene derivatives; molecular structure of $[\{Re(\eta\text{-}C_6H_6)\}_2(\mu\text{-}CHBu^t)(\mu\text{-H})_2]$, 391–402

ALKANYLIDYNE
Reactions of co-ordinated ligands. Part 39. The synthesis of carbyne complexes from alkyne molybdenum and tungsten cations; formation and crystal structures of $[Mo(\equiv CCH_2Bu^t)\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$ and $[Mo\{=C(SiMe_3)CH_2\}\{P(OMe)_3\}_2(\eta^5\text{-}C_9H_7)]$, 591–604

ALKENE
trans to *cis* isomerisations of platinum(II) olefin phosphine complexes, $[PtX_2(ol)(PR_3)]$ (X = Cl or Br; ol = ethene, propene, or 1-heptene; PR_3 = PBu_3 or PMe_2Ph), 139–44
Co-ordination of highly substituted alkenes to transition-metal complexes: preparation and crystal and molecular structures of the complexes $[\{RhCl(Ph_2PO_2CCH{=}CMe_2)\}_2]$ and $[\{RhCl(Ph_2PO_2CCH{=}CHCH{=}CHMe)\}_2]$, 443–50
Preparation, isomerization, and reactions of hydride complexes of ruthenium(II), 2989–94

ALKENYL
Crystal and molecular structures of $[MCl(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf) and $[TiCl_{0.5}Br_{0.5}(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$: an unusual π-interaction with the metals and molecular orbital calculations on model compounds, 1641–6

ALKOXIDE
Synthesis of the alkoxo(hydrido)-clusters $[Ru_3(\mu\text{-H})(CO)_{10}(\mu\text{-OR})]$ [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4
Synthesis, characterization, and reactions of chromyl(VI) alkoxides, 1253–6
Synthesis of chelate stabilised alcohol- and alkoxo–platinum(II) complexes. *X*-Ray crystal structure of $[Pt(PPh_2CH_2CMe_2O)_2]\cdot 3.5H_2O$, 2273–80

ALKYL
Organotin biocides. Part 8. The crystal structure of triphenyltin formate and a comparative variable-temperature tin-119 Mössbauer spectroscopic study of organotin formates and acetates, 101–6
The synthesis and structure of Group 5 metal alkyl and alkylidene complexes containing 2,6-dialkylphenoxide ligands: *X*-ray crystal structures of $[Ta(OC_6H_3Me_2\text{-}2,6)_2(CH_2Ph)_3]$, $[Ta(OC_6H_3Me_2\text{-}2,6)_4Me]$, and $[Ta(OC_6H_3Bu^t_2\text{-}2,6)_2(=CHSiMe_3)(CH_2SiMe_3)]$, 155–62
Electronic absorption spectra and photochemical reactivity of Group 5 metal alkyl compounds: photochemical α-hydrogen abstraction, 163–8
The configuration and lattice dynamics of complexes of dialkyltin(IV) with adenosine 5′-monophosphate and phenyl phosphates, 789–94
Syntheses and crystal structures of complexes $[M_2R_2]$, [M = Cu, Ag, or Au; R = 2-C(SiMe$_3$)$_2$C$_5$H$_4$N] and $[Cu_4R'_4]$[R′ = 2-CH(SiMe$_3$)C$_5$H$_4$N]; electrochemical generation of $[Cu_2R_2]^{2+}$, 3085–92

ALKYLIDENE
The synthesis and structure of Group 5 metal alkyl and alkylidene complexes containing 2,6-dialkylphenoxide ligands: *X*-ray crystal structures of $[Ta(OC_6H_3Me_2\text{-}2,6)_2(CH_2Ph)_3]$, $[Ta(OC_6H_3Me_2\text{-}2,6)_4Me]$, and $[Ta(OC_6H_3Bu^t_2\text{-}2,6)_2(=CHSiMe_3)(CH_2SiMe_3)]$, 155–62
Activation of alkanes involving rhenium atoms: synthesis and electronic structure of binuclear μ-alkylidene derivatives; molecular structure of $[\{Re(\eta\text{-}C_6H_6)\}_2(\mu\text{-}CHBu^t)(\mu\text{-H})_2]$, 391–402

ALKYLIDYNE
Chemistry of polynuclear metal complexes with bridging carbene or

ALKYLIDYNE (cont)
carbyne ligands. Part 54. Compounds derived from $[W(\equiv CC_6H_4Me\text{-}4)(CO)_2(\eta\text{-}C_5Me_5)]$ with bonds between tungsten and iron, cobalt, or nickel; crystal structure of $[FeW\{\mu\text{-}C(C_6H_4Me\text{-}4)C(O)O\}(CO)_5(\eta\text{-}C_5Me_5)]$, 1191–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 55. Coupling of methylene and alkylidyne groups at a trimetal centre; crystal structures of $[Ru_2W\{\mu\text{-}\sigma\text{:}\eta\text{-}trans\text{-}CH{=}C(H)R\}(\mu\text{-}CO)(CO)_3(\eta\text{-}C_5H_5)_2(\eta\text{-}C_5Me_5)]\cdot CH_2Cl_2$ and $[Ru_2W(\mu_3\text{-}CCH_2R)(\mu\text{-}CO)_3(CO)(\eta\text{-}C_5H_5)_3]$ ($R = C_6H_4Me\text{-}4$), 1201–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 56. Synthesis of iron–molybdenum compounds; crystal structures of $[FeMo(\mu\text{-}CR)(CO)_6(\eta\text{-}C_5H_5)]$ and $[FeMo_2(\mu_3\text{-}RC_2R)(CO)_6(\eta\text{-}C_5H_5)_2]$ ($R = C_6H_4Me\text{-}4$), 1209–14
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 57. Reactions of iron– molybdenum complexes with oxygen and sulphur; crystal structure of $[FeMo(\mu\text{-}\eta^2\text{-}SCC_6H_4Me\text{-}4)(CO)_5(\eta\text{-}C_5H_5)]$, 1215–20
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 58. Some reactions of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (M = Mo or W); crystal structures of the complexes $[W_2(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_2(L)(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$-$[L = N_2C(C_6H_4Me\text{-}4)_2$ or O], 1221–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 59. Reactions of the alkylidyne compounds $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Cr, Mo, or W, $R = C_6H_4Me\text{-}4$; M = W, R = Me, $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate] with d^{10} complexes; crystal structure of $[PtW_2(\mu\text{-}CMe)_2(CO)_4\{HB(pz)_3\}_2]$, 1229–34
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 60. Di- and tri-metal compounds prepared from the metal alkylidyne complexes $[W(\equiv CR)(CO)_2(\eta\text{-}C_5Me_5)]$ ($R = Me$ or $C_6H_4Me\text{-}4$) and $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Mo, $R = C_6H_4Me\text{-}4$; M = W, R = Me; $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate], 1235–42
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 61. Reactions of iron– molybdenum compounds with tertiary phosphines, 1243–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 62. Synthesis of penta-, hexa-, and hepta-heteronuclear metal cluster compounds involving tungsten or molybdenum with platinum or nickel: crystal structures of the chain compounds $[Pt_3W_3(\mu\text{-}CMe)(\mu_3\text{-}CMe)_2(CO)_6(cod)(\eta\text{-}C_5Me_5)_3]\cdot 0.5CH_2Cl_2$ and $[Pt_3W_4(\mu\text{-}CC_6H_4Me\text{-}4)_2(\mu_3\text{-}CC_6H_4Me\text{-}4)_2(CO)_8(\eta\text{-}C_5H_5)_4]$, 2177–88
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 63. Synthesis of eight-membered-ring metallacycles: *X*-ray crystal structures of $[Pt_4W_4(\mu\text{-}CR)(\mu_3\text{-}CR)_3(\mu\text{-}CO)(CO)_7(\eta\text{-}C_5H_5)_4]\cdot CH_2Cl_2$, $[Ni_2Pt_2W_4(\mu\text{-}CR)(\mu_3\text{-}CR)_3(\mu\text{-}CO)(CO)_7(\eta\text{-}C_5H_5)_4]\cdot CH_2Cl_2$ ($R = C_6H_4Me\text{-}4$), and $[Ni_2Pt_2W_4(\mu_3\text{-}CPh)_4(CO)_8(\eta\text{-}C_5H_5)_4]$, 2189–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 64. Addition of methylene groups to iron–molybdenum complexes; crystal structures of $[FeMo(\mu\text{-}CH_2)\{\mu\text{-}\sigma\text{:}\eta\text{-}C(C_6H_4Me\text{-}4){=}CH_2\}(CO)_5(\eta\text{-}C_5H_5)]$ and $[FeMo\{\mu\text{-}C(C_6H_4Me\text{-}4)C(OMe)C(H)\}(CO)_5(\eta\text{-}C_5H_5)]$, 2201–10
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 65. Reactions of the complexes $[FeW(\mu\text{-}CR)(CO)_5(\eta\text{-}C_5Me_5)]$ and $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5Me_5)]$ ($R = C_6H_4Me\text{-}4$) with alkynes R′C$_2$R′ (R′ = Me or Ph); crystal structures of $[FeW\{\mu\text{-}C(R)C(O)C(Me)C(Me)\}(CO)_5(\eta\text{-}C_5Me_5)]\cdot CH_2Cl_2$ and $[FeW\{\mu\text{-}C(R)C(Et)C(H)C(Me)C(Me)\}(\mu\text{-}CO)(CO)_3(\eta\text{-}C_5Me_5)]$, 2211–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 66. Carbaboranetungsten– platinum complexes. Polyhedral rearrangements of a 12-vertex cage system; crystal structures of $[PtW(CO)_2(PEt_3)_2\{\eta^6\text{-}C_2B_9H_8\text{-}(CH_2C_6H_4Me\text{-}4)Me_2\}]\cdot CH_2Cl_2$, $[PtW(\mu\text{-}H)\{\mu\text{-}\sigma\text{:}\eta^5\text{-}C_2B_9H_7(CH_2C_6H_4Me\text{-}4)Me_2\}(CO)_2(PMe_3)(PEt_3)_2]$, and related compounds, 2219–34
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 67. Reactions of the salts $[X][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$[X = $P(CH_2Ph)Ph_3$, $R = C_6H_4Me\text{-}4$; X = NEt_4, R = Ph] with bis(cyclo-octa-1,5-diene)platinum, 2661–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 68. Reactions between nonacarbonyldi-iron and the salts $[NEt_4][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = Me, Ph, or $C_6H_4Me\text{-}4$); crystal structures of $[NEt_4][FeW\{\mu\text{-}CH(C_6H_4Me\text{-}4)\}(\mu\text{-}\sigma\text{:}\eta^5\text{-}C_2B_9H_8Me_2)(\mu\text{-}CO)(CO)_5]$ and $[NEt_4][FeW(\mu_3\text{-}CPh)(\mu\text{-}\sigma\text{:}\sigma',\eta^5\text{-}C_2B_9H_7Me_2)(CO)_8]$, 2917–26

ALKYNE
Thermal and photochemical behaviour of organotetraruthenium clusters: solution structures and dynamics of phosphine-substituted derivatives, 349–54
Reactions of co-ordinated ligands. Part 39. The synthesis of carbyne complexes from alkyne molybdenum and tungsten cations; formation and crystal structures of $[Mo(\equiv CCH_2Bu^t)\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$ and $[Mo\{{=}C(SiMe_3)CH_2\}\{P(OMe)_3\}_2(\eta^5\text{-}C_9H_7)]$, 591–604
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η^2-vinyl complexes of tungsten, $[W\{\eta^3\text{-}C(CF_3)C(CF_3)(SR)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (R = Me, Et, Pr^n, Pr^i, or Bu^t), 895–906
New dicyclopentadienyl phosphine, phosphite, and acetylene niobium(III) complexes. Crystal structure of $[Nb(\eta\text{-}C_5H_4SiMe_3)_2Cl(PhC{\equiv}CPh)]$, 975–80
Carbene complexes in organic synthesis: a new tungstacarbene-promoted alkyne insertion–olefin cyclopropanation reaction, 1041–50
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 56. Synthesis of iron–molybdenum compounds; crystal structures of $[FeMo(\mu\text{-}CR)(CO)_6(\eta\text{-}C_5H_5)]$ and $[FeMo_2(\mu_3\text{-}RC_2R)(CO)_6(\eta\text{-}C_5H_5)_2]$ ($R = C_6H_4Me\text{-}4$), 1209–14
Reactions of co-ordinated ligands. Part 41. Synthesis and reactions of cationic functionalised alkyne molybdenum complexes; formation of carbyne and vinylidene complexes and crystal structure of $[Mo(C{=}CHPh)Br\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$, 1319–28
Synthesis and reactions of the co-ordinatively unsaturated hexafluorobut-2-yne complex $[WBr_2(CO)(CF_3C{\equiv}CCF_3)_2]$. The crystal and molecular structure of $[\{W[C(CF_3){=}C(CF_3)P(OEt)_2O](\mu\text{-}Br)(CO)(CF_3C{\equiv}CCF_3)\}_2]$, 2165–70

ALKYNYL
Some chemistry of the heterobimetallic complexes $[(OC)_3Mo(\mu\text{-}dppm)_2Rh(CO)Cl]$ and $[(OC)_3Mo(\mu\text{-}dppm)_2\text{-}M(CO)_2]PF_6$ (M = Rh or Ir, dppm = $Ph_2PCH_2PPh_2$). Structure of $[(OC)_3Mo(\mu\text{-}dppm)_2(\mu\text{-}C{\equiv}CMe)Rh(CO)]$, 2171–6
σ-Alkynyl complexes of manganese(I) as η^2-bonding ligands for Group 1B metal–ligand fragments. *X*-Ray crystal structure of $[Mn_2Cu(\mu\text{-}CCBu^t)_2(CO)_6(dppe)_2]PF_6\cdot 0.5CH_2Cl_2$, 2867–74
The chemistry of cycloheptatrienyl complexes of molybdenum and tungsten: the synthesis and reactions of some vinylidene and alkynyl derivatives, 3049–54

ALLENE
Reactions of allene with triosmium and triruthenium clusters including coupling and oxidative addition: *X*-ray structure of the coupled allene compound $[Os_3(C_6H_8)(CO)_{10}]$, 2935–40

ALLYL
Synthesis and characterization of palladium(II)–η^3-allyl–ylide complexes. *X*-Ray crystal structure of $[PdCl(\eta^3\text{-}2\text{-}MeC_3H_4)\{Ph_3PC(H)COMe\}]$, 1381–8

ALLYLIDENE
Reactions of co-ordinated ligands. Part 40. Synthesis, structure, and reactivity of dirhenium μ-allylidene complexes; crystal structures of $[Re_2(\mu\text{-}\sigma\text{:}\eta^3\text{-}CHCHCMe_2)(CO)_8]$ and $[Re_2(\mu\text{-}\sigma\text{:}\eta^3\text{-}CHCHCMe_2)(CO)_9]$, 1313–8

ALUMINIUM
Molecular orbital study of the bridging dioxygen ligand in $[Al_2(CH_3)_6(\cup_2)]^-$, 355–60
Reactions of aluminopolyhydride complexes of tungsten. *X*-Ray crystal structures of $[(Me_3P)_3H_3W(\mu\text{-}H)_2AlCl(\mu\text{-}N{=}CHEt)]_2$, $\{[(Me_3P)_3WH_5]Li\}_4$, $WH_2I(PMe_3)_3(SiMe_3)$, and $(Me_3P)_3H_2W(\mu\text{-}H)_3AlCl_2(NMe_3)$, 837–46
Aluminium speciation studies in biological fluids. A new investigation of aluminium hydroxide equilibria under physiological conditions, 1145–8
Oxygen-17 and aluminium-27 nuclear magnetic resonance spectroscopic investigations of aluminium(III) hydrolysis products, 2317–22

ALUMINIUM-27
Oxygen-17 and aluminium-27 nuclear magnetic resonance spectroscopic investigations of aluminium(III) hydrolysis products, 2317–22

AMIDE
Reactions of aluminopolyhydride complexes of tungsten. *X*-Ray crystal structures of $[(Me_3P)_3H_3W(\mu\text{-}H)_2AlCl(\mu\text{-}N{=}CHEt)]_2$, $\{[(Me_3P)_3WH_5]Li\}_4$, $WH_2I(PMe_3)_3(SiMe_3)$, and $(Me_3P)_3H_2W(\mu\text{-}H)_3AlCl_2(NMe_3)$, 837–46
Reactions of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}X_2]$ (M = Mo, X = I; M = W, X = Cl) with some amine-substituted pyridines. The molecular structures of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}(2\text{-}NHC_5NH_4)_2]$, 1063–70
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1283–4
Systematic cluster build up by the reaction of $[Os_3H(X)(CO)_9(MeCN)]$ (X = SR or NHMe; R = Me, Et, or Ph) with $[OsH_2(CO)_4]$, 1289–92
Substitution reactions on triosmium clusters of the type $[Os_3(\mu\text{-}H)(\mu\text{-}X)(CO)_{10}]$ [X = NHMe, O_2CR, C_5H_4N, or OC(NHMe); R = H, Me, or Ph], 1293–8
Reduction–oxidation properties of organotransition-metal complexes. Part 26. The synthesis of paramagnetic triazenido-and aryl(1-aryliminoethyl)amido-bridged dirhodium carbonyl derivatives containing the $[Rh_2]^{4+}$ core, 1403–8
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1583–4
Tetranuclear amido complexes of copper(I): a synthetic and structural study, 1883–8
Insertion of 2,6-dimethylphenyl isocyanide into the U–N bond of $[U(\eta\text{-}C_5H_5)_3(NEt_2)]$. Crystal structure of $[U(\eta\text{-}C_5H_5)_3\{C(NEt_2){=}NC_6H_3Me_2\text{-}2,6\}]$, 2039–44
Methylindium dialkylamido compounds: mono- and bis-dialkylamides involving four- and five-co-ordinated indium(III). *X*-Ray crystal structures of $\{Me_2In[N(CH_2)_2N(Me)CH_2CH_2]\}_2$, $MeIn[MeNC(CH)_4N]_2$, and $MeIn\{[MeN(CH_2)_2NMe]\text{-}InMe_2\}_2$, 2159–64
Phosphorus–nitrogen compounds. Part 51. Carbon-13 and phosphorus-31 nuclear magnetic resonance properties of phenylphosphonothioic di(monoalkylamides). Determination of aromatic substituent constants by ^{13}C nuclear magnetic resonance spectroscopy, 2479–84

AMIDINO
Synthesis and study of monomeric and dimeric amidine complexes of nickel(II) and platinum(II). Crystal structure of bis(*N*,*N*′-diphenylbenzamidino)platinum(II), $[Pt\{PhNC(Ph)NPh\}_2]$, 2687–94

AMIDOXIME
Infrared spectroscopic studies of uranyl(VI) species adsorbed from aqueous $[UO_2(CO_3)_3]^{4-}$ solutions on to a polymer bearing amidoxime groups, 2489–92

AMINE
Equilibria in aqueous solutions of some iron(II) complexes, 255–8
Studies of phosphazenes. Part 30. Reactions of hexachlorocyclotriphosphazene with aromatic primary amines: interplay of geminal and non-geminal modes of chlorine replacement, 579–84
Studies of phosphazenes. Part 31. Kinetic studies of the reactions of hexachlorocyclotriphosphazene with aromatic primary amines: evidence for the intermediacy of a reactive three-co-ordinated phosphorus(V) species, 585–90
Thermodynamics of complex formation of silver with amines in dimethyl sulphoxide, 657–60
Electron diffraction study of the molecular structure of *N*-methylbis(trimethylsilyl)amine in the gas phase, 785–8
Phosphorus–nitrogen compounds. Part 50. Further studies on the formation of bicyclic cyclotetraphosphazatetraene derivatives, 2023–30
New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states, 2621–30
Complexes formed between zinc(II) and diethylenetriamine: a carbon-13 nuclear magnetic resonance study, 2889–96
Carbon-13 nuclear magnetic resonance study of the zinc(II) complexes of 3,7-diazanonane-1,9-diamine and 1,2-diaminoethane, 2897–904

AMINO ACID
Co-ordination abilities of thioamide analogues of amino acids and peptides. Copper(II) and nickel(II) complexes, 1–4
Magnetic and electrochemical studies on the series $[Cr_nFe_{3-n}(\mu_3\text{-}O)(gly)_6(H_2O)_3][NO_3]_{7-n}\cdot xH_2O$(gly = glycine; *n* = 0, 1, 2, or 3) and on the mixed-valence complex $[Fe^{II}Fe^{III}_2(\mu_3\text{-}O)(gly)_6(H_2O)_3]Cl_6$, 97–100
Electron spin resonance and electronic spectra and crystal and molecular structures of copper(II) amino acid complexes, 457–66
A thermodynamic and spectroscopic study of the proton and copper(II) complexes of L-prolyl-L-histidine, D-prolyl-L-histidine, L-histidyl-L-histidine, and D-histidyl-L-histidine, 661–6
Uranyl(VI) complexes of 3-aminopropanoic acid and 4-aminobutanoic acid in aqueous solution: a potentiometric and calorimetric study, 695–8
Amino-acid complexation to a cobalt(III) complex of optically-active (2*R*,5*R*,8*R*,11*R*)-2,5,8,11-tetraethyl-1,4,7,10-tetra-azacyclododecane (L). Preparation, characterization, and structure determinations of $[Co^{III}\{(R)\text{- or }(S)\text{-alaO}\}L]Br\text{-}[ClO_4]\cdot H_2O$, 721–8
Copper(II) complexes of tyrosine-containing tripeptides, 1263–6
Spectroscopic, magnetic, and structural properties of diaquadimethanolbis(*N*-tosylvalinato)copper(II) and its 2,2′-bipyridine adduct, 1333–40
Studies on transition-metal–peptide complexes. Part 13. Copper(II) and nickel(II) complexes of amino acids and peptides containing a thioether group, 1717–20
Metal ion-bridged intramolecular stacking interaction between the tryptophyl residue and the aromatic heterocyclic amine within the ternary complex (1,10-phenanthroline)(L-tryptophanato-*O*,*N*)copper(II) perchlorate–water (1/2.5): an *X*-ray study, 2017–22
Synthesis, characterisation, and aquation kinetics of [*N*-(*o*-carboxyphenyl)iminodiacetato](glycinato)chromate(III) complex ion in acidic media, 2095–8

AMINOMETHYLIDYNE
Triosmium clusters containing the ligands CNH_2, NCHCN, and NCMeCN derived from cyanide ion, cyanogen, and acetonitrile, 1457–62

AMINOSILANE
Preparations, properties, and vibrational spectra of some (dimethylamino)halogenosilanes, 3029–34
Molecular structures of some (dimethylamino)halogenosilanes in the gas phase by electron diffraction and the crystal and molecular structures of mono-and di-chloro(dimethylamino)-silane by *X*-ray diffraction at low temperatures, 3035–42
The structure of bis(dimethylamino)silane in the gas phase, determined by electron diffraction, 3061–6

AMMONIA
Spectrochemistry of solutions. Part 19. Complexation of gold(I) by cyanide in liquid ammonia: an infrared and Raman spectroscopic study, 673–6

ANTIMONY
The crystal structures of the diphenyldithiophosphinates of antimony(III) and bismuth(III): $M(S_2PPh_2)_3$ (M = Sb or Bi), 145–50
Crystal and molecular structure of the six-co-ordinate antimony(III) complex $SbCl_3L$, where L = *N*,*N*′-dimethyloxamide. Unusual O,O ligand binding leading to four-membered $Sb(\mu\text{-}O)_2Sb$ ring formation, 263–6
Stereochemical role of lone pairs in main-group elements. Part 3. Structure and bonding in trichloro(1,4,7,10,13-pentaoxacyclopentadecane)antimony(III) studied by means of *X*-ray crystallography at 120 K, 427–30
Bonding in and gas-phase pyrolysis of the arsoranes and stibanes EMe_3X_2 (E = As or Sb, X = F or Cl) and $SbMe_4F$: ultraviolet photoelectron and field-ionization mass spectrometric studies, 481–8
Complexes of 2,5,7,9-tetrathiabicyclo[4.3.0]non-1(6)-ene-8-thione (L^1). Part 1. Synthesis and antimony-121 Mössbauer spectroscopic characterisation of $SbCl_3(L^1)_{1.5}$, crystal and molecular structures of $SbCl_3(L^1)_{1.5}$ and uncomplexed L^1, 1605–10
Iron antimonate, 1727–32
The preparation and crystal structure of $5SbF_3\cdot 3SbF_5$, 1989–98

ANTIMONY (contd)
The reaction of di-2-pyridyl ketone with antimony(III) fluoride. Preparation, properties, Mössbauer spectrum and crystal structure of difluoro(methoxydi-2-pyridylmethoxo-*NN'O*)antimony(III), 2337–40
Metal(II) hexafluoroantimonates: preparation and characterization of $MF_2{\cdot}2SbF_5$ (M = Mg, Ni, Zn, Fe, Co, Cu, Cr, Ag, Cd, or Pb) and the *X*-ray structure determination of $AgF_2{\cdot}2SbF_5$, 2379–84
Preparation, crystal and molecular structure, and antimony-121 Mössbauer spectrum of trifluoro(1,10-phenanthroline)(thiourea)-antimony(III), 2451–4
AQUATION
Synthesis, characterisation, and aquation kinetics of [*N*-(*o*-carboxyphenyl)iminodiacetato](glycinato)chromate(III) complex ion in acidic media, 2095–8
ARENE
Synthesis of η-arene rhenium compounds using rhenium atoms. Crystal structures of $[Re_2(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)_2(\mu\text{-}H)_2]$ and $[Re(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)H]$, 403–10
Isometallocenes. Part 1. Polymethylated isocobaltocenium salts; a general route to $[Co(C_4Me_4)(arene)]$, 2757–60
Synthesis using metal atoms of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[OsH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[MnH(\eta\text{-}C_6H_6)\text{-}(PMe_3)_2]$, and $[Mn(\eta\text{-}C_5H_5)(PMe_3)_3]$: arene ligand-exchange reactions of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, 2823–30
ARENEDIAZO
Chelate ring-opening of rhenium–diphosphine complexes by diazonium salts, 3067–72
ARSENIC
Oxidation and reduction of phosphorus–phosphorus and arsenic–arsenic double bonds. An electrochemical study of two diphosphenes and a diarsene, 249–52
Bonding in and gas-phase pyrolysis of the arsoranes and stibanes EMe_3X_2 (E = As or Sb, X = F or Cl) and $SbMe_4F$: ultraviolet photoelectron and field-ionization mass spectrometric studies, 481–8
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid, 573–8
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid (1987, 573), 2291–2
Mean phosphorus– and arsenic–sulphur bond enthalpies in diethyldithiocarbamate chelate complexes, 2955–8
ARSINE
Sulphur–arsenic ligands. Part 1. Dimethyl(methylthio)arsine complexes of halogenotrimethylplatinum(IV) and tricarbonylhalogenorhenium(I), 417–20
Reactions of phosphine, arsine, and stibine with carbonylbis(triethylphosphine)iridium(I) halides. Part 1. Reactions in toluene; *X*-ray crystal structures of $[Ir(CO)ClH(PEt_3)_2(AsH_2)]$ and $[Ir(CO)XH(PEt_3)_2(\mu\text{-}ZH_2)RuCl_2(\eta^6\text{-}MeC_6H_4CHMe_2\text{-}p)]$ (X = Br, Z = P; X = Cl, Z = As), 2831–8
ARYL
Organotin biocides. Part 8. The crystal structure of triphenyltin formate and a comparative variable-temperature tin-119 Mössbauer spectroscopic study of organotin formates and acetates, 101–6
Oxoaryls of rhenium-(V) and -(VI) and osmium(VI). *X*-Ray crystal structures of dimesityldioxorhenium(VI), tetramesityloxorhenium(VI) and dimesityldioxo-osmium(VI), 169–76
Secondary bonding. Part 14. Structural isomerism in diaryliodonium halides and the structure of di(*p*-tolyl)iodonium bromide, 193–6
The synthesis and *X*-ray crystal structures of homoleptic tetrahedral aryls of osmium(IV) and of cyclohexyls of ruthenium(IV), osmium(IV), and chromium(IV), 557–62
Neutral and anionic 2-methyl-6-nitrophenyl palladium(II) complexes. The *X*-ray crystal structure of bis(benzyltriphenylphosphonium) *trans*-di-μ-chloro-bis[chloro(2-methyl-6-nitrophenyl)palladate(II)]–acetone (1/2), 1655–8
ARYLIMIDE
Chemistry of molybdenum. Part 5. Synthesis, spectroscopic and electrochemical properties of bis(2-aminocyclopent-1-ene-1-dithiocarboxylato)(arylimido)dihalogenomolybdenum(VI) complexes, 3043–8
ASCORBIC ACID
Crystal structure of a copper complex of 2-carboxypentonic acid; a decomposition product of dehydroascorbic acid, 2905–10
AZATHIENYL
Rational synthesis of metalla–sulphur–nitrogen complexes: use of $[Me_2SnS_2N_2]_2$ in the preparation of complexes of the type $[M(S_2N_2H)(PR_3)_2]X$ (M = Pt or Pd). *X*-Ray structures of $[Pt(S_2N_2H)(PEt_3)_2][Me_2SnCl_3]$, $[Pt(S_2N_2H)(PBu^n_3)_2]_2[PF_6]Cl$, and $[\{Co(S_2N_2H)_2\}_2\{Ph_2P(O)CH_2CH_2P(O)Ph_2\}]$, 907–14
AZIDE
Electron spin resonance and optical spectroscopic studies on Cu^{2+} and Co^{2+} in $Cs_2[Zn(N_3)_4]$, 939–42
The relative nucleophilic efficiency of water and the azide and thiocyanate ions in the aqueous substitution of chloropentacyanocobaltate(III), 1059–62
AZOLIDINYLIDENE
Syntheses and reactions of acyclic and cyclic carbene complexes *cis*-$[MI(=CMeNHMe)(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo or W) and *cis*-$[MI\{=\overline{C(CH_2)_nNMe}\}(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo, *n* = 3 or 4; M = W, *n* = 3) obtained through alkyl migration to methyl isocyanide co-ordinated to molybdenum or tungsten. *X*-Ray structure of *cis*-$[MI\{=\overline{C(CH_2)_3NMe}\}(CO)_2(\eta\text{-}C_5H_5)]$ (1986, 2127), 479–80
BARIUM
Complexes of ligands providing endogenous bridges. Part 1. The syntheses and crystal structures of barium and lead(II) complexes of macrocyclic Schiff bases derived from heterocyclic dicarbonyls and 1,*n*-diamino-*n'*-hydroxyalkanes (*n*,*n'* = 3,2; 4,2; or 5,3), 207–18
Complexes of ligands providing endogenous bridges. Part 3. Non-transition metal complexes of macrocycles derived from 2,6-diacetyl-4-methylphenol; the crystal structure of a mononuclear barium complex, 1865–8
BASICITY
Analysis and prediction of acid–base reactions between oxides and oxysalts using the optical basicity concept, 2323–8
BENZENE
Activation of alkanes involving rhenium atoms: synthesis and electronic structure of binuclear μ-alkylidene derivatives; molecular structure of $[\{Re(\eta\text{-}C_6H_6)\}_2(\mu\text{-}CHBu^t)(\mu\text{-}H)_2]$, 391–402
Synthesis of η-arene rhenium compounds using rhenium atoms. Crystal structures of $[Re_2(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)_2(\mu\text{-}H)_2]$ and $[Re(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)H]$, 403–10
(η-Benzene)bis(trimethylphosphine)iron: a useful precursor to polyene iron derivatives. Crystal structure of 2—4-η:2′— 4′-η-1,1′-bi(cyclohex-3-en-2-yl)bis(trimethylphosphine)iron(II), 2031–8
Dinitrogen complexes derived from tricarbonyl(η^6-hexaethylbenzene)chromium(0): crystal and molecular structure of μ-dinitrogen-bis[dicarbonyl(η^6-hexaethylbenzene)-chromium(0)]–toluene (1/1), 2789–92
BENZIMIDAZOLE
New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states, 2621–30
BENZIMIDAZOLYL
Copper(I) and copper(II) compounds of a dinucleating, octadentate tetrakisbenzimidazole ligand. Crystal and molecular structures of three dihalogeno{1,8-bis[bis(1′-methylbenzimidazol-2′-ylmethyl)-amino]-3,6-dioxaoctane}dicopper(II) bis(anion) compounds (halogen = F, Cl, or Br; anion = BF_4, CF_3SO_3, or Cl), 2579–94
BENZOATE
The cyclometallation of benzoic acid to give rhodium, iridium, and osmium *C*,*O*-benzoates. *X*-Ray structure determination of the dibenzoate $[(C_5Me_5)Rh(OOCPh)_2(H_2O)]$, 2459–66
BENZOIC ACID
The cyclometallation of benzoic acid to give rhodium, iridium, and osmium *C*,*O*-benzoates. *X*-Ray structure determination of the dibenzoate $[(C_5Me_5)Rh(OOCPh)_2(H_2O)]$, 2459–66
BENZOTHIAZOLE
New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states, 2621–30
BENZOYL
Phenyl complexes of ruthenium(II): competition between carbonyl substitution and formation of benzoyl complexes, and an *X*-ray study of a benzoyl complex, 27–34
BERYLLIUM
Complex formation of beryllium(II) with salicylate and hydroxide ions in 1 mol dm^{-3} sodium perchlorate aqueous solution at 25 °C, 1853–6

BERYLLIUM (contd)
Beryllium(II) hydrolysis in 3.0 mol dm^{-3} perchlorate, 2431–8
Studies of metal carbonate equilibria. Part 15. The beryllium(II)–water–carbon dioxide(g) system in acidic 3.0 mol dm^{-3} perchlorate media, 2439–44
Studies of metal carbonate equilibria. Part 16. The beryllium(II)–water–carbon dioxide(g) system in neutral-to-alkaline 3.0 mol dm^{-3} perchlorate media at 25 °C, 2445–50
BIACETYL
Photo-oxidations of acetoin and biacetyl catalysed by tetrabutylammonium decatungstate(4−) in acetonitrile under excess of oxygen, 961–2
BICYCLOHEPTANE
Carbene complexes in organic synthesis: a new tungstacarbene-promoted alkyne insertion–olefin cyclopropanation reaction, 1041–50
BICYCLOHEXENE
(η-Benzene)bis(trimethylphosphine)iron: a useful precursor to polyene iron derivatives. Crystal structure of 2—4-η:2′— 4′-η-1,1′-bi(cyclohex-3-en-2-yl)bis(trimethylphosphine)iron(II), 2031–8
BIPHENYLYLENEDIMETHYLENE
Axially asymmetric metal alkyls. Part 5. Synthesis and reduction to Zr^{III} species of Group 4 metallepines $[ML(\eta\text{-}C_5H_5)_2]$[L = $(2\text{-}CHRC_6H_4)_2^{2-}$, R = H, M = Ti, Zr, or Hf), isomers of $[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), and *meso*-$[M\{1,8\text{-}(CHSiMe_3)_2C_{10}H_6\}(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf); *X*-ray crystal structures of $[TiL(\eta\text{-}C_5H_5)_2]$ (R = H) and λ-and δ-*RS*-$[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), 2347–58
BIPYRAZOLE
Reactivity of ruthenium(II) complexes with 1-hydroxymethyl- or 1-benzyl-3,5-dimethylpyrazole and similar functionalized bipyrazoles. *X*-Ray crystal structure of carbonylchlorohydrido(3,5-dimethylpyrazole-N^2)bis(triphenylphosphine)ruthenium(II), 183–6
BIPYRIDYL
The first authenticated example of geometrical isomers in organotin(IV) co-ordination chemistry. The crystal and molecular structures of octahedral *cis*- and *trans*-SnR_2 isomeric adducts of dichlorobis(4-chlorophenyl)tin(IV) with 4,4′-dimethyl-2,2′-bipyridyl, 129–38
Mono- and di-carbonyl complexes of molybdenum(II) with mixed bidentate ligands. *X*-Ray crystal structure of $[MoBr_2(CO)(bipy)(Ph_2PCH_2PPh_2)]$ (bipy = 2,2′-bipyridine): a novel type of $[M(bidentate)_2(unidentate)_3]$ structure, 819–24
Ternary metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of copper(II)–2,2′-bipyridyl– pyridoxic acid and copper(II)–2,2′-bipyridyl–pyridoxic acid γ-lactone complexes, 1009–16
Water photolysis. Part 2. An investigation of the relative advantages of various components of the sensitiser–electron relay–metal colloid system for the photoproduction of hydrogen from water, and the use of these systems in the photohydrogenation of unsaturated organic substrates, 1299–304
Spectroscopic, magnetic, and structural properties of diaquadimethanolbis(*N*-tosylvalinato)copper(II) and its 2,2′-bipyridine adduct, 1333–40
Synthesis, spectroscopic, and electrochemical properties of bis(2,2′-bipyridyl)ruthenium compounds of some pyridyl-1,2,4-triazoles, 1389–96
Aqua(2,2′-bipyridyl)difluorocopper(II) dihydrate: *X*-ray structure reveals short hydrogen bonds and other unusual features, 2397–400
Characterizations of the redox orbitals of mixed-ligand 2,2′-bipyridine–2-(2′-pyridyl)quinoline ruthenium(II) complexes, 2467–72
Synthesis, characterisation, and photochemical properties of a series of ruthenium containing metallopolymers based on poly-*N*-vinylimidazole, 3073–8
BISMUTH
The crystal structures of the diphenyldithiophosphinates of antimony(III) and bismuth(III): $M(S_2PPh_2)_3$ (M = Sb or Bi), 145–50
The crystal structure of bismuth diethyldithiophosphinate–benzene (1/1), $Bi(S_2PEt_2)_3{\cdot}C_6H_6$, 1257–60
BORANE
Reaction of monoborane with tris(cyclopentadienyl)uranium(IV) derivatives, 691–4
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$], 1805–12
The methyl/trifluoromethyl substitution effect in boranes. A gas-phase electron diffraction and vibrational study of bis(dimethylamino)(trifluoromethyl)borane and dimethylaminobis(trifluoromethyl)borane, 1839–46
Regioselective synthesis of $[1\text{-}B_{10}H_9(SH)]^{2-}$ and $[2\text{-}B_{10}H_9(SH)]^{2-}$: potential agents for boron–neutron capture therapy of brain tumours, 1953–6
Synthesis, molecular structure, and nuclear magnetic resonance spectra of the anionic monocarbametallaborane $[NEt_4][(CO)_3(PPh_3)(OH)\text{-}closo\text{-}MoCB_{10}H_{10}]{\cdot}CH_2Cl_2$, 2385–90
Pentamethylcyclopentadienylrhodaborane chemistry. Part 3. The reaction of $[\{Rh(\eta^5\text{-}C_5Me_5)Cl_2\}_2]$ with *closo*-$[B_{10}H_{10}]^{2-}$ and *arachno*-$B_{10}H_{12}$-6,9-$(PMe_2Ph)_2$, and the characterisation of some novel rhodaboranes by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 2417–30
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 67. Reactions of the salts $[X][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$[X = $P(CH_2Ph)Ph_3$, R = C_6H_4Me-4; X = NEt_4, R = Ph] with bis(cyclo-octa-1,5-diene)platinum, 2661–8
Gold–boron chemistry. Part 1. Synthetic, structural, and spectroscopic studies on the compounds [5,6-μ-$(AuPR_3)$-*nido*-$B_{10}H_{13}$] (R = cyclo-C_6H_{11} or C_6H_4Me-2), 2761–8
Organoruthenaborane chemistry. Part 5. Products of the reaction between *closo*-$[B_{10}H_{10}]^{2-}$ and $[(\eta^6\text{-}C_6Me_6)ClRuB_3H_8]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[\{(\eta^6\text{-}C_6Me_6)_2Ru_2H_4\}RuB_{10}H_8(OEt)_2]$, 2781–8
The syntheses and crystal and molecular structures of some ten-vertex (MCB_8; M = Ru or Os) metallacarbaboranes: 9,9,9-$(CO)(PPh_3)_2$-*nido*-9,6-MCB_8H_{10}-5-PPh_3)] (M = Ru or Os) and [9,9,9-(η-C_5H_5)(H)(PPh_3)-*arachno*-9,6-$RuCB_8H_{12}$], 2793–802
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 68. Reactions between nonacarbonyldi-iron and the salts $[NEt_4][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = Me, Ph, or C_6H_4Me-4); crystal structures of $[NEt_4][FeW\{\mu\text{-}CH(C_6H_4Me\text{-}4)\}(\mu\text{-}\sigma\text{:}\eta^5\text{-}C_2B_9H_8Me_2)(\mu\text{-}CO)(CO)_5]$ and $[NEt_4][FeW(\mu_3\text{-}CPh)(\mu\text{-}\sigma\text{:}\sigma',\eta^5\text{-}C_2B_9H_7Me_2)(CO)_8]$, 2917–26
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$] (1987, 1805), 3107–8
BORATE
A molybdenum-95 and nitrogen-14 nuclear magnetic resonance study of six-co-ordinate hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes containing a sixteen-electron $\{Mo(NO)\}^4$ core, 619–22
Studies on borate esters. Part 8. Interactions of cations with oxyacid anion-bridged esters of D-glucarate in alkaline media, 2051–8
BORON
Matrix isolation studies of monomeric boron trisulphide, 267–70
Ruthenium(II) complexes of *closo* dodecaboranyl anions and the molecular structure of the *pileo* thirteen-vertex ruthenaborane $[(PPh_3)_2ClRuB_{12}H_{11}(NEt_3)]$, 451–6
Reactions of heteroboranes. Synthesis of [2,2-$(PPh_3)_2$-1,2-$SePtB_{10}H_{10}$]-CH_2Cl_2, its crystal and molecular structure and that of $SeB_{11}H_{11}$, 699–704
Stability constants for hypochloritoborate and hypobromitoborate complex ions in aqueous solution, 943–6
Extended Hückel molecular orbital study of the effects of edge-bridging hydrogen atoms on the lengths of boron–boron and metal–metal bonds in cluster compounds, and the crystal structure of benzyltrimethylammonium octahydrotriborate(1−), 1017–26
The preparation of $[Ru(NCMe)_6][7\text{-}(\eta^6\text{-}C_6Me_6)\text{-}nido\text{-}7\text{-}RuB_{10}H_{13}]_2$ and its characterization by single-crystal *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy: a triply hydrogen-bridged eleven-vertex *nido*-7-metallaundecaborane, 1169–74
Pentamethylcyclopentadienylrhodaborane chemistry. Part 2. The reaction of [6-$(\eta^5\text{-}C_5Me_5)$-*nido*-6-RhB_9H_{13}] with dimethylphenylphosphine, and the characterization of [5-$(\eta^5\text{-}C_5Me_5)$-*nido*-5-RhB_9H_{11}-7-(PMe_2Ph)], [2-$(\eta^5\text{-}C_5Me_5)$-*closo*-2-RhB_9H_7-3,10-$(PMe_2Ph)_2$], and [2-$(\eta^5\text{-}C_5Me_5)$-*nido*-2-RhB_8H_{10}-5-(PMe_2Ph)] by *X*-ray diffraction analysis and nuclear magnetic

BORON (contd)
resonance spectroscopy, 1431–44
Identification of the *endo,exo* isomer of 6,9-$(PMe_2Ph)_2$-*arachno*-$B_{10}H_{12}$ by nuclear magnetic resonance spectroscopy, 1573–6
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$], 1805–12
The methyl/trifluoromethyl substitution effect in boranes. A gas-phase electron diffraction and vibrational study of bis(dimethylamino)(trifluoromethyl)borane and dimethylaminobis(trifluoromethyl)borane, 1839–46
Regioselective synthesis of $[1\text{-}B_{10}H_9(SH)]^{2-}$ and $[2\text{-}B_{10}H_9(SH)]^{2-}$: potential agents for boron–neutron capture therapy of brain tumours, 1953–6
Studies on borate esters. Part 8. Interactions of cations with oxyacid anion-bridged esters of D-glucarate in alkaline media, 2051–8
Synthesis, molecular structure, and nuclear magnetic resonance spectra of the anionic monocarbametallaborane $[NEt_4][(CO)_3(PPh_3)(OH)$-*closo*-$MoCB_{10}H_{10}]\cdot CH_2Cl_2$, 2385–90
Pentamethylcyclopentadienylrhodaborane chemistry. Part 3. The reaction of $[\{Rh(\eta^5\text{-}C_5Me_5)Cl_2\}_2]$ with *closo*-$[B_{10}H_{10}]^{2-}$ and *arachno*-$B_{10}H_{12}$-6,9-$(PMe_2Ph)_2$, and the characterisation of some novel rhodaboranes by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 2417–30
Gold–boron chemistry. Part 1. Synthetic, structural, and spectroscopic studies on the compounds [5,6-μ-$(AuPR_3)$-*nido*-$B_{10}H_{13}$] (R = cyclo-C_6H_{11} or C_6H_4Me-2), 2761–8
Organoruthenaborane chemistry. Part 5. Products of the reaction between *closo*-$[B_{10}H_{10}]^{2-}$ and $[(\eta^6\text{-}C_6Me_6)ClRuB_3H_8]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[\{(\eta^6\text{-}C_6Me_6)_2Ru_2H_4\}RuB_{10}H_8(OEt)_2]$, 2781–8
The syntheses and crystal and molecular structures of some ten-vertex (MCB_8; M = Ru or Os) metallacarbaboranes: 9,9,9-$(CO)(PPh_3)_2$-*nido*-9,6-MCB_8H_{10}-5-PPh_3)] (M = Ru or Os) and [9,9,9-$(\eta\text{-}C_5H_5)(H)(PPh_3)$-*arachno*-9,6-$RuCB_8H_{12}$], 2793–802
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$] (1987, 1805), 3107–8

BORON-11
Studies on borate esters. Part 8. Interactions of cations with oxyacid anion-bridged esters of D-glucarate in alkaline media, 2051–8

BROMIDE
Electrolytic behaviour of transition metal bromides in acetonitrile solution, 2863–6

BUTADIENE
Organic chemistry of dinuclear metal centres. Part 9. Diruthenium complexes of buta-1,3-diene. *X*-Ray crystal structure of $[Ru_2(CO)_2(\mu\text{-}s\text{-}trans\text{-}\eta^2\text{:}\eta^{2'}\text{-}CH_2CHCHCH_2)(\eta\text{-}C_5H_5)_2]$, 2703–8

BUTADIENOLATE
Reactions of co-ordinated ligands. Part 42. The synthesis and reactions of cationic η^4-tetraphenylcyclobutadiene-ruthenium complexes; *X*-ray crystal structures of $[Ru(NCMe)(\eta^4\text{-}C_4Ph_4)(\eta\text{-}C_5H_5)][BF_4]$ and $[Ru\{\eta^5\text{-}CH(Ph)C(Ph)C(Ph)C(Ph)O\}(\eta\text{-}C_5H_5)]$, 2803–14

BUTANONE OXIMATE
Syntheses, spectroscopic, and electrochemical properties of dinuclear copper(II) complexes bridged by a single thiolate sulphur atom; *X*-ray crystal structures of $[Cu_2(L^1)_2(SC_6H_4Cl\text{-}4)][PF_6]$ and $[Cu_2(L^2)_2(SC_6H_4CH_3\text{-}4)][ClO_4]$, 241–8

BUTENYL
Synthesis and reactions of the co-ordinatively unsaturated hexafluorobut-2-yne complex $[WBr_2(CO)(CF_3C{\equiv}CCF_3)_2]$. The crystal and molecular structure of $[\{W[C(CF_3){=}C(CF_3)P(OEt)_2O](\mu\text{-}Br)(CO)(CF_3C{\equiv}CCF_3)\}_2]$, 2165–70

BUTYNE
Synthesis and reactions of the co-ordinatively unsaturated hexafluorobut-2-yne complex $[WBr_2(CO)(CF_3C{\equiv}CCF_3)_2]$. The crystal and molecular structure of $[\{W[C(CF_3){=}C(CF_3)P(OEt)_2O](\mu\text{-}Br)(CO)(CF_3C{\equiv}CCF_3)\}_2]$, 2165–70

BUTYNYL
Unsymmetrical bridging of an acetylide ligand in the cation $[Pt_2(C{\equiv}CBu^t)_3(\mu\text{-}dppm)_2]^+$ (dppm = $Ph_2PCH_2PPh_2$): *X*-ray structure determination of the tetraphenylborate salt, 1659–64

CADMIUM
Investigation into aroylhydrazones as chelating agents. Part 8. Synthesis and spectroscopic characterization of complexes of Co, Ni, Cu, Zn, and Cd with 2,6-diacetylpyridine bis(salicyloylhydrazone); *X*-ray crystal structure of dichloro[2,6-diacetylpyridine bis(salicyloylhydrazone)]cadmium(II)–chloroform–methanol (1/1/1), 177–82
Reactions of chromium(III) complexes of 1,10-phenanthroline, 2,2′-bipyridyl, and oxalate with the pulse radiolytically generated aquated electron, zinc(I), and cadmium(I), 1425–30
Preparation and crystal structures of $[PPh_4]_2[M_2(SPh)_6]$ (M = Zn or Cd), 1577–80
Comments on the attempts by Wickens and Abrams to synthesize manganese(II) phosphine complexes: a comment, 1581–2
Exchange of cadmium into the sodium and ammonium forms of the natural zeolites clinoptilolite, mordenite, and ferrierite, 1911–6
Thermal investigation and stereochemical studies of some cyclic diamine complexes of nickel(II), zinc(II), and cadmium(II) in the solid state, 2089–94
Polymeric thiolate complexes of Group 12 metals. Crystal and molecular structures of *catena*-[μ-(3-dimethylammonio-1-propanethiolate)]-dichlorocadmium(II) and bis[3-(dimethylammonio)propyl] disulphide tetrabromocadmate(II), 2391–6

CALCIUM
Studies on alkaline-earth metal sulphites. Part 9. Solid solutions of $Ca_3(SO_4)_2SO_3\cdot 12H_2O$ in $Ca_3(SO_3)_2SO_4\cdot 12H_2O$: structure, stability, and dehydration products, 235–40

CARBAMATE
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S{=}C(NHPh)\text{-}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O{-}C(NHPh){=}S\}]$, 2299–304
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S{=}C(NHPh)\text{-}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O{-}C(NHPh){=}S\}]$ (1987, 2299), 3107–8

CARBENE
Syntheses and reactions of acyclic and cyclic carbene complexes *cis*-$[MI(=CMeNHMe)(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo or W) and *cis*-$[MI\{=\overline{C(CH_2)_n}NMe\}(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo, *n* = 3 or 4; M = W, *n* = 3) obtained through alkyl migration to methyl isocyanide co-ordinated to molybdenum or tungsten. *X*-Ray structure of *cis*-$[MI\{=\overline{C(CH_2)_3}NMe\}(CO)_2(\eta\text{-}C_5H_5)]$ (1986, 2127), 479–80
Carbene complexes in organic synthesis: a new tungstacarbene-promoted alkyne insertion–olefin cyclopropanation reaction, 1041–50
Carbene complexes. Part 21. Synthesis and characterisation of bis(carbene)molybdenum(II) complexes and dimetal(0) complexes of the Group 6 elements containing novel bridging bis(carbene) ligands; *X*-ray structures of $[Mo(CO)_2(L^{Et})_2(OSO_2CF_3)_2]$ $[L^{Et} = =\overline{CN(Et)(CH_2)_2}NEt]$ and $[W(CO)_5\{\overline{C(OEt)CH_2C_6H_4\text{-}CH_2C(OEt)\text{-}o}\}W(CO)_5]$, 2843–52

CARBIDE
High nuclearity osmium cluster radicals, 1051–4
Transformation of $[PPh_4][Fe_5RhC(CO)_{16}]$ under hydroformylation conditions: synthesis and crystal structure of $[PPh_4][Fe_3Rh_3C(CO)_{15}]$, 1983–8
Reactivity of a cluster-co-ordinated acyl. Reduction of a μ-acetyl ligand to acetaldehyde, 2839–42

CARBON-13
Dithio-oxamides as ligands: crystal structures and vibrational analyses of bis(*N,N′*-dicyclohexyldithio-oxamidato)palladium(II), bis(*N,N′*-dibenzyldithio-oxamide)copper(II) diperchlorate, and dichloro(*N,N′*-dimethyldithio-oxamide)zinc(II) and the carbon-13 nuclear magnetic resonance spectra of related compounds, 1921–8
Activation energies for molecular tumbling and cyclopentadienyl rotation in $[M(\eta^5\text{-}C_5H_5)(\eta^4\text{-cod})]$ (M = Rh or Ir; cod = cyclo-octa-1,5-diene) and the *X*-ray crystal structure of and bonding in $[Rh(\eta^5\text{-}C_5H_5)(\eta^4\text{-cod})]$, 1947–52

CARBON-13 (contd)
Preparation and characterisation of mononuclear organoimido complexes of tungsten: *X*-ray crystal structures of $[WCl_4(NC_6H_4Me\text{-}p)(thf)]$, $[P(CH_2Ph)Ph_3][WCl_5(NC_6H_4Me\text{-}p)]$, and $[WCl_2(NPh)_2(bipy)]$, 2067–76
Carbon-13 nuclear magnetic resonance study of the properties of the unsaturated anion $[Re_3(\mu\text{-}H)_4(CO)_9(NCMe)]^-$ at low temperature, 2083–6
Complexes formed between zinc(II) and diethylenetriamine: a carbon-13 nuclear magnetic resonance study, 2889–96
Carbon-13 nuclear magnetic resonance study of the zinc(II) complexes of 3,7-diazanonane-1,9-diamine and 1,2-diaminoethane, 2897–904

CARBONATE
Molten lithium carbonate–sodium carbonate–potassium carbonate eutectic: electronic spectroscopy of first-row transition metal compounds, 2329–32
Studies of metal carbonate equilibria. Part 15. The beryllium(II)–water–carbon dioxide(g) system in acidic 3.0 mol dm^{-3} perchlorate media, 2439–44
Studies of metal carbonate equilibria. Part 16. The beryllium(II)–water–carbon dioxide(g) system in neutral-to-alkaline 3.0 mol dm^{-3} perchlorate media at 25 °C, 2445–50
Infrared spectroscopic studies of uranyl(VI) species adsorbed from aqueous $[UO_2(CO_3)_3]^{4-}$ solutions on to a polymer bearing amidoxime groups, 2489–92
Reactions of tetra-μ-carboxylato-diruthenium(II,II) compounds. *X*-Ray crystal structures of $Ru_2(\mu\text{-}O_2CCF_3)_4(thf)_2$, $Ru_2(\mu\text{-}O_2CR)_4(NO)_2$ (R = Et or CF_3), and $\{Na_3[Ru_2(\mu\text{-}O_2CO)_4]\cdot 6H_2O\}_n$, 2723–36

CARBON DIOXIDE
Electrocatalytic reduction of carbon dioxide to methanol. Part 5. Relationship between the ability of metal complexes to engage in homogeneous catalysis and their co-ordination chemistry, 1377–80
Photochemical reduction of carbon dioxide to carbon monoxide in water using a nickel(II) tetra-azamacrocycle complex as catalyst, 2105–10
Ab initio study of stable bis(carbon dioxide)molybdenum complexes, 2373–8
Studies of metal carbonate equilibria. Part 15. The beryllium(II)–water–carbon dioxide(g) system in acidic 3.0 mol dm^{-3} perchlorate media, 2439–44
Studies of metal carbonate equilibria. Part 16. The beryllium(II)–water–carbon dioxide(g) system in neutral-to-alkaline 3.0 mol dm^{-3} perchlorate media at 25 °C, 2445–50

CARBON DISULPHIDE
Chemistry of metallacyclic complexes containing the FeCSCS ring obtained by metal-promoted CS_2–CS coupling. *X*-Ray crystal structures of $[(CO)(cp)FeC\{SW(CO)_5\}SC(FeL_n)S]$ and $[(CO)(cp)FeC(SEt)SC(FeL_n)S]SO_3CF_3$ [FeL_n = $Fe(cp)(CO)_2$, cp = $\eta\text{-}C_5H_5$], 1133–44
Crystal and molecular structure of $[Co_8(CO)_{21}(\mu_5\text{-}CS_2)(\mu_5\text{-}C_2S)]$: a thiocarbonyl group as a six-electron donor, 1707–12

CARBON MONOXIDE
Solvent effect in the insertion reaction of iron(II) carbonyls, 519–22
Oxygen- and carbon monoxide-binding to a lipophilic diporphyrinatocopperiron complex solubilized in an aqueous medium with a micelle, 2963–8

CARBONYL
Isomers in *fac*-$[ReBr(CO)_3(L_c)_2]$ (L_c = *cis*-3,4-dihydro-2,3,4,5-tetraphenyl-2*H*-1,2,3-diazaphosphole) arising from chirality of phosphorus ligands: separation and *X*-ray structural studies, 21–6
Phenyl complexes of ruthenium(II): competition between carbonyl substitution and formation of benzoyl complexes, and an *X*-ray study of a benzoyl complex, 27–34
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 49. Synthesis of mixed-metal complexes with bonds between Cr, Mo or W and Co, Rh, Ir or Re; crystal structure of $[CrReRh_2(\mu_3\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}CO)(CO)_9(\mu\text{-}C_9H_7)_2]$, 41–50
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 50. Synthesis and crystal structures of the complexes $[RuW_3(\mu\text{-}CO)(\mu_3\text{-}\eta\text{-}CO)(\mu\text{-}CPh)_2(\mu_3\text{-}CPh)(CO)_2(\eta\text{-}C_5H_5)_3]$ and $[RuW_3(\mu\text{-}CO)(\mu_3\text{-}\eta\text{-}CO)\{\mu\text{-}C(Ph)C(O)\}(\mu\text{-}CPh)(\mu_3\text{-}CPh)(CO)(PMePh_2)(\eta\text{-}C_5H_5)_3]$, 51–60
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 51. Alkylidyne(carbaborane)tungsten-gold and -rhodium complexes; crystal structures of $[AuW(\mu\text{-}CR)(CO)_2(PPh_3)(\eta^5\text{-}C_2B_9H_9Me_2)]$, $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$, and $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2\{\eta^5\text{-}C_2B_9(C_7H_9)H_8Me_2\}]$ (R = C_6H_4Me-4), 61–72
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 52. Synthesis and reactions of the compounds $[RuW(\mu\text{-}CR)(CO)_3(\eta\text{-}C_5H_5)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[RuW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta\text{-}C_5H_5)(\eta^5\text{-}C_2B_9H_9Me_2)]$ and $[N(PPh_3)_2][RuW(\mu\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}\sigma,\eta^5\text{-}C_2B_9H_8Me_2)(CO)_3(\eta\text{-}C_5H_5)]$, 73–80
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 53. Alkylidyne(carbaborane)dimetal complexes with tungsten–molybdenum and –tungsten bonds; crystal structures of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ [M = Mo (two isomers) or W], 81–90
High-yield syntheses of sulphido triruthenium carbonyl cluster complexes containing bis(diphenylphosphino)methane, 91–6
Synthesis and properties of heterotrimetallic complexes containing M–Pt–Hg (M = Mo, W, Fe, Mn, or Co) arrays, 119–22
Bimetallic systems. Part 16. Formation of Group 6 metal(0)–platinum(II) hydride complexes, by ring opening of *fac*- or *mer*-$[M(CO)_3(dppm\text{-}PP')(dppm\text{-}P)]$ (M = Cr, Mo, or W; dppm = $Ph_2PCH_2PPh_2$). Reactions of $[(CO)_3Mo(\mu\text{-}dppm)_2PtH(Cl)]$, 221–6
Synthesis of the alkoxo(hydrido)-clusters $[Ru_3(\mu\text{-}H)(CO)_{10}(\mu\text{-}OR)]$ [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4
Synthesis and *X*-ray crystal structure of a cuboctahedral platinum cluster, 259–62
Reactions of metal carbonyl cluster complexes with multidentate phosphine ligands; the preparation and thermolysis of bis(dimethylphosphino)methane derivatives of $[Ru_3(CO)_{12}]$, and the *X*-ray crystal structure of $[Ru_3(CO)_9(H)(Me_2PCHPMe_2)]$, 277–80
Reaction of $[Os_5H_2(CO)_{15}]$ with pyridine: crystal and molecular structures of $[Os_5H_3(CO)_{14}(C_5H_4N)]$ and $[Os_5H_2(CO)_{14}(C_5H_5N)]$, 327–34
Chemistry of platinum sulphido-complexes. Part 6. Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with the rhodium complexes $[(RhL_2Cl)_2]$ (L = CO or C_2H_4) and $[Rh(2,6\text{-}Me_2C_6H_3NC)_4]PF_6$. Crystal and molecular structures of $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(CO)_2][Rh(CO)_2Cl_2]\cdot C_3H_6O$ and $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(2,6\text{-}Me_2C_6H_3NC)_2]PF_6$, 335–40
The addition of isonitriles to $[\{M(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (M = Mo or W). Syntheses and properties of $[M_2(\mu\text{-}\eta^2\text{-}CNR)(CO)_4(\eta\text{-}C_5H_5)_2]$ (M = Mo or W, R = Me or Bu^t) and $[\{Mo(CNR)(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (R = Me or Bu^t). *X*-Ray crystal structures of $[Mo_2(\mu\text{-}\eta^2\text{-}CNBu^t)(CO)_4(\eta\text{-}(\eta\text{-}C_5H_5)_2]$ and $[\{Mo(CNBu^t)(CO)_2(\eta\text{-}C_5H_5)\}_2]$, 341–8
Thermal and photochemical behaviour of organotetraruthenium clusters: solution structures and dynamics of phosphine-substituted derivatives, 349–54
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au), 379–90
Sulphur–arsenic ligands. Part 1. Dimethyl(methylthio)arsine complexes of halogenotrimethylplatinum(IV) and tricarbonylhalogenorhenium(I), 417–20
Formation of a methoxycarbonyl group by attack of methanol on an iron(II) carbonyl complex, assisted by intramolecular proton transfer: *X*-ray structure determination of $[Fe(CO)(PMe_3)_2(CNC_6H_{11})\{CO(OCH_3)\}\{C(CH_3)=NHC_6H_{11}\}]BPh_4$, 473–6
Kinetics and mechanism of ring closure for a series of photoproduced $[M(CO)_5L]$ intermediates (M = Cr, Mo, or W; L = pyridine 2-carbaldehyde imine ligand), 513–8
Solvent effect in the insertion reaction of iron(II) carbonyls, 519–22
Phosphorus-tellurium compounds. Part 3. Cyclopentadienyliron tellurophosphorane complexes: tellurium-bridged iron phosphine compounds?, 541–4
Triosmium clusters containing ligands derived from tropone

CARBONYL (contd)
(cycloheptatrienone): molecular structure of an oxidative addition product $[Os_3H(\mu_3\text{-}C_7H_5O)(CO)_9]$, 551–6
The vibrational spectrum of tricarbonyl(trichlorostibine)nickel(0), $[Ni(CO)_3(SbCl_3)]$, 681–2
Vibrational studies of interstitial hydrogen in metal carbonyl clusters, 685–90
Rhenium carbonyl fluorides: preparation of $[Re(CO)_6][ReF_6]$ and some reactions of $[Re(CO)_5F{\cdot}ReF_5]$, 741–6
Heteronuclear 1,3-intramolecular metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of 2-thia-4-selenapentane. A dynamic nuclear magnetic resonance investigation, 757–60
Bimetallic systems. Part 17. Synthesis of Group 6 metal(0)–copper(I), –silver(I), or –gold(I) complexes with two $Ph_2PCH_2PPh_2$ (dppm) bridging ligands: crystal structure of $[(OC)_2W(\mu\text{-}CO)(\mu\text{-dppm})_2(\mu\text{-}Cl)Cu]{\cdot}CH_2Cl_2$, 769–76
Synthesis and characterisation of $[N(PPh_3)_2][RuRh_5(CO)_{16}]$ by *X*-ray crystallography and by natural-abundance carbon-13 nuclear magnetic resonance spectroscopy, 781–4
Mono- and di-carbonyl complexes of molybdenum(II) with mixed bidentate ligands. *X*-Ray crystal structure of $[MoBr_2(CO)(bipy)(Ph_2PCH_2PPh_2)]$ (bipy = 2,2′-bipyridine): a novel type of $[M(bidentate)_2(unidentate)_3]$ structure, 819–24
Transition-metal mediated asymmetric synthesis. Part 5. Alternative methods for the formation of tricarbonyl(η^5-cyclohexadienyl)-iron(1+) salts, 885–8
Trinuclear osmium clusters, containing bidentate $Ph_2PCH_2PPh_2$ and $Me_2PCH_2PMe_2$ ligands: *X*-ray crystal structures of the complexes $[Os_3H(CO)_9(Me_2PCHPMe_2)]$ and $[Os_3H(OH)(CO)_8(Ph_2PCH_2PPh_2)]$, 931–8
Complexes of the platinum metals. Part 32. Synthesis and chemistry of some iridium sulphonate derivatives, 947–52
Phosphinoalkylsilyl complexes. Part 7. Synthesis and reactivity of octahedral, bis(chelate)dicarbonyl complexes of ruthenium(II) and osmium(II). Conformational properties of the bis(chelate) framework and *X*-ray crystal structures of $[M(PPh_2CH_2CH_2SiMe_2)_2(CO)_2]$ (M = Ru or Os), 969–74
2-Pyridonate (Opy) rhodium(I) carbonyl complexes. Crystal structure of the unusual tetranuclear complex $[Rh_4(\mu\text{-Opy})_4(\mu\text{-}CO)_2(CO)_4]$, 981–4
The heteronuclear cluster chemistry of the Group 1B metals. Part 3. Synthesis, structures, and chemical properties of the trimetallic hexanuclear Group 1B metal cluster compounds $[MM'Ru_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]$ (M = Cu, M′ = Ag or Au; M = Ag, M′ = Au). *X*-Ray crystal structure of the complex $[AgCuRu_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]{\cdot}CH_2Cl_2$, 1001–8
Extended Hückel molecular orbital study of the effects of edge-bridging hydrogen atoms on the lengths of boron–boron and metal–metal bonds in cluster compounds, and the crystal structure of benzyltrimethylammonium octahydrotriborate(1−), 1017–26
High nuclearity osmium cluster radicals, 1051–4
Chemistry of metallacyclic complexes containing the $\overline{FeCSCS}$ ring obtained by metal-promoted CS_2–CS coupling. *X*-Ray crystal structures of $[(CO)(cp)\overline{FeC\{SW(CO)_5\}SC(FeL_n)S}]$ and $[(CO)(cp)\overline{FeC(SEt)SC(FeL_n)S}]SO_3CF_3$ [FeL_n = Fe(cp)(CO)$_2$, cp = η-C_5H_5], 1133–44
Photoactivation of methane by η^5-cyclopentadienyl and substituted η^5-cyclopentadienyl Group 8 metal dicarbonyl complexes, $[M(\eta^5\text{-}C_5R_5)(CO)_2]$ (M = Rh or Ir, R = H or Me), and dicarbonyl(η^5-indenyl)iridium: a matrix isolation study, 1181–90
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 54. Compounds derived from $[W(\equiv CC_6H_4Me\text{-}4)(CO)_2(\eta\text{-}C_5Me_5)]$ with bonds between tungsten and iron, cobalt, or nickel; crystal structure of $[FeW\{\mu\text{-}C(C_6H_4Me\text{-}4)C(O)O\}(CO)_5(\eta\text{-}C_5Me_5)]$, 1191–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 55. Coupling of methylene and alkylidyne groups at a trimetal centre; crystal structures of $[Ru_3W\{\mu\text{-}\sigma\text{:}\eta\text{-}trans\text{-}CH{=}C(H)R\}(\mu\text{-}CO)(CO)_3(\eta\text{-}C_5H_5)_2(\eta\text{-}C_5Me_5)]{\cdot}CH_2Cl_2$ and $[Ru_2W(\mu_3\text{-}CCH_2R)(\mu\text{-}CO)_3(CO)(\eta\text{-}C_5H_5)_3]$ (R = C_6H_4Me-4), 1201–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 56. Synthesis of iron–molybdenum compounds; crystal structures of $[FeMo(\mu\text{-}CR)(CO)_6(\eta\text{-}C_5H_5)]$ and $[FeMo_2(\mu_3\text{-}RC_2R)(CO)_6(\eta\text{-}C_5H_5)_2]$ (R = C_6H_4Me-4), 1209–14
carbyne ligands. Part 57. Reactions of iron– molybdenum complexes with oxygen and sulphur; crystal structure of $[FeMo(\mu\text{-}\eta^2\text{-}SCC_6H_4Me\text{-}4)(CO)_5(\eta\text{-}C_5H_5)]$, 1215–20
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 58. Some reactions of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (M = Mo or W); crystal structures of the complexes $[W_2(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_2(L)(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$-[L = $N_2C(C_6H_4Me\text{-}4)_2$ or O], 1221–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 59. Reactions of the alkylidyne compounds $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Cr, Mo, or W, R = C_6H_4Me-4; M = W, R = Me, $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate] with d^{10} complexes; crystal structure of $[PtW_2(\mu\text{-}CMe)_2(CO)_4\{HB(pz)_3\}_2]$, 1229–34
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 60. Di- and tri-metal compounds prepared from the metal alkylidyne complexes $[W(\equiv CR)(CO)_2(\eta\text{-}C_5Me_5)]$ (R = Me or C_6H_4Me-4) and $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Mo, R = C_6H_4Me-4; M = W, R = Me; $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate], 1235–42
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 61. Reactions of iron– molybdenum compounds with tertiary phosphines, 1243–8
Carbon-13 and phosphorus-31 nuclear magnetic resonance studies of the fluxional behaviour of the complexes $[Fe_3(CO)_{10}\{P(OMe)_3\}_2]$ and $[Fe_3(CO)_{10}\{P(OMe)_3\}\{P(OPh)_3\}]$, 1249–52
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au) (1987, 379), 1283–4
The preparation, characterisation, and some reactions of $[Os_3H(X)(CO)_9(NMe_3)]$ (X = Cl, Br, or I), 1285–8
Systematic cluster build up by the reaction of $[Os_3H(X)(CO)_9(MeCN)]$ (X = SR or NHMe; R = Me, Et, or Ph) with $[OsH_2(CO)_4]$, 1289–92
Substitution reactions on triosmium clusters of the type $[Os_3(\mu\text{-}H)(\mu\text{-}X)(CO)_{10}]$ [X = NHMe, O_2CR, C_5H_4N, or OC(NHMe); R = H, Me, or Ph], 1293–8
Reactions of co-ordinated ligands. Part 40. Synthesis, structure, and reactivity of dirhenium μ-allylidene complexes; crystal structures of $[Re_2(\mu\text{-}\sigma\text{:}\eta^3\text{-}CHCHCMe_2)(CO)_8]$ and $[Re_2(\mu\text{-}\sigma\text{:}\eta^3\text{-}CHCHCMe_2)(CO)_9]$, 1313–8
Reduction–oxidation properties of organotransition-metal complexes. Part 26. The synthesis of paramagnetic triazenido-and aryl(1-aryliminoethyl)amido-bridged dirhodium carbonyl derivatives containing the $[Rh_2]^{3+}$ core, 1403–8
Triosmium clusters containing the ligands CNH_2, NCHCN, and NCMeCN derived from cyanide ion, cyanogen, and acetonitrile, 1457–62
Bimetallic systems. Part 19. Chromium(0)–, molybdenum(0)–, or tungsten(0)–iron(0) carbonyl complexes containing one or two bridging $Ph_2PCH_2PPh_2$ ligands: crystal structure of $[(OC)_3Mo(\mu\text{-}Ph_2PCH_2PPh_2)_2Fe(CO)_3]$, 1489–94
Bimetallic systems. Part 20. Synthesis of Group 6 metal(0)–iridium(I) or –rhodium(I) complexes bridged by $Ph_2PCH_2PPh_2$, 1495–500
Bimetallic systems. Part 22. Molybdenum(0)– and manganese(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the reversible uptake of carbon monoxide: crystal structure of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)Mo(CO)_4]{\cdot}0.5C_6H_5Cl$, 1509–14
Intermediates in the conversion of $[Os(CO)_{11}(PRPh_2)]$ (R = Me or Ph) into $[Os_3(\mu_3\text{-}C_6H_4)(\mu_3\text{-}PR)(CO)_9]$: crystal and molecular structures of $[Os_3(\mu\text{-}H)(\mu_3\text{-}C_6H_4PMePh\text{-}o)(CO)_9]$ and $[Os_3(\mu_3\text{-}C_6H_4PMe\text{-}o)(CO)_{10}]$, 1529–34
The triphenylphosphine cone angle and restricted rotation about the chromium–phosphorus bond in dicarbonyl(η^6-hexaalkylbenzene)(triphenylphosphine)chromium(0) complexes. Crystal and molecular structure of dicarbonyl(η^6-hexa-n-propylbenzene)(triphenylphosphine)chromium(0), 1545–50
Preparation and reactivity of halide-bridged binuclear carbonyl manganese and rhenium anions, $[M_2(\mu\text{-}X)(CO)_{8-2n}\{(EtO)_2POP(OEt)_2\}_n]^-$ (X = halide; M = Mn, *n* = 1 or 2; M = Re, *n* = 1). *X*-Ray crystal structure of $[Mn_2Au(\mu\text{-}$

CARBONYL (contd)
Br)(CO)$_6$(μ-tedip)(PPh$_3$)], 1551–6
Synthesis and substitution reactions of a heterodimetallic molybdenum–manganese complex, [MoMn(μ-H)(μ-PPh$_2$)(η^5-C$_5$H$_5$)(CO)$_6$]; *X*-ray crystal structure of [MoMn(μ-H)(μ-PPh$_2$)(η^5-C$_5$H$_5$)(CO)$_4$(dppm-*PP'*)], 1557–64
Synthesis and characterization of new heterobimetallic iron–cobalt complexes with ethenyl bridges CR^2=CHR^1 [R^1 = H or C(O)OMe, R^2 = C(O)OMe or C(O)OEt]. *X*-Ray structure of [FeCo(CO)$_6$\{μ-MeOC(O)C=C(H)C(O)OMe\}], 1619–22
Substituted osmium clusters containing labile ligands. The preparation, characterisation, and some reactions of [Os$_3$H(OR)(CO)$_9$(MeCN)] R = H, Me, Et, or Ph) and the isomers of [Os$_3$H(SR)(CO)$_9$(MeCN)] (R = Me, Et, or Ph); the crystal and molecular structure of [Os$_3$H(SPh)(CO)$_9$(PEt$_3$)], 1623–30
Crystal and molecular structure of [Co$_8$(CO)$_{21}$(μ$_5$-CS$_2$)(μ$_5$-C$_2$S)]: a thiocarbonyl group as a six-electron donor, 1707–12
The reversible one-electron oxidation of *trans*-[Mn^{I}L(CO)(Ph$_2$PCH$_2$PPh$_2$)$_2$]$^{+/+}$: synthesis and characterisation of stable manganese(II) derivatives, 1769–74
Cationic ruthenium(II) phosphite complexes. Preparation and properties of monocarbonyl and isocyanide derivatives, 1813–6
Decarboxylation of vinylene carbonate at triosmium clusters; *X*-ray crystal structure of the μ$_3$-formylmethylidene compound [Os$_3$(CHCHO)(CO)$_{10}$], 1861–4
Three isomeric hydride compounds formed by protonation of decacarbonylbis(dimethylphenylphosphine)triosmium and the *X*-ray structures of two of these: 1,1- and 1,2-[Os$_3$H(CO)$_{10}$-(PMe$_2$Ph)$_2$][PF$_6$], 1869–76
A general synthetic route for platinum cluster compounds containing carbonyl and tertiary phosphine ligands and a study of their reactions with unsaturated inorganic molecules, 1889–96
Electrophilic attack on a series of dinuclear diphosphazane-bridged derivatives of iron by halogens. *X*-Ray crystal structures of [Fe$_2$I(CO)$_5$\{μ-(MeO)$_2$PN(Et)P(OMe)$_2$\}$_2$]PF$_6$ and [Fe$_2$(μ-Br)(CO)$_4$\{μ-(PhO)$_2$PN(Et)P(OPh)$_2$\}$_2$]PF$_6$, 1933–46
The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of [M$_2$Ru$_3$(μ$_3$-S)(μ-Ph$_2$PCH$_2$PPh$_2$)-(CO)$_9$] (M = Cu or Au), 1967–76
Transformation of [PPh$_4$][Fe$_5$RhC(CO)$_{16}$] under hydroformylation conditions: synthesis and crystal structure of [PPh$_4$][Fe$_3$Rh$_3$C(CO)$_{15}$], 1983–8
Alkylation of tricarbonyliron complexes by trimethylsilyl cyanide: synthetic and kinetic studies, 2013–6
Carbon-13 nuclear magnetic resonance study of the properties of the unsaturated anion [Re$_3$(μ-H)$_4$(CO)$_9$(NCMe)]$^-$ at low temperature, 2083–6
A trigonal bipyramidal pentaruthenium cluster, [Ru$_5$(CO)$_3$(μ$_3$-CO)$_3$(η-C$_5$H$_5$)$_4$], 2087–8
Reactions of main group homopolynuclear cations with transition-metal carbonyls. Stabilisation of μ-Se$_2^{2+}$ and η^3-Te$_3^{2+}$ dicationic ligands and crystal structures of [Mo(Te$_3$)(CO)$_4$][SbF$_6$]$_2$ and [FeW(Se$_2$)(CO)$_8$][SbF$_6$]$_2$, 2111–6
Some chemistry of the heterobimetallic complexes [(OC)$_3$Mo(μ-dppm)$_2$Rh(CO)Cl] and [(OC)$_3$Mo(μ-dppm)$_2$-M(CO)$_2$]PF$_6$ (M = Rh or Ir, dppm = Ph$_2$PCH$_2$PPh$_2$). Structure of [(OC)$_3$Mo(μ-dppm)$_2$(μ-C≡CMe)Rh(CO)], 2171–6
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 62. Synthesis of penta-, hexa-, and hepta-heteronuclear metal cluster compounds involving tungsten or molybdenum with platinum or nickel: crystal structures of the chain compounds [Pt$_3$W$_3$(μ-CMe)(μ$_3$-CMe)$_2$(CO)$_6$(cod)(η-C$_5$Me$_5$)$_3$]·0.5CH$_2$Cl$_2$ and [Pt$_3$W$_4$(μ-CC$_6$H$_4$Me-4)$_2$(μ$_3$-CC$_6$H$_4$Me-4)$_2$(CO)$_8$(η-C$_5$H$_5$)$_4$], 2177–88
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 63. Synthesis of eight-membered-ring metallacycles: *X*-ray crystal structures of [Pt$_4$W$_4$(μ-CR)(μ$_3$-CR)$_3$(μ-CO)(CO)$_7$(η-C$_5$H$_5$)$_4$]·CH$_2$Cl$_2$, [Ni$_2$Pt$_2$W$_4$(μ-CR)(μ$_3$-CR)$_3$(μ-CO)(CO)$_7$(η-C$_5$H$_5$)$_4$]·CH$_2$Cl$_2$ (R = C$_6$H$_4$Me-4), and [Ni$_2$Pt$_2$W$_4$(μ$_3$-CPh)$_4$(CO)$_8$(η-C$_5$H$_5$)$_4$], 2189–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 64. Addition of methylene groups to iron–molybdenum complexes; crystal structures of [FeMo(μ-CH$_2$)\{μ-σ:η-C(C$_6$H$_4$Me-4)=CH$_2$\}(CO)$_5$(η-C$_5$H$_5$)] and [FeMo\{μ-C(C$_6$H$_4$Me-4)C(OMe)C(H)\}(CO)$_5$(η-C$_5$H$_5$)], 2201–10
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 65. Reactions of the complexes [FeW(μ-CR)(CO)$_5$(η-C$_5$Me$_5$)] and [Fe$_2$W(μ$_3$-CR)(μ-CO)(CO)$_8$(η-C$_5$Me$_5$)] (R = C$_6$H$_4$Me-4) with alkynes R'C$_2$R' (R' = Me or Ph); crystal structures of [FeW\{μ-C(R)C(O)C(Me)C(Me)\}(CO)$_5$-(η-C$_5$Me$_5$)]·CH$_2$Cl$_2$ and [FeW\{μ-C(R)C(Et)C(H)C(Me)C(Me)\}-(μ-CO)(CO)$_3$(η-C$_5$Me$_5$)], 2211–8
A reversible metal framework rearrangement in high-nuclearity osmium-platinum cluster compounds. *X*-Ray crystal structures of [Os$_6$Pt(CO)$_{17}$(μ$_3$-NCMe)(C$_8$H$_{12}$)], [Os$_6$Pt(CO)$_{17}$(μ$_4$-NCMe)(C$_8$H$_{12}$)](C$_8$H$_{12}$ = cyclo-octa-1,5-diene), and [Os$_6$(CO)$_{19}$(μ$_3$-NCMe)]: compounds with novel metal–acetonitrile bonding modes, 2245–52
Synthesis, reactivity, and *X*-ray crystal structures of [Os$_6$Pt$_2$H$_2$(CO)$_{16}$(μ-C$_8$H$_{10}$)(C$_8$H$_{12}$)]·CH$_2$Cl$_2$ and [Os$_6$H(CO)$_{16}$(μ-C$_8$H$_{11}$)]. Model compounds for C–H activation, 2253–60
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC$_6$H$_4$) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as [Re(CO)$_2$(PPh$_3$)$_2$\{S=C(NHPh)-S\}]·[Re(CO)$_2$(PPh$_3$)$_2$\{O–C(NHPh)=S\}], 2299–304
Determination of the relative signs of $^2J(^{31}P–^{31}P)$ in complexes of tungsten(0) and molybdenum(0) using two-dimensional [^{31}P–^{31}P]-COSY-45 nuclear magnetic resonance chemical shift correlation, 2401–6
Substitution reactions of the osmium(I) dinuclear compound [Os$_2$(MeCO$_2$)$_2$(CO)$_6$]. *X*-Ray crystal structure of [N(PPh$_3$)$_2$][Os$_2$(MeCO$_2$)$_2$Cl(CO)$_5$], 2473–8
Metallo-ligands containing terminal oxygen donors bonded to a manganese–carbonyl moiety, 2599–604
The reaction of [Mn$_2$\{σ:η^2-C(R)=CHR\}(μ-PPh$_2$)(CO)$_7$] (R = H or Ph) with nucleophiles: *X*-ray crystal structures of [N(PPh$_3$)$_2$][Mn$_2$(μ-H)\{μ-C(O)C(Ph)=CHPh\}(μ-PPh$_2$)(CO)$_6$], [Mn$_2$\{PPh$_2$C(Ph)=CHPh\}(CO)$_9$], [Mn$_2$(μ-PPh$_2$-CH=CH$_2$)(CO)$_8$], [Mn$_2$\{μ-C(O)C(Ph)=CHPh\}(μ-PPh$_2$)(CO)$_6$(CNBu^t)$_2$], [Mn$_2$\{μ-σ:η^2-C(Ph)=CHPh\}(μ-PPh$_2$)(CO)$_6$(CNBu^t)], and [Mn(μ-PPh$_2$CH=CH$_2$)(CO)$_7$(PEt$_3$)], 2669–82
Heteronuclear complexes containing Group 6 transition metals; chemical, spectroscopic, and theoretical studies on some binuclear complexes and the *X*-ray crystal structure determination of [WNi(CO)$_3$(PPh$_3$)$_2$(η^2-C$_5$H$_5$)], 2741–50
Rhodium(I)–iron(0) carbonyl complexes containing one bridging Ph$_2$PCH$_2$PPh$_2$ ligand, including the formation of a tetranuclear cluster from a heterobimetallic precursor: crystal structures of [(OC)$_4$Fe(μ-Ph$_2$PCH$_2$PPh$_2$)RhCl(CO)] and [Fe$_2$Rh$_2$(μ-Ph$_2$PCH$_2$PPh$_2$)$_2$(CO)$_8$], 2751–6
Reactivity of a cluster-co-ordinated acyl. Reduction of a μ-acetyl ligand to acetaldehyde, 2839–42
σ-Alkynyl complexes of manganese(I) as η^2-bonding ligands for Group 1B metal–ligand fragments. *X*-Ray crystal structure of [Mn$_2$Cu(μ-CCBu^t)$_2$(CO)$_6$(dppe)$_2$]PF$_6$·0.5CH$_2$Cl$_2$, 2867–74
Manganese(I) complexes with (tricyclohexylphosphonio)dithiocarboxylate as chelate and unidentate ligand. *X*-Ray crystal structure of *fac*-[Mn(CO)$_3$\{S$_2$CP(C$_6$H$_{11}$)$_3$\}$_2$]ClO$_4$·H$_2$O, 2875–80
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 68. Reactions between nonacarbonyldi-iron and the salts [NEt$_4$][W(≡CR)(CO)$_2$(η^5-C$_2$B$_9$H$_9$Me$_2$)] (R = Me, Ph, or C$_6$H$_4$Me-4); crystal structures of [NEt$_4$][FeW\{μ-CH(C$_6$H$_4$Me-4)\}(μ-σ:η^5-C$_2$B$_9$H$_8$Me$_2$)(μ-CO)(CO)$_5$] and [NEt$_4$][FeW(μ$_3$-CPh)(μ-σ:σ',η^5-C$_2$B$_9$H$_7$Me$_2$)(CO)$_8$], 2917–26
Dienophile cycloaddition to cycloheptatriene and related complexes of tricarbonyliron. *X*-Ray crystal structures of [Fe(CO)$_3$(η^4-C$_7$H$_7$CN)]·C$_2$(CN)$_4$ and [Fe(CO)$_3$(η^4-C$_7$H$_7$–C$_7$H$_7$)], 2927–34
Reactions of allene with triosmium and triruthenium clusters including coupling and oxidative addition: *X*-ray structure of the coupled allene compound [Os$_3$(C$_6$H$_8$)(CO)$_{10}$], 2935–40
Synthesis and chemistry of *trans*-[RhX(CO)L$_2$] (X = anionic ligand, L = tertiary phosphine), 2969–74
Preparation, isomerization, and reactions of hydride complexes of ruthenium(II), 2989–94
Instant ligands. Part 2. Preparation of some bidentate ligands containing PF$_2$O Groups from aromatic hydroxy compounds and their reactions to form molybdenum and tungsten complexes, 3023–8
Chelate ring-opening of rhenium–diphosphine complexes by diazonium salts, 3067–72

CARBONYL (contd)
Platinum(II)– and palladium(II)–iron(0) complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand. Crystal structures of $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}CO)PtBr_2]$ and $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}I)PtI]\cdot CH_2Cl_2$, 3079–84
Preparation of two mixed-metal carb· ·yl clusters based on the platinumtriosmium core. *X*-Ray crystal structure of $[Os_3PtH_2(cod)(CH_2)(CO)_9]$ (cod = cyclo-octa-1,5-diene), 3103–6
Reactivity of rhenium(I) pseudo-allyl complexes with hererocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S{=}C(NHPh)\text{–}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O\text{–}C(NHPh){=}S\}]$ (1987, 2299), 3107–8
CARBOXYLATE
Organotin biocides. Part 8. The crystal structure of triphenyltin formate and a comparative variable-temperature tin-119 Mössbauer spectroscopic study of organotin formates and acetates, 101–6
Equilibria in aqueous solutions of some iron(II) complexes, 255–8
Studies on transition-metal peroxo complexes. Part 7. Molybdenum(VI) and tungsten(VI) carboxylato peroxo complexes, and the *X*-ray crystal structure of $K_2[MoO(O_2)_2(glyc)]\cdot 2H_2O$, 991–6
Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNN'N'*-tetra-acetate (cdta): structural and magnetic characterization of $[(H_2O)_4Cu(cdta)Ni]\cdot 3H_2O$ and $[(H_2O)_5Ni(cdta)Cu]\cdot H_2O$, 1847–52
Preparation and characterisation of dicarboxylatobis(η-cyclopentadienyl)tungsten(IV) complexes: a novel and convenient synthesis from $[W(\eta\text{-}C_5H_5)_2H_2]$ using molecular oxygen as a hydrogen acceptor, 1857–60
Reactions of tetra-μ-carboxylato-diruthenium(II,II) compounds. *X*-Ray crystal structures of $Ru_2(\mu\text{-}O_2CCF_3)_4(thf)_2$, $Ru_2(\mu\text{-}O_2CR)_4(NO)_2$ (R = Et or CF_3), and $\{Na_3[Ru_2(\mu\text{-}O_2CO)_4]\cdot 6H_2O\}_n$, 2723–36
CARBOXYPENTONIC ACID
Crystal structure of a copper complex of 2-carboxypentonic acid; a decomposition product of dehydroascorbic acid, 2905–10
CATALASE
A realistic model for heme-containing catalases and peroxidases: the *X*-ray structural characterisation of a non-porphyrin iron(III) macrocyclic complex, and the mechanism of its peroxidation of aromatic substrates, 2505–10
CATALYSIS
Synthesis of $[Mo(\eta\text{-}C_5H_4Pr^i)(PR_3)_2H_3](R_3 = Me_3$ or $Me_2Ph)$, $[Mo(\eta\text{-}C_5H_4Pr^i)(Pr^i{}_2PCH_2CH_2PPr^i{}_2)H_3]$, and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of $[\{Mo(\eta\text{-}C_5H_4Pr^i)(\mu\text{-}Cl)_2\}_2]$, 9–20
Synthesis of the alkoxo(hydrido)-clusters $[Ru_3(\mu\text{-}H)(CO)_{10}(\mu\text{-}OR)]$ [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid (1987, 573), 573–8
Photo-oxidations of acetoin and biacetyl catalysed by tetrabutylammonium decatungstate(4−) in acetonitrile under excess of oxygen, 961–2
Water photolysis. Part 2. An investigation of the relative advantages of various components of the sensitiser–electron relay–metal colloid system for the photoproduction of hydrogen from water, and the use of these systems in the photohydrogenation of unsaturated organic substrates, 1299–304
Electrocatalytic reduction of carbon dioxide to methanol. Part 5. Relationship between the ability of metal complexes to engage in homogeneous catalysis and their co-ordination chemistry, 1377–80
Di-2-pyridyl ketone complexes of copper(I): efficient photocatalysts for norbornadiene–quadricyclane conversion, 1797–800
Transformation of $[PPh_4][Fe_5RhC(CO)_{16}]$ under hydroformylation conditions: synthesis and crystal structure of $[PPh_4][Fe_3Rh_3C(CO)_{15}]$, 1983–8
Photochemical reduction of carbon dioxide to carbon monoxide in water using a nickel(II) tetra-azamacrocycle complex as catalyst, 2105–10
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid (1987, 573), 2291–2
The formation and characterisation of (η^2-ethene)hydrido(η^5-pentamethylcyclopentadienyl)(trisubstituted-silyl)rhodium complexes; intermediates in catalytic dehydrogenative silylation reactions, 2709–14
CERIUM
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid, 573–8
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid (1987, 573), 2291–2
CHAIN
Linear-chain platinum(II) dithiocarboxylates: crystal structure of tetrakis(phenyldithioacetato)diplatinum(II), 35–40
Synthesis and properties of heterotrimetallic complexes containing M–Pt–Hg (M = Mo, W, Fe, Mn, or Co) arrays, 119–22
CHARGE DENSITY
Charge density in the $CoCl_4^{2-}$ ion: a comparison with spin density and theoretical calculations, 1737–46
Covalence in the $[Cr(CN)_6]^{3-}$ ion studied by *X*-ray diffraction from $Cs_2KCr(CN)_6$, 1747–54
CHIRALITY
Amino-acid complexation to a cobalt(III) complex of optically-active (2*R*,5*R*,8*R*,11*R*)-2,5,8,11-tetraethyl-1,4,7,10-tetra-azacyclododecane (L). Preparation, characterization, and structure determinations of $[Co^{III}\{(R)\text{- or }(S)\text{-alaO}\}L]Br[ClO_4]\cdot H_2O$, 721–8
Synthesis and stereochemical studies of the chiral ruthenium complexes $[Ru(\eta\text{-}C_5H_5)\{(S)dpompyr\text{-}PP'\}X]$ [dpompyr = *N*-diphenylphosphino-2-(diphenylphosphinoxymethyl)pyrrolidine, X = H or Cl]. Crystal structure of $[(S)Ru(\eta\text{-}C_5H_5)\{(S)dpompyr\text{-}PP'\}Cl]$, 1149–56
The chemistry of tetraphenylcyclopentadienone complexes of ruthenium and rhodium: the *X*-ray crystal structure of $[Ru\{\eta^5\text{-}C_5Ph_4OC(O)CH(OMe)Ph\}(CO)_2Cl]$, 2815–22
CHLORIDE
Crystal and molecular structure of the six-co-ordinate antimony(III) complex $SbCl_3L$, where L = *N*,*N'*-dimethyloxamide. Unusual O,O ligand binding leading to four-membered $Sb(\mu\text{-}O)_2Sb$ ring formation, 263–6
Trimethylphosphine complexes of niobium(IV) and tantalum(IV). Crystal, molecular, and electronic structures of $[M_2Cl_4(\mu\text{-}Cl)_4(PMe_3)_4]$ (M = Nb or Ta), 307–14
Matrix isolation and mass spectrometric studies on the vaporisation of chromium(III) chloride. The characterisation of molecular $CrCl_4$ and $CrCl_3$, 859–66
CHLOROPHENYL
The first authenticated example of geometrical isomers in organotin(IV) co-ordination chemistry. The crystal and molecular structures of octahedral *cis*- and *trans*-SnR_2 isomeric adducts of dichlorobis(4-chlorophenyl)tin(IV) with 4,4'-dimethyl-2,2'-bipyridyl, 129–38
CHROMATOGRAPHY
Chiral recognition in ion pairing of an optically active tris(ethylenediamine)cobalt(III) cation and application to chromatographic resolution of metal complex anions, 1977–82
CHROMIUM
Magnetic and electrochemical studies on the series $[Cr_nFe_{3-n}(\mu_3\text{-}O)(gly)_6(H_2O)_3][NO_3]_{7-}\cdot xH_2O$(gly = glycine; *n* = 0, 1, 2, or 3) and on the mixed-valence complex $[Fe^{II}Fe^{III}{}_2(\mu_3\text{-}O)(gly)_6(H_2O)_3]Cl_6$, 97–100
Bimetallic systems. Part 16. Formation of Group 6 metal(0)–platinum(II) hydride complexes, by ring opening of *fac*- or *mer*-$[M(CO)_3(dppm\text{-}PP')(dppm\text{-}P)]$ (M = Cr, Mo, or W; dppm = $Ph_2PCH_2PPh_2$). Reactions of $[(CO)_3Mo(\mu\text{-}dppm)_2PtH(Cl)]$, 221–6
Kinetics and mechanism of ring closure for a series of photoproduced $[M(CO)_5L]$ intermediates (M = Cr, Mo, or W; L = pyridine 2-carbaldehyde imine ligand), 513–8
The synthesis and *X*-ray crystal structures of homoleptic tetrahedral aryls of osmium(IV) and of cyclohexyls of ruthenium(IV), osmium(IV), and chromium(IV), 557–62
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid, 573–8
Heteronuclear 1,3-intramolecular metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of 2-thia-4-selenapentane. A dynamic nuclear magnetic resonance investigation, 757–60
Matrix isolation and mass spectrometric studies on the vaporisation of chromium(III) chloride. The characterisation of molecular $CrCl_4$ and $CrCl_3$, 859–66

CHROMIUM (contd)
Ferromagnetic tetrahalogenochromates(II) containing mixed halides (Cl and Br), 965–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 59. Reactions of the alkylidyne compounds $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Cr, Mo, or W, R = C_6H_4Me-4; M = W, R = Me, $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate] with d^{10} complexes; crystal structure of $[PtW_2(\mu\text{-}CMe)_2(CO)_4\{HB(pz)_3\}_2]$, 1229–34
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 60. Di- and tri-metal compounds prepared from the metal alkylidyne complexes $[W(\equiv CR)(CO)_2(\eta\text{-}C_5Me_5)]$ (R = Me or C_6H_4Me-4) and $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Mo, R = C_6H_4Me-4; M = W, R = Me; $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate], 1235–42
Synthesis, characterization, and reactions of chromyl(VI) alkoxides, 1253–6
Reaction paths and kinetics in the Cr^{2+} reduction of the Mo^{V}_2 and Mo^{IV}_3 ions $[Mo_2O_4(H_2O)_6]^{2+}$ and $[Mo_3O_4(H_2O)_9]^{4+}$, 1397–402
Reactions of chromium(III) complexes of 1,10-phenanthroline, 2,2′-bipyridyl, and oxalate with the pulse radiolytically generated aquated electron, zinc(I), and cadmium(I), 1425–30
Bimetallic systems. Part 19. Chromium(0)–, molybdenum(0)–, or tungsten(0)–iron(0) carbonyl complexes containing one or two bridging $Ph_2PCH_2PPh_2$ ligands: crystal structure of $[(OC)_3Mo(\mu\text{-}Ph_2PCH_2PPh_2)_2Fe(CO)_3]$, 1489–94
Bimetallic systems. Part 20. Synthesis of Group 6 metal(0)–iridium(I) or –rhodium(I) complexes bridged by $Ph_2PCH_2PPh_2$, 1495–500
The triphenylphosphine cone angle and restricted rotation about the chromium–phosphorus bond in dicarbonyl(η^6-hexaalkylbenzene)(triphenylphosphine)chromium(0) complexes. Crystal and molecular structure of dicarbonyl(η^6-hexa-n-propylbenzene)(triphenylphosphine)chromium(0), 1545–50
Covalence in the $[Cr(CN)_6]^{3-}$ ion studied by *X*-ray diffraction from $Cs_2KCr(CN)_6$, 1747–54
Crystal and molecular structure of chromium(II) sulphate pentahydrate and single-crystal electronic spectra and bonding of $CrSO_4{\cdot}5H_2O$, $CuSO_4{\cdot}5H_2O$, and $CuSO_4{\cdot}5D_2O$, 1817–22
Synthesis, characterisation, and aquation kinetics of [*N*-(*o*-carboxyphenyl)iminodiacetato](glycinato)chromate(III) complex ion in acidic media, 2095–8
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid (1987, 573), 2291–2
Kinetics of reduction of tris(ethylenediamine)cobalt(III) by ethylenediamine-*NNN′N′*-tetra-acetatochromate(II), 2485–8
Dinitrogen complexes derived from tricarbonyl(η^6-hexaethylbenzene)chromium(0): crystal and molecular structure of μ-dinitrogen-bis[dicarbonyl(η^6-hexaethylbenzene)-chromium(0)]–toluene (1/1), 2789–92
Carbene complexes. Part 21. Synthesis and characterisation of bis(carbene)molybdenum(II) complexes and dimetal(0) complexes of the Group 6 elements containing novel bridging bis(carbene) ligands; *X*-ray structures of $[Mo(CO)_2(L^{Et})_2(OSO_2CF_3)_2]$ $[L^{Et} = \equiv\overline{CN(Et)(CH_2)_2}NEt]$ and $[\overline{W(CO)_5\{C(OEt)CH_2\text{-}C_6H_4CH_2C(OEt)\text{-}o\}W(CO)_5}]$, 2843–52
The chemistry of chromium nitrile complexes of 1,2-bis(dimethylphosphino)ethane. *X*-Ray crystal structures of *trans*-$[Cr^{IV}Cl(NEt)(dmpe)_2]CF_3SO_3$, *trans*-$[Cr^{IV}(N{=}CHMe)_2(dmpe)_2]$-$[BPh_4]_2$, and *trans*-$[Co^{0}(H{\cdot}NCMe)(dmpe)_2][BPh_4]_2$, 2947–54

CIRCULAR DICHROISM
Reactions of copper(II) complexes of optically active *N*-substituted diamines with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of optically active macrocycles, 1127–32
On the induced circular dichroism of *d–d* transitions of a 14-membered macrocyclic tetra-amine nickel(II) complex in β-cyclodextrin, 1801–4

CLATHRATE
Mössbauer spectroscopic studies on molecular rotation and lattice dynamics of cyclodextrin clathrates of bis(η-cyclopentadienyl)-iron(II) and some of its ring-substituted derivatives, 627–32
Studies in Werner clathrates. Part 4. Structures of tetrakis(4-ethylpyridine)di-isothiocyanatonickel(II) and its clathrates with *p*-, *m*-, and *o*-xylene, carbon disulphide, and carbon tetrachloride, 2125–40

CLUSTER
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 49. Synthesis of mixed-metal complexes with bonds between Cr, Mo or W and Co, Rh, Ir or Re; crystal structure of $[CrReRh_2(\mu_3\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}CO)(CO)_9(\mu\text{-}C_9H_7)_2]$, 41–50
High-yield syntheses of sulphido triruthenium carbonyl cluster complexes containing bis(diphenylphosphino)methane, 91–6
Synthesis of the alkoxo(hydrido)-clusters $[Ru_3(\mu\text{-}H)(CO)_{10}(\mu\text{-}OR)]$ [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4
Synthesis and *X*-ray crystal structure of a cuboctahedral platinum cluster, 259–62
Reactions of metal carbonyl cluster complexes with multidentate phosphine ligands; the preparation and thermolysis of bis(dimethylphosphino)methane derivatives of $[Ru_3(CO)_{12}]$, and the *X*-ray crystal structure of $[\overline{Ru_3(CO)_9(H)(Me_2PCHPMe_2)}]$, 277–80
Preparations, crystal structures, and molybdenum-95 nuclear magnetic resonance spectroscopic studies of triphenylphosphine–gold–tetrathiomolybdate(VI) clusters, 303–6
Reaction of $[Os_5H_2(CO)_{15}]$ with pyridine: crystal and molecular structures of $[Os_5H_3(CO)_{14}(C_5H_4N)]$ and $[Os_5H_2(CO)_{14}(C_5H_5N)]$, 327–34
Thermal and photochemical behaviour of organotetraruthenium clusters: solution structures and dynamics of phosphine-substituted derivatives, 349–54
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au), 379–90
Determination of the deuterium position in DNb_6I_{11}, 431–4
Triosmium clusters containing ligands derived from tropone (cycloheptatrienone): molecular structure of an oxidative addition product $[Os_3H(\mu_3\text{-}C_7H_5O)(CO)_9]$, 551–6
A group theoretical paradigm for describing the skeletal molecular orbitals of cluster compounds. Part 1. Deltahedral clusters, 647–56
Vibrational studies of interstitial hydrogen in metal carbonyl clusters, 685–90
Synthesis and characterisation of $[N(PPh_3)_2][RuRh_5(CO)_{16}]$ by *X*-ray crystallography and by natural-abundance carbon-13 nuclear magnetic resonance spectroscopy, 781–4
Redox behaviour of the iron–sulphur cluster $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{2+}$. Synthesis and crystal structure of the new paramagnetic monopositive species $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{+}$ as its $[PF_6]^{-}$ salt, 831–6
Trinuclear osmium clusters, containing bidentate $Ph_2PCH_2PPh_2$ and $Me_2PCH_2PMe_2$ ligands: *X*-ray crystal structures of the complexes $[Os_3H(CO)_9(Me_2PCHPMe_2)]$ and $[Os_3H(OH)(CO)_8(Ph_2PCH_2PPh_2)]$, 931–8
The heteronuclear cluster chemistry of the Group 1B metals. Part 3. Synthesis, structures, and chemical properties of the trimetallic hexanuclear Group 1B metal cluster compounds $[MM'Ru_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]$ (M = Cu, M′ = Ag or Au; M = Ag, M′ = Au). *X*-Ray crystal structure of the complex $[AgCuRu_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]{\cdot}CH_2Cl_2$, 1001–8
Extended Hückel molecular orbital study of the effects of edge-bridging hydrogen atoms on the lengths of boron–boron and metal–metal bonds in cluster compounds, and the crystal structure of benzyltrimethylammonium octahydrotriborate(1−), 1017–26
High nuclearity osmium cluster radicals, 1051–4
The ring-stacking principle in organolithium chemistry: its development through the isolation and crystal structures of hexameric iminolithium clusters $(RR'C{=}NLi)_6$ (R′ = Ph, R = Bu^t or Me_2N; R = R′ = Me_2N or Bu^t), 1071–82
Carbon-13 and phosphorus-31 nuclear magnetic resonance studies of the fluxional behaviour of the complexes $[Fe_3(CO)_{10}\{P(OMe)_3\}_2]$ and $[Fe_3(CO)_{10}\{P(OMe)_3\}\{P(OPh)_3\}]$, 1249–52
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au) (1987, 379), 1283–4
The preparation, characterisation, and some reactions of $[Os_3H(X)(CO)_9(NMe_3)]$ (X = Cl, Br, or I), 1285–8
Systematic cluster build up by the reaction of

CLUSTER (contd)
$[Os_3H(X)(CO)_9(MeCN)]$ (X = SR or NHMe; R = Me, Et, or Ph) with $[OsH_2(CO)_4]$, 1289–92
Substitution reactions on triosmium clusters of the type $[Os_3(\mu\text{-}H)(\mu\text{-}X)(CO)_{10}]$ [X = NHMe, O_2CR, C_5H_4N, or OC(NHMe); R = H, Me, or Ph], 1293–8
A group theoretical paradigm for describing the skeletal molecular orbitals of cluster compounds. Part 2. Bispherical clusters, 1445–56
Triosmium clusters containing the ligands CNH_2, NCHCN, and NCMeCN derived from cyanide ion, cyanogen, and acetonitrile, 1457–62
Intermediates in the conversion of $[Os(CO)_{11}(PRPh_2)]$ (R = Me or Ph) into $[Os_3(\mu_3\text{-}C_6H_4)(\mu_3\text{-}PR)(CO)_9]$: crystal and molecular structures of $[Os_3(\mu\text{-}H)(\mu_3\text{-}C_6H_4PMePh\text{-}o)(CO)_9]$ and $[Os_3(\mu_3\text{-}C_6H_4PMe\text{-}o)(CO)_{10}]$, 1529–34
Substituted osmium clusters containing labile ligands. The preparation, characterisation, and some reactions of $[Os_3H(OR)(CO)_9(MeCN)]$ R = H, Me, Et, or Ph) and the isomers of $[Os_3H(SR)(CO)_9(MeCN)]$ (R = Me, Et, or Ph); the crystal and molecular structure of $[Os_3H(SPh)(CO)_9(PEt_3)]$, 1623–30
Crystal and molecular structure of $[Co_8(CO)_{21}(\mu_5\text{-}CS_2)(\mu_5\text{-}C_2S)]$: a thiocarbonyl group as a six-electron donor, 1707–12
Decarboxylation of vinylene carbonate at triosmium clusters; *X*-ray crystal structure of the μ_3-formylmethylidene compound $[Os_3(CHCHO)(CO)_{10}]$, 1861–4
Three isomeric hydride compounds formed by protonation of decacarbonylbis(dimethylphenylphosphine)triosmium and the *X*-ray structures of two of these: 1,1- and 1,2- $[Os_3H(CO)_{10}(PMe_2Ph)_2][PF_6]$, 1869–76
A general synthetic route for platinum cluster compounds containing carbonyl and tertiary phosphine ligands and a study of their reactions with unsaturated inorganic molecules, 1889–96
The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of $[M_2Ru_3(\mu_3\text{-}S)(\mu\text{-}Ph_2PCH_2PPh_2)(CO)_9]$ (M = Cu or Au), 1967–76
Transformation of $[PPh_4][Fe_5RhC(CO)_{16}]$ under hydroformylation conditions: synthesis and crystal structure of $[PPh_4][Fe_3Rh_3C(CO)_{15}]$, 1983–8
Carbon-13 nuclear magnetic resonance study of the properties of the unsaturated anion $[Re_3(\mu\text{-}H)_4(CO)_9(NCMe)]^-$ at low temperature, 2083–6
A trigonal bipyramidal pentaruthenium cluster, $[Ru_5(CO)_3(\mu_3\text{-}CO)_3(\eta\text{-}C_5H_5)_4]$, 2087–8
A reversible metal framework rearrangement in high-nuclearity osmium-platinum cluster compounds. *X*-Ray crystal structures of $[Os_6Pt(CO)_{17}(\mu_3\text{-}NCMe)(C_8H_{12})]$, $[Os_6Pt(CO)_{17}(\mu_4\text{-}NCMe)(C_8H_{12})]$ (C_8H_{12} = cyclo-octa-1,5-diene), and $[Os_6(CO)_{19}(\mu_3\text{-}NCMe)]$: compounds with novel metal–acetonitrile bonding modes, 2245–52
Synthesis, reactivity, and *X*-ray crystal structures of $[Os_6Pt_2H_2(CO)_{16}(\mu\text{-}C_8H_{10})(C_6H_{12})]\cdot CH_2Cl_2$ and $[Os_6H(CO)_{16}(\mu\text{-}C_8H_{11})]$. Model compounds for C–H activation, 2253–60
The synthesis, crystal and molecular structure of a copper(I) complex of 2(1*H*)-pyridinethione: an intermediate in Cu_4S_4 heterocubane build up, 2281–4
Gold-197 Mössbauer spectra of some gold–ruthenium cluster compounds, 2333–6
Reactivity of a cluster-co-ordinated acyl. Reduction of a μ-acetyl ligand to acetaldehyde, 2839–42
Reactions of allene with triosmium and triruthenium clusters including coupling and oxidative addition: *X*-ray structure of the coupled allene compound $[Os_3(C_6H_8)(CO)_{10}]$, 2935–40
Preparation of two mixed-metal carbonyl clusters based on the platinumtriosmium core. *X*-Ray crystal structure of $[Os_3PtH_2(cod)(CH_2)(CO)_9]$ (cod = cyclo-octa-1,5-diene), 3103–6

COBALAMIN
The chemistry of vitamin B_{12}. Part 27. The cage phosphite ester 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane and the anionic dimethyl phosphite as ligands for cobalt(III) corrinoids, 761–8

COBALT
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 5. Unsubstituted fifteen-membered-ring macrocycles containing 2,2′:6′,2″-terpyridine, 5–8
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution, 435–42
Reduction–oxidation properties of organotransition-metal complexes. Part 25. The one-electron reduction of cationic cyclopentadienyl–cobalt and –rhodium nitrosyl derivatives, 467–72
Magnetic exchange interactions in perovskite solid solutions. Part 7. A study of the non-stoicheiometric phase $SrCoO_{3-y}$ by ^{57}Fe Mössbauer spectroscopy, 667–72
Vibrational studies of interstitial hydrogen in metal carbonyl clusters, 685–90
Amino-acid complexation to a cobalt(III) complex of optically-active (2*R*,5*R*,8*R*,11*R*)-2,5,8,11-tetraethyl-1,4,7,10-tetra-azacyclododecane (L). Preparation, characterization, and structure determinations of $[Co^{III}\{(R)\text{- or }(S)\text{-alaO}\}L]Br\text{-}[ClO_4]\cdot H_2O$, 721–8
The chemistry of vitamin B_{12}. Part 27. The cage phosphite ester 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane and the anionic dimethyl phosphite as ligands for cobalt(III) corrinoids, 761–8
Rational synthesis of metalla–sulphur–nitrogen complexes: use of $[Me_2SnS_2N_2]_2$ in the preparation of complexes of the type $[M(S_2N_2H)(PR_3)_2]X$ (M = Pt or Pd). *X*-Ray structures of $[Pt(S_2N_2H)(PEt_3)_2][Me_2SnCl_3]$, $[Pt(S_2N_2H)(PBu^n_3)_2]_2[PF_6]Cl$, and $[\{Co(S_2N_2H)_2\}_2\{Ph_2P(O)CH_2CH_2P(O)Ph_2\}]$, 907–14
Electron spin resonance and optical spectroscopic studies on Cu^{2+} and Co^{2+} in $Cs_2[Zn(N_3)_4]$, 939–42
Direct synthesis of alkali trifluorocobaltate(II) monohydrates, $A[CoF_3]\cdot H_2O$ (A = NH_4, Na, or K). First synthesis and structural assessment of mixed-ligand fluoro complexes of cobalt(II), $[Co(NH_3)_4F_2]$ and $A[CoF(SO_4)(H_2O)_2]$ (A = NH_4, Na, or K), 1055–8
The relative nucleophilic efficiency of water and the azide and thiocyanate ions in the aqueous substitution of chloropentacyanocobaltate(III), 1059–62
Photoactivation of methane by η^5-cyclopentadienyl and substituted η^5-cyclopentadienyl Group 8 metal dicarbonyl complexes, $[M(\eta^5\text{-}C_5R_5)(CO)_2]$ (M = Rh or Ir, R = H or Me), and dicarbonyl(η^5-indenyl)iridium: a matrix isolation study, 1181–90
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 54. Compounds derived from $[W(\equiv CC_6H_4Me\text{-}4)(CO)_2(\eta\text{-}C_5Me_5)]$ with bonds between tungsten and iron, cobalt, or nickel; crystal structure of $[FeW\{\mu\text{-}C(C_6H_4Me\text{-}4)C(O)O\}(CO)_5(\eta\text{-}C_5Me_5)]$, 1191–200
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1283–4
Mixed oxidation state trinuclear cobalt complexes with bridging sulphito and Schiff-base ligands. Part 1. Preparation of the complexes $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}L)_2Co^{III}_2(ROH)_2]$ (L = Schiff base anion, R = alkyl) and structure determination of $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}\alpha,\alpha'\text{-}Me_2\text{-}salpd)_2Co^{III}_2(Pr^nOH)_2]\cdot 2Pr^nOH$, 1305–12
The chemistry of vitamin B_{12}. Part 28. Crystal structure of dicyanocobyrinic acid heptamethyl ester and its interaction with alcohols: the effects of hydrogen bonding to co-ordinated cyanide, 1349–58
Reaction of elemental sulphur with cobalt(II)–Schiff-base complexes: synthesis of μ-disulphido and μ-tetrasulphido binuclear cobalt(III) complexes. Crystal structures of two binuclear cobalt(III)–Schiff-base complexes, 1367–76
Magnetic exchange interactions in perovskite solid solutions. Part 8. A study of the solid solution $SrFe_{1-x}Co_xO_{3-y}$ by iron-57 Mössbauer spectroscopy, 1419–24
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1583–4
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), and hydrogen ions in aqueous solution (1987, 435), 1583–4
Synthesis and characterization of new heterobimetallic iron–cobalt complexes with ethenyl bridges $CR^2{=}CHR^1$ [R^1 = H or C(O)OMe, R^2 = C(O)OMe or C(O)OEt]. *X*-Ray structure of $[FeCo(CO)_6\{\mu\text{-}MeOC(O)C{=}C(H)C(O)OMe\}]$, 1619–22
Crystal and molecular structure of $[Co_8(CO)_{21}(\mu_5\text{-}CS_2)(\mu_5\text{-}C_2S)]$: a

COBALT (contd)
thiocarbonyl group as a six-electron donor, 1707–12
Charge density in the $CoCl_4^{2-}$ ion: a comparison with spin density and theoretical calculations, 1737–46
The base catalysed hydrolysis of *trans*-dichloro-bis(1,2-diaminoethane)cobalt(III) and -(1,3-diaminopropane)cobalt(III) cations: rate-limiting proton transfer, 1775–8
Co-oligomerization of diphenylketene and ethylene promoted by (η^5-cyclopentadienyl)bis(ethylene)cobalt(I): synthesis and *X*-ray structure of a heterocycle containing cobalt(III), 1789–92
Chiral recognition in ion pairing of an optically active tris(ethylenediamine)cobalt(III) cation and application to chromatographic resolution of metal complex anions, 1977–82
The relative nucleophilic efficiency of water and dimethyl sulphoxide in the aqueous substitution of aquapentacyano-cobaltate(III) and chloropentacyanocobaltate(III), 2311–6
An electron spin resonance study of the dioxygen adduct of bis[2-(2′-pyridylmethylenehydrazonomethyl)phenolato]cobalt(II) and related materials, 2341–6
A structural comparison of the two polymorphs of $Sr_2Co_2O_5$ by cobalt *K*-edge extended *X*-ray absorption fine structure spectroscopy, 2359–64
Kinetics of reduction of tris(ethylenediamine)cobalt(III) by ethylenediamine-*NNN′N′*-tetra-acetatochromate(II), 2485–8
Disproportionation, electrochemistry, and electronic coupling involving mononuclear and binuclear cobalt phthalocyanine derivatives, 2511–8
Studies of metal-ion recognition. The interaction of Co^{II}, Ni^{II}, and Cu^{II} with new oxygen–nitrogen donor macrocycles; *X*-ray structures of complexes of Cu^{II} and Ni^{II} with a 15-membered O_2N_3 derivative, 2543–54
Kinetic studies on 1:1 electron-transfer reactions involving blue copper proteins. Part 14. Reactions of poplar plastocyanin with inorganic complexes, 2573–8
Complexes of cobalt(II) and nickel(II) with 3- and 4-acetylpyridine, 2595–8
Isometallocenes. Part 1. Polymethylated isocobaltocenium salts; a general route to $[Co(C_4Me_4)(arene)]$, 2757–60
Electrolytic behaviour of transition metal bromides in acetonitrile solution, 2863–6
Structure and electrochemical properties of bis(pyridine)cobalt(III) complexes of *N,N′*-ethylenebis(salicylideneimine) homologues with an aryl substituent on the ethylene backbone. Crystal structures of $[Co(salphen)(py)_2]PF_6 \cdot 0.5Et_2O$ and $[Co(salcppn)(py)_2]PF_6 \cdot 2py$ (py = pyridine), 3055–60
An electron spin resonance study of frozen aqueous solutions containing 5,10,15,20-tetrakis(*N*-methyl-4′-pyridinio)-porphyrinatocobalt(II), 3099–102

CONDUCTIVITY
Linear-chain platinum(II) dithiocarboxylates: crystal structure of tetrakis(phenyldithioacetato)diplatinum(II), 35–40
Bis[bis(diphenylphosphino)methane]platinum(II) salts: evidence for the formation of a five-co-ordinate bis[bis(diphenylphosphino)-methane]halogenoplatinum(II) cation, 705–8
Structure and basicity. Part 10. Conductometric and potentiometric titrations of some cyclotriphosphazatrienes in non-aqueous media. The conductivities of alkali-metal acetates on titration with perchloric acid in acetic acid, 925–30
Preparation, structure, and properties of $[Ni\{S_2C_2(CF_3)_2\}_2]^-$ and $[Ni\{Se_2C_2(CF_3)_2\}_2]^-$ salts with small cations, 1683–6
Electrolytic behaviour of transition metal bromides in acetonitrile solution, 2863–6

CONFIGURATION
Menthyl-substituted organophosphorus compounds. Part 6. Preparation of P(L-men)RCl and P(L-men)R(S)Cl (R = Me, Et, Pr^i Bu^t, or Ph; men = *cyclo*-C_6H_9-2-Pr^i-5-Me) and nuclear magnetic resonance studies of configuration and halogen exchange, 795–806

CONFORMATION
Ab initio study of stable bis(carbon dioxide)molybdenum complexes, 2373–8

COPPER
Co-ordination abilities of thioamide analogues of amino acids and peptides. Copper(II) and nickel(II) complexes, 1–4
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 5. Unsubstituted fifteen-membered-ring macrocycles containing 2,2′:6′,2″-terpyridine, 5–8
Thiosemicarbazones as co-ordinating agents. Part 2. Synthesis, spectroscopic characterization, and *X*-ray structure of aquachloro(pyridoxal thiosemicarbazone)manganese(II) chloride and aqua(pyridoxal thiosemicarbazonato)copper(II) chloride monohydrate, 227–34
Syntheses, spectroscopic, and electrochemical properties of dinuclear copper(II) complexes bridged by a single thiolate sulphur atom; *X*-ray crystal structures of $[Cu_2(L^1)_2(SC_6H_4Cl\text{-}4)][PF_6]$ and $[Cu_2(L^2)_2(SC_6H_4CH_3\text{-}4)][ClO_4]$, 241–8
Magnetostructural correlations in parallel square-planar chloride bridged copper(II) dimers: structure, dynamic nuclear magnetic resonance study, and magnetic properties of $[Cu_2(terpy)_2Cl_2][PF_6]_2$, 285–92
Studies on transition-metal macrocyclic complexes. Single-crystal *X*-ray structure and electrochemistry of the bis-macrocyclic complex $[Cu(L)_2][ClO_4]_2 \cdot 2H_2O$ (L = 1,4,7-triazacyclononane), 373–8
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au), 379–90
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution, 435–42
Electron spin resonance and electronic spectra and crystal and molecular structures of copper(II) amino acid complexes, 457–66
Synthetic, spectroscopic, and *X*-ray crystallographic studies on binuclear copper(II) complexes with a tridentate NNS-bonding 2-formylpyridine thiosemicarbazone ligand. The characterization of both neutral and deprotonated co-ordinated ligand structures, 493–500
Studies of pendant-arm macrocyclic ligands. Part 5. Synthesis of two pyridine-containing penta-aza macrocycles with single pyrrolidinyl pendant arms and characterisation of their nickel(II) and copper(II) complexes. Crystal structure of perchlorato{7-[2′-(1″-pyrrolidinyl)ethyl]-3,7,11,17-tetra-azabicyclo[11.3.1]-heptadeca-1(17),13,15-triene}nickel(II) perchlorate, 545–50
A thermodynamic and spectroscopic study of the proton and copper(II) complexes of L-prolyl-L-histidine, D-prolyl-L-histidine, L-histidyl-L-histidine, and D-histidyl-L-histidine, 661–6
Bimetallic systems. Part 17. Synthesis of Group 6 metal(0)–copper(I), –silver(I), or –gold(I) complexes with two $Ph_2PCH_2PPh_2$ (dppm) bridging ligands: crystal structure of $[(OC)_2W(\mu\text{-}CO)(\mu\text{-}dppm)_2(\mu\text{-}Cl)Cu] \cdot CH_2Cl_2$, 769–76
Dicopper(II) complexes of a 28-membered N_8 macrocycle. Evidence for proton transfer from co-ordinated secondary amine to thiocyanate, leading to formation of an isothiocyanic acid complex, 873–80
Synthesis of pyridine-containing tetra-aza macrocycles: 3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), its 3,11-dibenzyl (L^2) and 3,7,11-tribenzyl (L^3) derivatives, and their nickel(II), copper(II), and zinc(II) complexes: crystal structures of $L^2 \cdot HCl$ and $[Ni(L^2)Cl]ClO_4 \cdot H_2O$, 889–94
Electron spin resonance and optical spectroscopic studies on Cu^{2+} and Co^{2+} in $Cs_2[Zn(N_3)_4]$, 939–42
Copper(II) complexes of the 14-membered *cis*-N_2S_2 macrocycle 14,14,16-trimethyl-2,6-dithia-13,17-diazatricyclo[16.4.0.0^{7,12}]-docosa-1(18),7(12),8,10,16,19,21-heptaene, 963–4
The heteronuclear cluster chemistry of the Group 1B metals. Part 3. Synthesis, structures, and chemical properties of the trimetallic hexanuclear Group 1B metal cluster compounds $[MM'Ru_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]$ (M = Cu, M′ = Ag or Au; M = Ag, M′ = Au). *X*-Ray crystal structure of the complex $[AgCuRu_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2] \cdot CH_2Cl_2$, 1001–8
Ternary metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of copper(II)–2,2′-bipyridyl– pyridoxic acid and copper(II)–2,2′-bipyridyl–pyridoxic acid γ-lactone complexes, 1009–16
Reactions of copper(II) with $[S_2P(OR)_2]^-$ (R = Et or Pr^i) and single-crystal *X*-ray studies of $Cu[S_2P(OEt)_2] \cdot bipy$ and $Cu[S_2P(OEt)_2] \cdot 2PPh_3$, 1027–34
A structural comparison of bis(quinoxaline-2,3-dithiolato)cuprate complexes, 1083–8
Lewis-base adducts of Group 11 metal(I) compounds. Part 27. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance, far-infrared, and structural studies on the mononuclear 2:1 adducts of triphenylphosphine with copper(I) and gold(I) halides, 1089–98
Lewis-base adducts of Group 11 metal(I) compounds. Part 28. Solid-state phosphorus-31 cross-polarization magic-angle

COPPER (contd)
spinning nuclear magnetic resonance and structural studies on the mononuclear 3:1 adducts of triphenylphosphine with copper(I) halides, 1099–106
Studies of the pyridine-containing tetra-azamacrocycle 3,7,11-trimethyl-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1) and its complexes with Ni^{2+}, Cu^{2+}, and Zn^{2+}. Crystal structures of the unsymmetric isomer of $[Ni(L^1)][ClO_4]_2$, and the five-co-ordinate complexes $[Ni(L^1)(dmso)][ClO_4]_2$ (dmso = dimethyl sulphoxide) and $[Ni(L^1)Cl]ClO_4$, 1107–14
Reactions of copper(II) complexes of optically active *N*-substituted diamines with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of optically active macrocycles, 1127–32
Copper(II) complexes of tyrosine-containing tripeptides, 1263–6
Binuclear copper(I) complexes containing bis(diphenylphosphino)methane bridging ligands: crystal structure of $[Cu_2(\mu\text{-dppm})_2(MeCN)_4][ClO_4]_2$, 1275–8
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au) (1987, 379), 1283–4
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1283–4
Spectroscopic, magnetic, and structural properties of diaquadimethanolbis(*N*-tosylvalinato)copper(II) and its 2,2′-bipyridine adduct, 1333–40
Design of transition metal chelates with biological activity. Potentiometric study of complex formation equilibria between 2-amino-*N*-hydroxy-3-(*p*-hydroxyphenyl)propanamide and nickel(II), copper(II), and hydrogen ions in aqueous solution, 1409–14
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1583–4
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), and hydrogen ions in aqueous solution (1987, 435), 1583–4
Synthetic, spectroscopic, and *X*-ray crystallographic studies on binuclear copper(II) complexes with a tridentate NNS-bonding 2-formylpyridine thiosemicarbazone ligand. The characterization of both neutral and deprotonated co-ordinated ligand structures (1987, 493), 1583–4
Structure and magnetism of analogous *O*-bridged (phenolato) and *S*-bridged (thiophenolato) binuclear copper(II) complexes. Crystal structure of the binuclear *S*-bridged complex μ-[2,6-bis(4′-cyclohexyl-4′-hydroxy-2′,3′-diazabuta-1′,3′-dien-1′-yl)-4-methylthiophenolato(3−)-*S*,N^2,*O*($Cu^{1,2}$)]-methanol-μ-(pyrazolyl-*N*,*N*′)-dicopper(II), 1585–92
Studies on transition-metal–peptide complexes. Part 13. Copper(II) and nickel(II) complexes of amino acids and peptides containing a thioether group, 1717–20
An electron spin resonance study of the copper(II) chelate of 2-(2′-pyridylmethylenehydrazonomethyl)phenol and related compounds, 1755–62
Di-2-pyridyl ketone complexes of copper(I): efficient photocatalysts for norbornadiene–quadricyclane conversion, 1797–800
Crystal and molecular structure of chromium(II) sulphate pentahydrate and single-crystal electronic spectra and bonding of $CrSO_4{\cdot}5H_2O$, $CuSO_4{\cdot}5H_2O$, and $CuSO_4{\cdot}5D_2O$, 1817–22
Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNN′N′*-tetra-acetate (cdta): structural and magnetic characterization of $[(H_2O)_4Cu(cdta)Ni]{\cdot}3H_2O$ and $[(H_2O)_5Ni(cdta)Cu]{\cdot}H_2O$, 1847–52
Tetranuclear amido complexes of copper(I): a synthetic and structural study, 1883–8
Dithio-oxamides as ligands: crystal structures and vibrational analyses of bis(*N*,*N*′-dicyclohexyldithio-oxamidato)palladium(II), bis(*N*,*N*′-dibenzyldithio-oxamide)copper(II) diperchlorate, and dichloro(*N*,*N*′-dimethyldithio-oxamide)zinc(II) and the carbon-13 nuclear magnetic resonance spectra of related compounds, 1921–8
The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of $[M_2Ru_3(\mu_3\text{-}S)(\mu\text{-}Ph_2PCH_2PPh_2)(CO)_9]$ (M = Cu or Au), 1967–76
Metal ion-bridged intramolecular stacking interaction between the tryptophyl residue and the aromatic heterocyclic amine within the ternary complex (1,10-phenanthroline)(L-tryptophanato-*O*,*N*)copper(II) perchlorate–water (1/2.5): an *X*-ray study, 2017–22
Lamellar inorganic ion exchangers. Proton–copper(II) ion exchange in γ-titanium bis(hydrogenphosphate), 2045–50
The synthesis, crystal and molecular structure of a copper(I) complex of 2(1*H*)-pyridinethione: an intermediate in Cu_4S_4 heterocubane build up, 2281–4
Aqua(2,2′-bipyridyl)difluorocopper(II) dihydrate: *X*-ray structure reveals short hydrogen bonds and other unusual features, 2397–400
Complexes of ligands providing endogenous bridges. Part 4. Copper(II) complexes of macrocyclic Schiff bases derived from 2,6-diacetylpyridine and 1-*n*-diamino-*n*′-hydroxyalkanes (*n*,*n*′ = 3,2; 4,2; and 5,3): synthesis, properties, and structures, 2519–30
Studies of metal-ion recognition. The interaction of Co^{II}, Ni^{II}, and Cu^{II} with new oxygen–nitrogen donor macrocycles; *X*-ray structures of complexes of Cu^{II} and Ni^{II} with a 15-membered O_2N_3 derivative, 2543–54
Molecular mechanics studies of a series of dicopper complexes of a 20-membered macrocycle containing differing types of bridges between the metal centres, 2563–72
Kinetic studies on 1:1 electron-transfer reactions involving blue copper proteins. Part 14. Reactions of poplar plastocyanin with inorganic complexes, 2573–8
Copper(I) and copper(II) compounds of a dinucleating, octadentate tetrakisbenzimidazole ligand. Crystal and molecular structures of three dihalogeno{1,8-bis[bis(1′-methylbenzimidazol-2′-ylmethyl)amino]-3,6-dioxaoctane}dicopper(II) bis(anion) compounds (halogen = F, Cl, or Br; anion = BF_4, CF_3SO_3, or Cl), 2579–94
Synthetic and equilibrium studies on copper(II), nickel(II), and palladium(II) complexes of *N*,*N*′,*N*″,*N*‴-tetrakis(2-hydroxyethyl)-1,4,8,11-tetra-azacyclotetradecane (L); kinetics of the acid dissociation of the copper(II) complex and *X*-ray structural characterisation of the nickel(II) complex $[NiLH_{-1}][ClO_4]$, 2605–14
Bicyclam [6,6′-bi(1,4,8,11-tetra-azacyclotetradecane)]: a ditopic receptor for homo- and hetero-bimetallic complexes, 2631–4
Nickel(II), copper(II), and zinc (II) complexes of penta-azamacrocyclic ligands. Crystal and molecular structure of 3,6,9,12-tetramethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-trienezinc(II) perchlorate, 2643–8
σ-Alkynyl complexes of manganese(I) as η²-bonding ligands for Group 1B metal–ligand fragments. *X*-Ray crystal structure of $[Mn_2Cu(\mu\text{-}CCBu^t)_2(CO)_6(dppe)_2]PF_6{\cdot}0.5CH_2Cl_2$, 2867–74
Crystal structure of a copper complex of 2-carboxypentonic acid; a decomposition product of dehydroascorbic acid, 2905–10
Oxygen- and carbon monoxide-binding to a lipophilic diporphyrinatocopperiron complex solubilized in an aqueous medium with a micelle, 2963–8
Binuclear copper(I) complexes of a 24-membered macrocycle (L^1) derived from 2,5-diformylthiophene and 3-thiapentane-1,5-diamine. Crystal and molecular structure of $[Cu_2L^1(NCS)][ClO_4]$; a unique example of a single end-to-end bridging thiocyanate between copper(I) atoms, 2975–80
Metal–nucleotide interactions: crystal structures of Ca[Cu(5′-GMP)$_2$(en)$(H_2O)_2$]·8H_2O(5′-GMP = guanosine 5′-monophosphate) and Ca[Cu(5′-IMP)$_2(H_2O)_4$]·6.7H_2O (5′-IMP = inosine 5′-monophosphate), 2999–3002
Syntheses and crystal structures of complexes $[M_2R_2]$, [M = Cu, Ag, or Au; R = 2-C($SiMe_3$)$_2C_5H_4N$] and $[Cu_4R'_4]$[R′ = 2-CH($SiMe_3$)C_5H_4N]; electrochemical generation of $[Cu_2R_2]^{2+}$, 3085–92

CORRINOID
The chemistry of vitamin B_{12}. Part 27. The cage phosphite ester 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane and the anionic dimethyl phosphite as ligands for cobalt(III) corrinoids, 761–8
The chemistry of vitamin B_{12}. Part 28. Crystal structure of dicyanocobyrinic acid heptamethyl ester and its interaction with alcohols: the effects of hydrogen bonding to co-ordinated cyanide, 1349–58

COVALENCE
Covalence in the $[Cr(CN)_6]^{3-}$ ion studied by *X*-ray diffraction from $Cs_2KCr(CN)_6$, 1747–54

CROWN ETHER

Stereochemical role of lone pairs in main-group elements. Part 3. Structure and bonding in trichloro(1,4,7,10,13-pentaoxacyclopentadecane)antimony(III) studied by means of *X*-ray crystallography at 120 K, 427–30

Correlation between thermal stability and molecular and crystal structure in a series of potassium–dibenzo-18-crown-6 solid compounds, 1779–84

Synthesis of *SS'*-bis(4'-benzo[15-crown-5])dithioglyoxime and its complexes with copper(II), nickel(II), cobalt(II), cobalt(III), palladium(II), platinum(II), and platinum(IV), 1827–32

CRYPTAND

A structural study of the complexation of the lithium ion by the cryptand 4,7,13-trioxa-1,10-diazabicyclo[8.5.5]icosane, 489–92

CRYSTAL STRUCTURE

Synthesis of $[Mo(\eta\text{-}C_5H_4Pr^i)(PR_3)_2H_3](R_3 = Me_3$ or $Me_2Ph)$, $[Mo(\eta\text{-}C_5H_4Pr^i)(Pr^i{}_2PCH_2CH_2PPr^i{}_2)H_3]$, and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of $[\{Mo(\eta\text{-}C_5H_4Pr^i)(\mu\text{-}Cl)_2\}_2]$, 9–20

Isomers in *fac*-$[ReBr(CO)_3(L_c)_2]$ (L_c = *cis*-3,4-dihydro-2,3,4,5-tetraphenyl-2*H*-1,2,3-diazaphosphole) arising from chirality of phosphorus ligands: separation and *X*-ray structural studies, 21–6

Phenyl complexes of ruthenium(II): competition between carbonyl substitution and formation of benzoyl complexes, and an *X*-ray study of a benzoyl complex, 27–34

Linear-chain platinum(II) dithiocarboxylates: crystal structure of tetrakis(phenyldithioacetato)diplatinum(II), 35–40

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 49. Synthesis of mixed-metal complexes with bonds between Cr, Mo or W and Co, Rh, Ir or Re; crystal structure of $[CrReRh_2(\mu_3\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}CO)(CO)_9(\mu\text{-}C_9H_7)_2]$, 41–50

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 50. Synthesis and crystal structures of the complexes $[RuW_3(\mu\text{-}CO)(\mu_3\text{-}\eta\text{-}CO)(\mu\text{-}CPh)_2(\mu_3\text{-}CPh)(CO)_2(\eta\text{-}C_5H_5)_3]$ and $[RuW_3(\mu\text{-}CO)(\mu_3\text{-}\eta\text{-}CO)\{\mu\text{-}C(Ph)C(O)\}(\mu\text{-}CPh)(\mu_3\text{-}CPh)(CO)(PMePh_2)(\eta\text{-}C_5H_5)_3]$, 51–60

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 51. Alkylidyne(carbaborane)tungsten-gold and -rhodium complexes; crystal structures of $[AuW(\mu\text{-}CR)(CO)_2(PPh_3)(\eta^5\text{-}C_2B_9H_9Me_2)]$, $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$, and $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2\{\eta^5\text{-}C_2B_9(C_7H_9)H_8Me_2\}]$ (R = C_6H_4Me-4), 61–72

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 52. Synthesis and reactions of the compounds $[RuW(\mu\text{-}CR)(CO)_3(\eta\text{-}C_5H_5)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[RuW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta\text{-}C_5H_5)(\eta^5\text{-}C_2B_9H_9Me_2)]$ and $[N(PPh_3)_2][RuW(\mu\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}\sigma,\eta^5\text{-}C_2B_9H_8Me_2)(CO)_3(\eta\text{-}C_5H_5)]$, 73–80

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 53. Alkylidyne(carbaborane)dimetal complexes with tungsten–molybdenum and –tungsten bonds; crystal structures of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ [M = Mo (two isomers) or W], 81–90

High-yield syntheses of sulphido triruthenium carbonyl cluster complexes containing bis(diphenylphosphino)methane, 91–6

Organotin biocides. Part 8. The crystal structure of triphenyltin formate and a comparative variable-temperature tin-119 Mössbauer spectroscopic study of organotin formates and acetates, 101–6

Binucleating N_6 24- and 26-membered macrocyclic ligands. Part 1. Dilead complexes: *X*-ray crystal structure determination of a macrocyclic dilead complex containing nitrogen-only bridging thiocyanate, 123–8

The first authenticated example of geometrical isomers in organotin(IV) co-ordination chemistry. The crystal and molecular structures of octahedral *cis*- and *trans*-SnR_2 isomeric adducts of dichlorobis(4-chlorophenyl)tin(IV) with 4,4'-dimethyl-2,2'-bipyridyl, 129–38

The crystal structures of the diphenyldithiophosphinates of antimony(III) and bismuth(III): $M(S_2PPh_2)_3$ (M = Sb or Bi), 145–50

The synthesis and structure of Group 5 metal alkyl and alkylidene complexes containing 2,6-dialkylphenoxide ligands: *X*-ray crystal structures of $[Ta(OC_6H_3Me_2\text{-}2,6)_2(CH_2Ph)_3]$, $[Ta(OC_6H_3Me_2\text{-}2,6)_4Me]$, and $[Ta(OC_6H_3Bu^t{}_2\text{-}2,6)_2(=CHSiMe_3)(CH_2SiMe_3)]$, 155–62

Oxoaryls of rhenium-(V) and -(VI) and osmium(VI). *X*-Ray crystal structures of dimesityldioxorhenium(VI), tetramesityloxorhenium(VI) and dimesityldioxo-osmium(VI), 169–76

Investigation into aroylhydrazones as chelating agents. Part 8. Synthesis and spectroscopic characterization of complexes of Co, Ni, Cu, Zn, and Cd with 2,6-diacetylpyridine bis(salicyloylhydrazone); *X*-ray crystal structure of dichloro[2,6-diacetylpyridine bis(salicyloylhydrazone)]cadmium(II)–chloroform–methanol (1/1/1), 177–82

Reactivity of ruthenium(II) complexes with 1-hydroxymethyl- or 1-benzyl-3,5-dimethylpyrazole and similar functionalized bipyrazoles. *X*-Ray crystal structure of carbonylchlorohydrido-(3,5-dimethylpyrazole-N^2)bis(triphenylphosphine)ruthenium(II), 183–6

Preparation and magnetism of the binuclear iron(II) complexes $[\{Fe(C_9H_{21}N_3)X_2\}_2]$ (X = NCS, NCO, or N_3) and their reaction with NO. Crystal structures of $[\{Fe(C_9H_{21}N_3)(NCS)_2\}_2]$ and $[Fe(C_9H_{21}N_3)(NO)(N_3)_2]$, 187–92

Secondary bonding. Part 14. Structural isomerism in diaryliodonium halides and the structure of di(*p*-tolyl)iodonium bromide, 193–6

Bis(dialkylphosphino)methane complexes of palladium(II) and the molecular structure of $[Pd_2Cl_3(\mu\text{-}Pr^i PCH_2PPr^i{}_2)(\mu\text{-}Pr^i{}_2PCH_2PHPr^i)]\cdot 2CHCl_3$ containing an unusual phosphidophosphine bridging ligand, 201–6

Complexes of ligands providing endogenous bridges. Part 1. The syntheses and crystal structures of barium and lead(II) complexes of macrocyclic Schiff bases derived from heterocyclic dicarbonyls and 1,*n*-diamino-*n'*-hydroxyalkanes (*n*,*n'* = 3,2; 4,2; or 5,3), 207–18

Thiosemicarbazones as co-ordinating agents. Part 2. Synthesis, spectroscopic characterization, and *X*-ray structure of aquachloro(pyridoxal thiosemicarbazone)manganese(II) chloride and aqua(pyridoxal thiosemicarbazonato)copper(II) chloride monohydrate, 227–34

Syntheses, spectroscopic, and electrochemical properties of dinuclear copper(II) complexes bridged by a single thiolate sulphur atom; *X*-ray crystal structures of $[Cu_2(L^1)_2(SC_6H_4Cl\text{-}4)][PF_6]$ and $[Cu_2(L^2)_2(SC_6H_4CH_3\text{-}4)][ClO_4]$, 241–8

Synthesis and *X*-ray crystal structure of a cuboctahedral platinum cluster, 259–62

Crystal and molecular structure of the six-co-ordinate antimony(III) complex $SbCl_3L$, where L = *N,N'*-dimethyloxamide. Unusual O,O ligand binding leading to four-membered $Sb(\mu\text{-}O)_2Sb$ ring formation, 263–6

Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of $[5,6,6\text{-}(PPh_3)_3\text{-}6\text{-}H\text{-}nido\text{-}6\text{-}RuB_9H_{12}]$ (1985, 2397), 275–6

Reactions of metal carbonyl cluster complexes with multidentate phosphine ligands; the preparation and thermolysis of bis(dimethylphosphino)methane derivatives of $[Ru_3(CO)_{12}]$, and the *X*-ray crystal structure of $[Ru_3(CO)_9(H)(Me_2PCHPMe_2)]$, 277–80

Magnetostructural correlations in parallel square-planar chloride bridged copper(II) dimers: structure, dynamic nuclear magnetic resonance study, and magnetic properties of $[Cu_2(terpy)_2Cl_2][PF_6]_2$, 285–92

Structure and properties of tris(quinoxaline-2,3-dithiolato)-molybdate(IV), 297–302

Preparations, crystal structures, and molybdenum-95 nuclear magnetic resonance spectroscopic studies of triphenylphosphine–gold–tetrathiomolybdate(VI) clusters, 303–6

Trimethylphosphine complexes of niobium(IV) and tantalum(IV). Crystal, molecular, and electronic structures of $[M_2Cl_4(\mu\text{-}Cl)_4(PMe_3)_4]$ (M = Nb or Ta), 307–14

Preparation of isocyanide and mixed dinitrogen–isocyanide complexes of rhenium(I) from reactions of *trans*-$[ReCl(N_2)(PMe_2Ph)_4]$, *mer*-$[Re(S_2PPh_2)(N_2)(PMe_2Ph)_3]$, or *mer*-$[Re(S_2CNEt_2)(N_2)(PMe_2Ph)_3]$ with methyl isocyanide; crystal structure of *mer*-$[Re(S_2PPh_2)(N_2)(CNMe)(PMe_2Ph)_3]$, 319–26

Reaction of $[Os_5H_2(CO)_{15}]$ with pyridine: crystal and molecular

CRYSTAL STRUCTURE (contd)
structures of $[Os_5H_3(CO)_{14}(C_5H_4N)]$ and $[Os_5H_2(CO)_{14}(C_5H_5N)]$, 327–34
Chemistry of platinum sulphido-complexes. Part 6. Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with the rhodium complexes $[(RhL_2Cl)_2]$ (L = CO or C_2H_4) and $[Rh(2,6\text{-}Me_2C_6H_3NC)_4]PF_6$. Crystal and molecular structures of $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(CO)_2]\cdot[Rh(CO)_2Cl_2]\cdot C_3H_6O$ and $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4\cdot(2,6\text{-}Me_2C_6\text{-}H_3NC)_2]PF_6$, 335–40
The addition of isonitriles to $[\{M(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (M = Mo or W). Syntheses and properties of $[M_2(\mu\text{-}\eta^2\text{-}CNR)(CO)_4(\eta\text{-}C_5H_5)_2]$ (M = Mo or W, R = Me or Bu^t) and $[\{Mo(CNR)(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (R = Me or Bu^t). *X*-Ray crystal structures of $[Mo_2(\mu\text{-}\eta^2\text{-}CNBu^t)(CO)_4(\eta\text{-}(\eta\text{-}C_5H_5)_2]$ and $[\{Mo(CNBu^t)(CO)_2(\eta\text{-}C_5H_5)\}_2]$, 341–8
Studies on transition-metal macrocyclic complexes. Single-crystal *X*-ray structure and electrochemistry of the bis-macrocyclic complex $[Cu(L)_2][ClO_4]_2\cdot 2H_2O$ (L = 1,4,7-triazacyclononane), 373–8
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au), 379–90
Activation of alkanes involving rhenium atoms: synthesis and electronic structure of binuclear μ-alkylidene derivatives; molecular structure of $[\{Re(\eta\text{-}C_6H_6)\}_2(\mu\text{-}CHBu^t)(\mu\text{-}H)_2]$, 391–402
Synthesis of η-arene rhenium compounds using rhenium atoms. Crystal structures of $[Re_2(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)_2(\mu\text{-}H)_2]$ and $[Re(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)H]$, 403–10
Stereochemical role of lone pairs in main-group elements. Part 3. Structure and bonding in trichloro(1,4,7,10,13-pentaoxacyclopentadecane)antimony(III) studied by means of *X*-ray crystallography at 120 K, 427–30
Co-ordination of highly substituted alkenes to transition-metal complexes: preparation and crystal and molecular structures of the complexes $[\{RhCl(Ph_2PO_2CCH{=}CMe_2)\}_2]$ and $[\{RhCl(Ph_2PO_2CCH{=}CHCH{=}CHMe)\}_2]$, 443–50
Ruthenium(II) complexes of *closo* dodecaboranyl anions and the molecular structure of the *pileo* thirteen-vertex ruthenaborane $[(PPh_3)_2ClRuB_{12}H_{11}(NEt_3)]$, 451–6
Electron spin resonance and electronic spectra and crystal and molecular structures of copper(II) amino acid complexes, 457–66
Formation of a methoxycarbonyl group by attack of methanol on an iron(II) carbonyl complex, assisted by intramolecular proton transfer: *X*-ray structure determination of $[Fe(CO)(PMe_3)_2(CNC_6H_{11})\{CO(OCH_3)\}\{C(CH_3){=}NHC_6H_{11}\}]\cdot BPh_4$, 473–6
Syntheses and reactions of acyclic and cyclic carbene complexes *cis*-$[MI(=CMeNHMe)(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo or W) and *cis*-$[MI\{=\overline{C(CH_2)_n}NMe\}(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo, n = 3 or 4; M = W, n = 3) obtained through alkyl migration to methyl isocyanide co-ordinated to molybdenum or tungsten. *X*-Ray structure of *cis*-$[MI\{=\overline{C(CH_2)_3}NMe\}(CO)_2(\eta\text{-}C_5H_5)]$ (1986, 2127), 479–80
A structural study of the complexation of the lithium ion by the cryptand 4,7,13-trioxa-1,10-diazabicyclo[8.5.5]icosane, 489–92
Synthetic, spectroscopic, and *X*-ray crystallographic studies on binuclear copper(II) complexes with a tridentate NNS-bonding 2-formylpyridine thiosemicarbazone ligand. The characterization of both neutral and deprotonated co-ordinated ligand structures, 493–500
Some binuclear molybdenum(VI) complexes obtained from condensation reactions: their crystal and molecular structures, 523–8
The chemistry of the hydrazido(1−)-ligand. Preparations and crystal structures of $[Mo(NHNHCO_2Me)(NNCO_2Me)(S_2CNMe_2)_2]$, $[Mo(NHNMePh)(NNMePh)(S_2CNMe_2)_2]BPh_4$, and $[ReCl_2(NHNHCOPh)(NNHCOPh)(PPh_3)_2]$. Mechanism of formation of substituted hydrazines from $[Mo(NNRPh)_2(S_2CNMe_2)_2]$ (R = Me or Ph), 529–40
Studies of pendant-arm macrocyclic ligands. Part 5. Synthesis of two pyridine-containing penta-aza macrocycles with single pyrrolidinyl pendant arms and characterisation of their nickel(II) and copper(II) complexes. Crystal structure of perchlorato{7-[2′-(1″-pyrrolidinyl)ethyl]-3,7,11,17-tetra-azabicyclo[11.3.1]-heptadeca-1(17),13,15-triene}nickel(II) perchlorate, 545–50
Triosmium clusters containing ligands derived from tropone (cycloheptatrienone): molecular structure of an oxidative addition product $[Os_3H(\mu_3\text{-}C_7H_5O)(CO)_9]$, 551–6
The synthesis and *X*-ray crystal structures of homoleptic tetrahedral aryls of osmium(IV) and of cyclohexyls of ruthenium(IV), osmium(IV), and chromium(IV), 557–62
Preparation of the nitrenes $ReF_5(NF)$, $ReF_5(NCl)$, and $ReF_5(NBr)$. *X*-Ray crystal structures of $ReF_5(NF)$ and $ReF_5(NCl)$, 567–72
Reactions of co-ordinated ligands. Part 39. The synthesis of carbyne complexes from alkyne molybdenum and tungsten cations; formation and crystal structures of $[Mo(\equiv CCH_2Bu^t)\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$ and $[\overline{Mo\{=C(SiMe_3)CH_2\}}\{P(OMe)_3\}_2(\eta^5\text{-}C_9H_7)]$, 591–604
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[\overline{Pt\{ON(Ph)C(NPh)S\}}(PPh_3)_2]$ and $[\overline{Pt\{ON(Ph)C(O)NPh\}}(PPh_3)_2]$, 605–8
Crystal structure of *trans*-dicarbonylbis(dimethylphenylphosphine)-(*N*-phenylsalicylideneaminato)rhenium(I) and *mer*-trichloro-(dimethylphenylphosphine)(*N*-phenylsalicylideneaminato)-rhenium(IV). Comparison of Re^I, Re^{III}, Re^{IV}, and Re^V complexes with the salicylideneaminato ligand and reaction mechanisms, 613–8
Synthesis and structure of μ-[*trans*-1,2-bis(diphenylphosphino)-ethylene]-bis[chlorogold(I)], 677–80
Methyl- and phenyl-mercury(II) derivatives of 2-mercaptopyridine. Crystal and molecular structure of methyl(pyridine-2-thiolato)-mercury(II) (1986, 1945), 683–4
Reactions of heteroboranes. Synthesis of $[2,2\text{-}(PPh_3)_2\text{-}1,2\text{-}SePtB_{10}H_{10}]\cdot CH_2Cl_2$, its crystal and molecular structure and that of $SeB_{11}H_{11}$, 699–704
A comparative synthetic and structural study of μ-oxo and μ-sulphido derivatives of *N*,*N*′-ethylenebis(acetylacetoneiminato)-iron(III), including the synthesis and structural characterization of the parent compound, 709–14
Amino-acid complexation to a cobalt(III) complex of optically-active (2*R*,5*R*,8*R*,11*R*)-2,5,8,11-tetraethyl-1,4,7,10-tetra-azacyclododecane (L). Preparation, characterization, and structure determinations of $[Co^{III}\{(R)\text{- or }(S)\text{-alaO}\}L]Br\cdot[ClO_4]\cdot H_2O$, 721–8
Metallacyclic complexes. Synthesis of metallathietane-3,3-dioxide (metallacyclosulphone) complexes of platinum(II), palladium(II), and nickel(II); crystal structure of *trans*-2,4-diphenyl-1,1-bis(triphenylarsine)platinathietane-3,3-dioxide, 733–40
The chemistry of trichloro[tris(trimethylsilyl)methyl] and trichloro[tris(dimethylphenylsilyl)methyl] complexes of gallium, indium, and thallium. Crystal and molecular structures of $[Li(thf)_2(\mu\text{-}Cl)_2GaCl\{C(SiMe_2Ph)_3\}]\cdot thf$, $[Li(thf)_3(\mu\text{-}Cl)InCl_2\{C(SiMe_3)_3\}]$, and $[\{(Me_3Si)_3C\}In(\mu\text{-}Cl)_2\{\mu\text{-}Fe(CO)_4\}In\{C(SiMe_3)_3\}]$ (thf = tetrahydrofuran), 747–56
Bimetallic systems. Part 17. Synthesis of Group 6 metal(0)–copper(I), –silver(I), or –gold(I) complexes with two $Ph_2PCH_2PPh_2$ (dppm) bridging ligands: crystal structure of $[(OC)_5W(\mu\text{-}CO)(\mu\text{-dppm})_2(\mu\text{-}Cl)Cu]\cdot CH_2Cl_2$, 769–76
Synthesis and characterisation of $[N(PPh_3)_2][RuRh_5(CO)_{16}]$ by *X*-ray crystallography and by natural-abundance carbon-13 nuclear magnetic resonance spectroscopy, 781–4
Crystal structure and mass spectrometry of dichlorodimethylbis[2(1*H*)-pyridinethione-*S*]tin(IV), 815–8
Mono- and di-carbonyl complexes of molybdenum(II) with mixed bidentate ligands. *X*-Ray crystal structure of $[MoBr_2(CO)(bipy)(Ph_2PCH_2PPh_2)]$ (bipy = 2,2′-bipyridine): a novel type of $[M(bidentate)_2(unidentate)_3]$ structure, 819–24
Redox behaviour of the iron–sulphur cluster $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{2+}$. Synthesis and crystal structure of the new paramagnetic monopositive species $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^+$ as its $[PF_6]^-$ salt, 831–6
Reactions of aluminopolyhydride complexes of tungsten. *X*-Ray crystal structures of $[(Me_3P)_3H_3W(\mu\text{-}H)_2AlCl(\mu\text{-}N{=}CHEt)]_2$, $\{[(Me_3P)_3WH_5]Li\}_4$, $WH_2I(PMe_3)_3(SiMe_3)$, and $(Me_3P)_3H_2W(\mu\text{-}H)_3AlCl_2(NMe_3)$, 837–46
New heterocyclic compounds containing niobium and molybdenum; crystal structure of a cyclomolybdaphosphazene, 847–50
Reactions of cyclobutenedione and cyclopentenetrione with zero-

CRYSTAL STRUCTURE (contd)
valent nickel triad complexes of isocyanide. *X*-Ray crystal structures of (3—4-η-3,4-diphenylcyclobut-3-ene-1,2-dione)- and (4—5-η-4,5-diphenylcyclopent-4-ene-1,2,3-trione)-tris(t-butyl isocyanide)nickel, 853–8
Synthesis and characterization of technetium-(IV) and -(III) complexes with bidentate Schiff bases. *X*-Ray molecular structure of chloro(dimethylphenylphosphine)bis(*N*-phenylsalicylideneiminato)technetium(III), 867–72
Dicopper(II) complexes of a 28-membered N_8 macrocycle. Evidence for proton transfer from co-ordinated secondary amine to thiocyanate, leading to formation of an isothiocyanic acid complex, 873–80
Synthesis of *o*-nitrophenylplatinum(IV) complexes; crystal and molecular structure of dichlorobis(2-nitrophenyl-*C'O*)-platinum(IV), 881–4
Synthesis of pyridine-containing tetra-aza macrocycles: 3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), its 3,11-dibenzyl (L^2) and 3,7,11-tribenzyl (L^3) derivatives, and their nickel(II), copper(II), and zinc(II) complexes: crystal structures of L^2·HCl and $[Ni(L^2)Cl]ClO_4·H_2O$, 889–94
Rational synthesis of metalla–sulphur–nitrogen complexes: use of $[Me_2SnS_2N_2]_2$ in the preparation of complexes of the type $[M(S_2N_2H)(PR_3)_2]X$ (M = Pt or Pd). *X*-Ray structures of $[Pt(S_2N_2H)(PEt_3)_2][Me_2SnCl_3]$, $[Pt(S_2N_2H)(PBu^n_3)_2]_2[PF_6]Cl$, and $[\{Co(S_2N_2H)_2\}_2\{Ph_2P(O)CH_2CH_2P(O)Ph_2\}]$, 907–14
Some investigations of $S_5N_5^+$ compounds; *X*-ray crystal structure of $[S_5N_5]Cl$, 915–24
Trinuclear osmium clusters, containing bidentate $Ph_2PCH_2PPh_2$ and $Me_2PCH_2PMe_2$ ligands: *X*-ray crystal structures of the complexes $[Os_3H(CO)_9(Me_2PCHPMe_2)]$ and $[Os_3H(OH)(CO)_8(Ph_2PCH_2PPh_2)]$, 931–8
Synthesis and characterization of monocyclopentadienyl niobium derivatives. *X*-Ray crystal structure of di-μ-chloro-μ-oxo-bis[dichloro(η-trimethylsilylcyclopentadienyl)niobium(V)], $[\{Nb(\eta\text{-}C_5H_4SiMe_3)Cl_2\}_2(\mu\text{-}O)(\mu\text{-}Cl)_2]$, 953–6
Phosphinoalkylsilyl complexes. Part 7. Synthesis and reactivity of octahedral, bis(chelate)dicarbonyl complexes of ruthenium(II) and osmium(II). Conformational properties of the bis(chelate) framework and *X*-ray crystal structures of $[M(PPh_2CH_2CH_2SiMe_2)_2(CO)_2]$ (M = Ru or Os), 969–74
New dicyclopentadienyl phosphine, phosphite, and acetylene niobium(III) complexes. Crystal structure of $[Nb(\eta\text{-}C_5H_4SiMe_3)_2Cl(PhC{\equiv}CPh)]$, 975–80
2-Pyridonate (Opy) rhodium(I) carbonyl complexes. Crystal structure of the unusual tetranuclear complex $[Rh_4(\mu\text{-}Opy)_4(\mu\text{-}CO)_2(CO)_4]$, 981–4
Studies on transition-metal peroxo complexes. Part 7. Molybdenum(VI) and tungsten(VI) carboxylato peroxo complexes, and the *X*-ray crystal structure of $K_2[MoO(O_2)_2(glyc)]·2H_2O$, 991–6
Thermally induced phase transition and *X*-ray crystal structure of bis(*N*,*N'*-dimethyl-1,2-ethanediamine)di-isothiocyanatonickel(II), 997–1000
The heteronuclear cluster chemistry of the Group 1B metals. Part 3. Synthesis, structures, and chemical properties of the trimetallic hexanuclear Group 1B metal cluster compounds $[MM'Ru_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]$ (M = Cu, M' = Ag or Au; M = Ag, M' = Au). *X*-Ray crystal structure of the complex $[AgCuRu_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]·CH_2Cl_2$, 1001–8
Ternary metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of copper(II)–2,2'-bipyridyl– pyridoxic acid and copper(II)–2,2'-bipyridyl–pyridoxic acid γ-lactone complexes, 1009–16
Extended Hückel molecular orbital study of the effects of edge-bridging hydrogen atoms on the lengths of boron–boron and metal–metal bonds in cluster compounds, and the crystal structure of benzyltrimethylammonium octahydrotriborate(1−), 1017–26
Reactions of copper(II) with $[S_2P(OR)_2]^-$ (R = Et or Pr^i) and single-crystal *X*-ray studies of $Cu[S_2P(OEt)_2]$·bipy and $Cu[S_2P(OEt)_2]·2PPh_3$, 1027–34
Electron-transfer platinum complexes of esters of dithiocarbazic acid. The crystal and molecular structure of $[Pt\{N(CH_2Ph)NC(S)SMe\}_2]$, 1035–40
Reactions of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}X_2]$ (M = Mo, X = I; M = W, X = Cl) with some amine-substituted pyridines. The molecular structures of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}(2\text{-}NHC_5NH_4)_2]$, 1063–70
The ring-stacking principle in organolithium chemistry. Its development through the isolation and crystal structures of hexameric iminolithium clusters $(RR'C{=}NLi)_6$ (R' = Ph, R = Bu^t or Me_2N; R = R' = Me_2N or Bu^t), 1071–82
A structural comparison of bis(quinoxaline-2,3-dithiolato)cuprate complexes, 1083–8
Lewis-base adducts of Group 11 metal(I) compounds. Part 27. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance, far-infrared, and structural studies on the mononuclear 2:1 adducts of triphenylphosphine with copper(I) and gold(I) halides, 1089–98
Lewis-base adducts of Group 11 metal(I) compounds. Part 28. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance and structural studies on the mononuclear 3:1 adducts of triphenylphosphine with copper(I) halides, 1099–106
Studies of the pyridine-containing tetra-azamacrocycle 3,7,11-trimethyl-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1) and its complexes with Ni^{2+}, Cu^{2+}, and Zn^{2+}. Crystal structures of the unsymmetric isomer of $[Ni(L^1)][ClO_4]_2$, and the five-co-ordinate complexes $[Ni(L^1)(dmso)][ClO_4]_2$ (dmso = dimethyl sulphoxide) and $[Ni(L^1)Cl]ClO_4$, 1107–14
The preparation, structure, and reactions of the mononuclear thiolate–hydride complexes $[MoH(SR)(dppe)_2]$ (R = bulkyl alkyl or aryl group, dppe = $Ph_2PCH_2CH_2PPh_2$); *X*-ray structure determination of *cis*-$[MoH(SC_6H_2Pr^i_3\text{-}2,4,6)(dppe)_2]$, 1115–22
Chemistry of metallacyclic complexes containing the FeCSCS ring obtained by metal-promoted CS_2–CS coupling. *X*-Ray crystal structures of $[(CO)(cp)FeC\{SW(CO)_5\}SC(FeL_n)S]$ and $[(CO)(cp)FeC(SEt)SC(FeL_n)S]SO_3CF_3$ [FeL_n = Fe(cp)$(CO)_2$, cp = η-C_5H_5], 1133–44
Synthesis and stereochemical studies of the chiral ruthenium complexes $[Ru(\eta\text{-}C_5H_5)\{(S)dpompyr\text{-}PP'\}X]$ [dpompyr = *N*-diphenylphosphino-2-(diphenylphosphinoxymethyl)pyrrolidine, X = H or Cl]. Crystal structure of $[(S)Ru(\eta\text{-}C_5H_5)\{(S)dpompyr\text{-}PP'\}Cl]$, 1149–56
Crystal structures of low- and high-spin iron(III) complexes with quadridentate Schiff bases, 1157–62
Dimeric tungsten(V) compounds containing $[(S)W(\mu\text{-}S)_2W(S)]^{2+}$. Syntheses and structures of di-μ-sulphido-bis[(diethyl dithiophosphato-*SS'*)sulphidotungsten(V)] and di-μ-sulphido-bis[(*N*,*N*-diethyldithiocarbamato-*SS'*)sulphidotungsten(V)], 1163–8
The preparation of $[Ru(NCMe)_6][7\text{-}(\eta^6\text{-}C_6Me_6)\text{-}nido\text{-}7\text{-}RuB_{10}H_{13}]_2$ and its characterization by single-crystal *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy: a triply hydrogen-bridged eleven-vertex *nido*-7-metallaundecaborane, 1169–74
Reactions of planar pentadentate ligands with palladium and platinum chlorides. Crystal structures of *cis*-dichlorobis(*N'*-isopropylidenebenzohydrazide)platinum(II) and the cyclopalladated dianion of *N'*,*N'''*-(pyridine-2,6-diyldiethylidyne)bis(benzohydrazide), 1175–80
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 54. Compounds derived from $[W{\equiv}CC_6H_4Me\text{-}4)(CO)_2(\eta\text{-}C_5Me_5)]$ with bonds between tungsten and iron, cobalt, or nickel; crystal structure of $[FeW\{\mu\text{-}C(C_6H_4Me\text{-}4)C(O)O\}(CO)_5(\eta\text{-}C_5Me_5)]$, 1191–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 55. Coupling of methylene and alkylidyne groups at a trimetal centre; crystal structures of $[Ru_2W\{\mu\text{-}\sigma{:}\eta\text{-}trans\text{-}CH{=}C(H)R\}(\mu\text{-}CO)(CO)_3(\eta\text{-}C_5H_5)_2(\eta\text{-}C_5Me_5)]·CH_2Cl_2$ and $[Ru_2W(\mu_3\text{-}CCH_2R)(\mu\text{-}CO)_3(CO)(\eta\text{-}C_5H_5)_3]$ (R = C_6H_4Me-4), 1201–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 56. Synthesis of iron–molybdenum compounds; crystal structures of $[FeMo(\mu\text{-}CR)(CO)_6(\eta\text{-}C_5H_5)]$ and $[FeMo_2(\mu_3\text{-}RC_2R)(CO)_6(\eta\text{-}C_5H_5)_2]$ (R = C_6H_4Me-4), 1209–14
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 57. Reactions of iron– molybdenum complexes with oxygen and sulphur; crystal structure of $[FeMo(\mu\text{-}\eta^2\text{-}SCC_6H_4Me\text{-}4)(CO)_5(\eta\text{-}C_5H_5)]$, 1215–20
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 58. Some reactions of the compounds

CRYSTAL STRUCTURE (contd)
$[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (M = Mo or W); crystal structures of the complexes $[W_2(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_2(L)(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$-[L = $N_2C(C_6H_4Me\text{-}4)_2$ or O], 1221–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 59. Reactions of the alkylidyne compounds $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Cr, Mo, or W, R = C_6H_4Me-4; M = W, R = Me, $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate] with d^{10} complexes; crystal structure of $[PtW_2(\mu\text{-}CMe)_2(CO)_4\{HB(pz)_3\}_2]$, 1229–34
The crystal structure of bismuth diethyldithiophosphinate–benzene (1/1), $Bi(S_2PEt_2)_3\cdot C_6H_6$, 1257–60
Crystal and molecular structure of a novel mixed-spin binuclear nickel(II) complex with *N,N′,N″,N‴*-tetrakis(2-aminoethyl)-1,4,8,11-tetra-azacyclotetradecane, 1261–2
Binuclear copper(I) complexes containing bis(diphenylphosphino)-methane bridging ligands: crystal structure of $[Cu_2(\mu\text{-}dppm)_2(MeCN)_4][ClO_4]_2$, 1275–8
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au) (1987, 379), 1283–4
Triaquahenicosatungstodiphosphate(6−) heteropolyanion, $[P_2W_{21}O_{71}(OH_2)_3]^{6-}$: *X*-ray crystallographic and ^{183}W nuclear magnetic resonance structural studies (1986, 2237), 1283–4
Mixed oxidation state trinuclear cobalt complexes with bridging sulphito and Schiff-base ligands. Part 1. Preparation of the complexes $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}L)_2Co^{III}{}_2(ROH)_2]$ (L = Schiff base anion, R = alkyl) and structure determination of $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}\alpha,\alpha'\text{-}Me_2\text{-}salpd)_2Co^{III}{}_2(Pr^nOH)_2]\cdot 2Pr^nOH$, 1305–12
Reactions of co-ordinated ligands. Part 40. Synthesis, structure, and reactivity of dirhenium μ-allylidene complexes; crystal structures of $[Re_2(\mu\text{-}\sigma;\eta^3\text{-}CHCHCMe_2)(CO)_8]$ and $[Re_2(\mu\text{-}\sigma:\eta^3\text{-}CHCHCMe_2)(CO)_9]$, 1313–8
Reactions of co-ordinated ligands. Part 41. Synthesis and reactions of cationic functionalised alkyne molybdenum complexes; formation of carbyne and vinylidene complexes and crystal structure of $[Mo(C{\equiv}CHPh)Br\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$, 1319–28
Gold(III) complexes with nitriles containing aromatic nuclei: crystal and molecular structure of trichloro(*p*-toluonitrile)gold(III), 1329–32
Spectroscopic, magnetic, and structural properties of diaquadimethanolbis(*N*-tosylvalinato)copper(II) and its 2,2′-bipyridine adduct, 1333–40
The chemistry of vitamin B_{12}. Part 28. Crystal structure of dicyanocobyrinic acid heptamethyl ester and its interaction with alcohols: the effects of hydrogen bonding to co-ordinated cyanide, 1349–58
Syntheses and characterization of dinuclear palladium and platinum complexes with a single unsupported bridging group. Crystal structure of $[\{C_6H_3(CH_2NMe_2)_2\text{-}o,o'\}Pd(\mu\text{-}Cl)Pd\{C_6H_3(CH_2NMe_2)_2\text{-}o,o'\}]BF_4$, 1359–66
Reaction of elemental sulphur with cobalt(II)–Schiff-base complexes: synthesis of μ-disulphido and μ-tetrasulphido binuclear cobalt(III) complexes. Crystal structures of two binuclear cobalt(III)–Schiff-base complexes, 1367–76
Synthesis and characterization of palladium(II)–η^3-allyl–ylide complexes. *X*-Ray crystal structure of $[PdCl(\eta^3\text{-}2\text{-}MeC_3H_4)\{Ph_3PC(H)COMe\}]$, 1381–8
Pentamethylcyclopentadienylrhodaborane chemistry. Part 2. The reaction of $[6\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}6\text{-}RhB_9H_{13}]$ with dimethylphenylphosphine, and the characterization of $[5\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}5\text{-}RhB_9H_{11}\text{-}7\text{-}(PMe_2Ph)]$, $[2\text{-}(\eta^5\text{-}C_5Me_5)\text{-}closo\text{-}2\text{-}RhB_9H_7\text{-}3,10\text{-}(PMe_2Ph)_2]$, and $[2\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}2\text{-}RhB_8H_{10}\text{-}5\text{-}(PMe_2Ph)]$ by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 1431–44
Metallocene derivatives of early transition elements. Part 4. Synthesis and crystal structures of a series of zirconocene(IV) halides $[Zr(\eta\text{-}C_5H_4SiMe_3)_2X_2]$ (X = Cl or Br) and $[Zr\{\eta\text{-}C_5H_3(SiMe_3)_2\text{-}1,3\}_2X_2]$ (X = F, Br, or I), 1463–72
Cyclopalladation to form planar tridentate [N–C–N]$^-$ intramolecular co-ordination systems involving pyridine donor groups, including ligand synthesis and *X*-ray structural studies., 1477–84
Bimetallic systems. Part 19. Chromium(0)–, molybdenum(0)–, or tungsten(0)–iron(0) carbonyl complexes containing one or two bridging $Ph_2PCH_2PPh_2$ ligands: crystal structure of $[(OC)_3Mo(\mu\text{-}Ph_2PCH_2PPh_2)_2Fe(CO)_3]$, 1489–94
Bimetallic systems. Part 21. Synthesis of vinylidene- and phenylene-bridged complexes of dinickel(II), nickel(II)–palladium(II), and nickel(II)–platinum(II) containing $Ph_2PCH_2PPh_2$ (dppm): crystal structures of $[(SCN)Ni(\mu\text{-}dppm)_2(\mu\text{-}C{=}CH_2)Ni(NCS)]$ and $[ClPd(\mu\text{-}dppm)_2(\mu\text{-}C{=}CCl_2)NiCl]$, 1501–8
Bimetallic systems. Part 22. Molybdenum(0)– and manganese(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the reversible uptake of carbon monoxide: crystal structure of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)Mo(CO)_4]\cdot 0.5C_6H_5Cl$, 1509–14
Intermediates in the conversion of $[Os(CO)_{11}(PRPh_2)]$ (R = Me or Ph) into $[Os_3(\mu_3\text{-}C_6H_4)(\mu_3\text{-}PR)(CO)_9]$: crystal and molecular structures of $[Os_3(\mu\text{-}H)(\mu_3\text{-}C_6H_4PMePh\text{-}o)(CO)_9]$ and $[Os_3(\mu_3\text{-}C_6H_4PMe\text{-}o)(CO)_{10}]$, 1529–34
The triphenylphosphine cone angle and restricted rotation about the chromium–phosphorus bond in dicarbonyl(η^6-hexa-alkylbenzene)(triphenylphosphine)chromium(0) complexes. Crystal and molecular structure of dicarbonyl(η^6-hexa-n-propylbenzene)(triphenylphosphine)chromium(0), 1545–50
Preparation and reactivity of halide-bridged binuclear carbonyl manganese and rhenium anions, $[M_2(\mu\text{-}X)(CO)_{8-2n}\{(EtO)_2POP(OEt)_2\}_n]^-$ (X = halide; M = Mn, *n* = 1 or 2; M = Re, *n* = 1). *X*-Ray crystal structure of $[Mn_2Au(\mu\text{-}Br)(CO)_6(\mu\text{-}tedip)(PPh_3)]$, 1551–6
Synthesis and substitution reactions of a heterodimetallic molybdenum–manganese complex, $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_6]$; *X*-ray crystal structure of $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_4(dppm\text{-}PP')]$, 1557–64
Crystal and molecular structure of $[\{Ag[S_2P(OEt)_2]\cdot PPh_3\}_2]$, 1569–72
Preparation and crystal structures of $[PPh_4]_2[M_2(SPh)_6]$ (M = Zn or Cd), 1577–80
Comments on the attempts by Wickens and Abrams to synthesize manganese(II) phosphine complexes: a comment, 1581–2
Synthetic, spectroscopic, and *X*-ray crystallographic studies on binuclear copper(II) complexes with a tridentate NNS-bonding 2-formylpyridine thiosemicarbazone ligand. The characterization of both neutral and deprotonated co-ordinated ligand structures (1987, 493), 1583–4
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\{ON(Ph)C(NPh)S\}(PPh_3)_2]$ and $[Pt\{ON(Ph)C(O)NPh\}(PPh_3)_2]$ (1987, 605), 1583–4
Structure and magnetism of analogous *O*-bridged (phenolato) and *S*-bridged (thiophenolato) binuclear copper(II) complexes. Crystal structure of the binuclear *S*-bridged complex μ-[2,6-bis(4′-cyclohexyl-4′-hydroxy-2′,3′-diazabuta-1′,3′-dien-1′-yl)-4-methylthiophenolato(3−)-*S,N*^{2},*O*(Cu^{1,2})]-methanol-μ-(pyrazolyl-*N,N′*)-dicopper(II), 1585–92
Complexes of 2,5,7,9-tetrathiabicyclo[4.3.0]non-1(6)-ene-8-thione (L^1). Part 1. Synthesis and antimony-121 Mössbauer spectroscopic characterisation of $SbCl_3(L^1)_{1.5}$, crystal and molecular structures of $SbCl_3(L^1)_{1.5}$ and uncomplexed L^1, 1605–10
Synthesis and characterization of new heterobimetallic iron–cobalt complexes with ethenyl bridges $CR^2{=}CHR^1$ [R^1 = H or C(O)OMe, R^2 = C(O)OMe or C(O)OEt]. *X*-Ray structure of $[FeCo(CO)_6\{\mu\text{-}MeOC(O)C{=}C(H)C(O)OMe\}]$, 1619–22
Substituted osmium clusters containing labile ligands. The preparation, characterisation, and some reactions of $[Os_3H(OR)(CO)_9(MeCN)]$ R = H, Me, Et, or Ph) and the isomers of $[Os_3H(SR)(CO)_9(MeCN)]$ (R = Me, Et, or Ph); the crystal and molecular structure of $[Os_3H(SPh)(CO)_9(PEt_3)]$, 1623–30
Ring-size and substituent effects in relatively rigid macrocyclic quinquedentate ligands. Structure determinations of $[ZnL^4(H_2O)_2][ClO_4]_2\cdot MeCN$ and $[ZnL^3(ClO_4)_2]$, 1631–6
Crystal and molecular structures of $[MCl(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf) and $[TiCl_{0.5}Br_{0.5}(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$: an unusual π-interaction with the metals and molecular orbital calculations on model compounds, 1641–6
Chemistry of 2,3-dihydro-2-metallaphenalenes. Part 2. Formation of $[CHMMe_2C(MMe_3)C_{10}H_6]^x$ *via* base decomposition of $[1,8\text{-}(Me_3MCH_2)_2C_{10}H_6]^y$ (M = Ge, *x* = 2−, *y* = 0; M = P, *x* = 0,

CRYSTAL STRUCTURE (contd)
$y = 2+$); crystal structures of the novel bis-ylide $[CHPMe_2C(PMe_3)C_{10}H_6]$ and of a distorted *peri*-ylide, 1-Me-8-$(Me_3PCH)C_{10}H_6$, 1647–54
Neutral and anionic 2-methyl-6-nitrophenyl palladium(II) complexes. The *X*-ray crystal structure of bis(benzyltriphenylphosphonium) *trans*-di-μ-chloro-bis[chloro(2-methyl-6-nitrophenyl)palladate(II)]–acetone (1/2), 1655–8
Unsymmetrical bridging of an acetylide ligand in the cation $[Pt_2(C{\equiv}CBu^t)_3(\mu\text{-dppm})_2]^+$ (dppm = $Ph_2PCH_2PPh_2$): *X*-ray structure determination of the tetraphenylborate salt, 1659–64
Syntheses, *X*-ray structures, and properties of complexes of macrocyclic hexaimines with lanthanide nitrates, 1665–74
Preparation and partial oxidation of tin(II) trifluoroacetate: spectroscopic properties and *X*-ray crystal structure of di-μ_3-oxo-octakis-μ-trifluoroacetato-tetratin(II)tin(IV), 1675–82
Technetium(V) and rhenium(V) complexes with *N*-(2-mercaptophenyl)salicylideneiminate. Crystal structure of chloro(oxo)[*N*-(2-sulphidophenyl)salicylideneiminato-*NOS*]technetium(V), 1693–700
Crystal and molecular structure of $[Co_8(CO)_{21}(\mu_5\text{-}CS_2)(\mu_5\text{-}C_2S)]$: a thiocarbonyl group as a six-electron donor, 1707–12
Charge density in the $CoCl_4^{2-}$ ion: a comparison with spin density and theoretical calculations, 1737–46
Covalence in the $[Cr(CN)_6]^{3-}$ ion studied by *X*-ray diffraction from $Cs_2KCr(CN)_6$, 1747–54
Synthesis and reactions of $[(OC)_2Re(\mu\text{-dppm})_2(\mu\text{-Cl})_2Rh(\eta^2\text{-}C_2H_4)]$ (dppm = $Ph_2PCH_2PPh_2$): crystal structure of the 1:2 chlorobenzene solvate, 1763–8
Correlation between thermal stability and molecular and crystal structure in a series of potassium–dibenzo-18-crown-6 solid compounds, 1779–84
Co-oligomerization of diphenylketene and ethylene promoted by (η^5-cyclopentadienyl)bis(ethylene)cobalt(I): synthesis and *X*-ray structure of a heterocycle containing cobalt(III), 1789–92
X-Ray crystal structure and properties of dibenzotetrathiafulvalenium trichlorodiethylstannate(IV), 1793–6
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$], 1805–12
Crystal and molecular structure of chromium(II) sulphate pentahydrate and single-crystal electronic spectra and bonding of $CrSO_4{\cdot}5H_2O$, $CuSO_4{\cdot}5H_2O$, and $CuSO_4{\cdot}5D_2O$, 1817–22
Mixed chloro-, *NN*-dialkyldithiocarbamato-, and *S*-alkyl *NN*-dialkyldithiocarbamate complexes of platinum(II). Crystal structure of $[PtCl(S_2CNEt_2)(MeS_2CNEt_2)]$, 1823–6
Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNN′N′*-tetra-acetate (cdta): structural and magnetic characterization of $[(H_2O)_4Cu(cdta)Ni]{\cdot}3H_2O$ and $[(H_2O)_5Ni(cdta)Cu]{\cdot}H_2O$, 1847–52
Decarboxylation of vinylene carbonate at triosmium clusters; *X*-ray crystal structure of the μ_3-formylmethylidene compound $[Os_3(CHCHO)(CO)_{10}]$, 1861–4
Complexes of ligands providing endogenous bridges. Part 3. Non-transition metal complexes of macrocycles derived from 2,6-diacetyl-4-methylphenol; the crystal structure of a mononuclear barium complex, 1865–8
Three isomeric hydride compounds formed by protonation of decacarbonylbis(dimethylphenylphosphine)triosmium and the *X*-ray structures of two of these: 1,1- and 1,2-$[Os_3H(CO)_{10}(PMe_2Ph)_2][PF_6]$, 1869–76
Tetranuclear amido complexes of copper(I): a synthetic and structural study, 1883–8
Co-ordination chemistry of higher oxidation states. Part 23. Synthesis and properties of tetrahalogenoiridium(IV) complexes, $[IrL_2X_4]$ [X = Cl or Br; L = pyridine, PR_3, AsR_3, SbR_3, SR_2, or SeR_2]. Crystal and molecular structure of *trans*-$[Ir(AsEt_3)_2Br_4]$, 1901–10
Dithio-oxamides as ligands: crystal structures and vibrational analyses of bis(*N*,*N′*-dicyclohexyldithio-oxamidato)palladium(II), bis(*N*,*N′*-dibenzyldithio-oxamide)copper(II) diperchlorate, and dichloro(*N*,*N′*-dimethyldithio-oxamide)zinc(II) and the carbon-13 nuclear magnetic resonance spectra of related compounds, 1921–8
Electrophilic attack on a series of dinuclear diphosphazane-bridged derivatives of iron by halogens. *X*-Ray crystal structures of $[Fe_2I(CO)_5\{\mu\text{-}(MeO)_2PN(Et)P(OMe)_2\}_2]PF_6$ and $[Fe_2(\mu\text{-}Br)(CO)_4\{\mu\text{-}(PhO)_2PN(Et)P(OPh)_2\}_2]PF_6$, 1933–46
Activation energies for molecular tumbling and cyclopentadienyl rotation in $[M(\eta^5\text{-}C_5H_5)(\eta^4\text{-cod})]$ (M = Rh or Ir; cod = cycloocta-1,5-diene) and the *X*-ray crystal structure of and bonding in $[Rh(\eta^5\text{-}C_5H_5)(\eta^4\text{-cod})]$, 1947–52
X-Ray structural study of [Fe(saltrien)]X (X = Br·$2H_2O$, BPh_4, or PF_6). Origin of unusual magnetic behaviour of the spin-crossover complex $[Fe(saltrien)]PF_6$, 1957–62
The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of $[M_2Ru_3(\mu_3\text{-}S)(\mu\text{-}Ph_2PCH_2PPh_2)(CO)_9]$ (M = Cu or Au), 1967–76
Transformation of $[PPh_4][Fe_5RhC(CO)_{16}]$ under hydroformylation conditions: synthesis and crystal structure of $[PPh_4][Fe_3Rh_3C(CO)_{15}]$, 1983–8
The preparation and crystal structure of $5SbF_3{\cdot}3SbF_5$, 1989–98
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 8. Irreversible oxidation of trimethylphosphinemanganese(II) dihalides, $[MnX_2(PMe_3)]$ (X = Cl, Br, or I), to bis(trimethylphosphine)manganese(III) trihalides, $[MnX_3(PMe_3)_2]$. *X*-Ray crystal structure of *trans*-triiodobis(trimethylphosphine)manganese(III), the first example of a trigonal-bipyramidal manganese(III) tertiary phosphine complex, 1999–2004
Metal ion-bridged intramolecular stacking interaction between the tryptophyl residue and the aromatic heterocyclic amine within the ternary complex (1,10-phenanthroline)(L-tryptophanato-*O*,*N*)copper(II) perchlorate–water (1/2.5): an *X*-ray study, 2017–22
(η-Benzene)bis(trimethylphosphine)iron: a useful precursor to polyene iron derivatives. Crystal structure of 2—4-η:2′—4′-η-1,1′-bi(cyclohex-3-en-2-yl)bis(trimethylphosphine)iron(II), 2031–8
Insertion of 2,6-dimethylphenyl isocyanide into the U–N bond of $[U(\eta\text{-}C_5H_5)_3(NEt_2)]$. Crystal structure of $[U(\eta\text{-}C_5H_5)_3\{C(NEt_2){=}NC_6H_3Me_2\text{-}2,6\}]$, 2039–44
Bis(organoimido)-complexes of tungsten(VI): the crystal and molecular structures of tetrachlorobis(t-butylamine)bis(t-butylimido)bis(μ-*p*-tolylimido)-ditungsten(VI) and tetrachlorobis(μ-phenylimido)-bis(t-butylamine)bis(t-butylimido)ditungsten(VI), 2059–66
Preparation and characterisation of mononuclear organoimido complexes of tungsten: *X*-ray crystal structures of $[WCl_4(NC_6H_4Me\text{-}p)(thf)]$, $[P(CH_2Ph)Ph_3][WCl_5(NC_6H_4Me\text{-}p)]$, and $[WCl_2(NPh)_2(bipy)]$, 2067–76
Reaction of elemental sulphur with bis(pentamethylcyclopentadienyl)vanadium derivatives: disulphido and μ-sulphido complexes. Crystal structures of $[V(\eta\text{-}C_5Me_5)_2(S_2)]$ and $[V_2(\eta\text{-}C_5Me_5)_2Cl_2(\mu\text{-}S)_2]$, 2099–104
Reactions of main group homopolynuclear cations with transition-metal carbonyls. Stabilisation of $\mu\text{-}Se_2^{2+}$ and $\eta^3\text{-}Te_3^{2+}$ dicationic ligands and crystal structures of $[Mo(Te_3)(CO)_4][SbF_6]_2$ and $[FeW(Se_2)(CO)_8][SbF_6]_2$, 2111–6
Platinum(II) complexes with ligands $(RO)_2PCH_2P(OR)_2$ (R = Me, Et, Ph, or C_6H_4Me-4); crystal structure of *cis*,*cis*-$[Pt_2Me_4\{\mu\text{-}(EtO)_2PCH_2P(OEt)_2\}_2]$, 2117–24
Studies in Werner clathrates. Part 4. Structures of tetrakis(4-ethylpyridine)di-isothiocyanatonickel(II) and its clathrates with *p*-, *m*-, and *o*-xylene, carbon disulphide, and carbon tetrachloride, 2125–40
The syntheses and crystal structures of two iminolithium adducts: $(Ph_2C{=}NLi{\cdot}NC_5H_5)_4$, a tetrameric pseudo-cubane formed by stacking two $(LiN)_2$ four-membered rings, and $[Bu^t_2C{=}NLi{\cdot}OP(NMe_2)_3]_2$, a dimer prevented sterically from stacking, 2141–8
Methylindium dialkylamido compounds: mono- and bis-dialkylamides involving four- and five-co-ordinated indium(III). *X*-Ray crystal structures of $\{Me_2In[N(CH_2)_2N(Me)CH_2CH_2]\}_2$, $MeIn[MeNC(CH)_4N]_2$, and $MeIn\{[MeN(CH_2)_2NMe]\text{-}InMe_2\}_2$, 2159–64
Synthesis and reactions of the co-ordinatively unsaturated hexafluorobut-2-yne complex $[WBr_2(CO)(CF_3C{\equiv}CCF_3)_2]$. The crystal and molecular structure of $[\{W[C(CF_3){=}C(CF_3)P(OEt)_2O](\mu\text{-}Br)(CO)(CF_3C{\equiv}CCF_3)\}_2]$, 2165–70
Some chemistry of the heterobimetallic complexes $[(OC)_3Mo(\mu\text{-}dppm)_2Rh(CO)Cl]$ and $[(OC)_3Mo(\mu\text{-}dppm)_2\text{-}M(CO)_2]PF_6$

CRYSTAL STRUCTURE (contd)
(M = Rh or Ir, dppm = $Ph_2PCH_2PPh_2$). Structure of $[(OC)_3Mo(\mu\text{-dppm})_2(\mu\text{-C}{\equiv}\text{CMe})Rh(CO)]$, 2171–6
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 62. Synthesis of penta-, hexa-, and hepta-heteronuclear metal cluster compounds involving tungsten or molybdenum with platinum or nickel: crystal structures of the chain compounds $[Pt_3W_3(\mu\text{-CMe})(\mu_3\text{-CMe})_2(CO)_6(cod)(\eta\text{-}C_5Me_5)_3]\cdot 0.5CH_2Cl_2$ and $[Pt_3W_4(\mu\text{-}CC_6H_4Me\text{-}4)_2(\mu_3\text{-}CC_6H_4Me\text{-}4)_2(CO)_8(\eta\text{-}C_5H_5)_4]$, 2177–88
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 63. Synthesis of eight-membered-ring metallacycles: *X*-ray crystal structures of $[Pt_4W_4(\mu\text{-CR})(\mu_3\text{-CR})_3(\mu\text{-CO})(CO)_7(\eta\text{-}C_5H_5)_4]\cdot CH_2Cl_2$, $[Ni_2Pt_2W_4(\mu\text{-CR})(\mu_3\text{-CR})_3(\mu\text{-CO})(CO)_7(\eta\text{-}C_5H_5)_4]\cdot CH_2Cl_2$ (R = C_6H_4Me-4), and $[Ni_2Pt_2W_4(\mu_3\text{-CPh})_4(CO)_8(\eta\text{-}C_5H_5)_4]$, 2189–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 64. Addition of methylene groups to iron–molybdenum complexes; crystal structures of $[FeMo(\mu\text{-}CH_2)\{\mu\text{-}\sigma{:}\eta\text{-}C(C_6H_4Me\text{-}4){=}CH_2\}(CO)_5(\eta\text{-}C_5H_5)]$ and $[FeMo\{\mu\text{-}C(C_6H_4Me\text{-}4)C(OMe)C(H)\}(CO)_5(\eta\text{-}C_5H_5)]$, 2201–10
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 65. Reactions of the complexes $[FeW(\mu\text{-CR})(CO)_5(\eta\text{-}C_5Me_5)]$ and $[Fe_2W(\mu_3\text{-CR})(\mu\text{-CO})(CO)_8(\eta\text{-}C_5Me_5)]$ (R = C_6H_4Me-4) with alkynes $R'C_2R'$ (R′ = Me or Ph); crystal structures of $[FeW\{\mu\text{-}C(R)C(O)C(Me)C(Me)\}(CO)_5(\eta\text{-}C_5Me_5)]\cdot CH_2Cl_2$ and $[FeW\{\mu\text{-}C(R)C(Et)C(H)C(Me)C(Me)\}(\mu\text{-CO})(CO)_3(\eta\text{-}C_5Me_5)]$, 2211–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 66. Carbaboranetungsten–platinum complexes. Polyhedral rearrangements of a 12-vertex cage system; crystal structures of $[PtW(CO)_2(PEt_3)_2\{\eta^6\text{-}C_2B_9H_8\text{-}(CH_2C_6H_4Me\text{-}4)Me_2\}]\cdot CH_2Cl_2$, $[PtW(\mu\text{-H})\{\mu\text{-}\sigma{:}\eta^5\text{-}C_2B_9H_7\text{-}(CH_2C_6H_4Me\text{-}4)Me_2\}(CO)_2(PMe_3)(PEt_3)_2]$, and related compounds, 2219–34
Crystal structures and properties of $[Au(phen)\{(CN)_{0.92}Br_{0.08}\}_2]Br$ and $[Au(phen)(CN)\{(CN)_{0.82}Br_{0.18}\}]\cdot 0.5$*trans*-$[Au(CN)_2Br_2]\cdot 0.5Br\cdot phen$ (phen = 1,10-phenanthroline) obtained by disproportionation of five-co-ordinate bromodicyano(1,10-phenanthroline)gold(III). Two examples of secondary co-ordination and CN/Br disorder in square-planar gold(III) complexes, 2235–40
A reversible metal framework rearrangement in high-nuclearity osmium-platinum cluster compounds. *X*-Ray crystal structures of $[Os_6Pt(CO)_{17}(\mu_3\text{-NCMe})(C_8H_{12})]$, $[Os_6Pt(CO)_{17}(\mu_4\text{-NCMe})(C_8H_{12})]$ (C_8H_{12} = cyclo-octa-1,5-diene), and $[Os_6(CO)_{19}(\mu_3\text{-NCMe})]$: compounds with novel metal–acetonitrile bonding modes, 2245–52
Synthesis, reactivity, and *X*-ray crystal structures of $[Os_6Pt_2H_2(CO)_{16}(\mu\text{-}C_8H_{10})(C_6H_{12})]\cdot CH_2Cl_2$ and $[Os_6H(CO)_{16}(\mu\text{-}C_8H_{11})]$. Model compounds for C–H activation, 2253–60
Crystal structure of tris(quinoxaline-2,3-dithiolato)molybdate(v); a distorted trigonal prismatic complex, 2267–72
Synthesis of chelate stabilised alcohol- and alkoxo–platinum(II) complexes. *X*-Ray crystal structure of

$[Pt(PPh_2CH_2CMe_2O)_2]\cdot 3.5H_2O$, 2273–80
The synthesis, crystal and molecular structure of a copper(I) complex of 2(1*H*)-pyridinethione: an intermediate in Cu_4S_4 heterocubane build up, 2281–4
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_3(PPh_3)_2\{S{=}C(NHPh)\text{-}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O\text{–}C(NHPh){=}S\}]$, 2299–304
Preparation and crystal structures of bis(tetraphenylarsonium) tris(oxalato)technetate(IV), and tetraphenylarsonium tris(benzene-1,2-dithiolato)technetate(V): octahedral *versus* trigonal-prismatic geometry for tris-bidentate complexes of technetium, 2305–10
The reaction of di-2-pyridyl ketone with antimony(III) fluoride. Preparation, properties, Mössbauer spectrum and crystal structure of difluoro(methoxydi-2-pyridylmethoxo-*NN′O*)antimony(III), 2337–40
Axially asymmetric metal alkyls. Part 5. Synthesis and reduction to Zr^{III} species of Group 4 metallepines $[ML(\eta\text{-}C_5H_5)_2][L = (2\text{-}CHRC_6H_4)_2^{2-}$, R = H, M = Ti, Zr, or Hf), isomers of $[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), and *meso*-$[M\{1,8\text{-}(CHSiMe_3)_2C_{10}H_6\}(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf); *X*-ray crystal structures of $[TiL(\eta\text{-}C_5H_5)_2]$ (R = H) and λ-and δ-*RS*-$[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), 2347–58
Metal(II) hexafluoroantimonates: preparation and characterization of $MF_2\cdot 2SbF_5$ (M = Mg, Ni, Zn, Fe, Co, Cu, Cr, Ag, Cd, or Pb) and the *X*-ray structure determination of $AgF_2\cdot 2SbF_5$, 2379–84
Synthesis, molecular structure, and nuclear magnetic resonance spectra of the anionic monocarbametallaborane $[NEt_4][(CO)_3(PPh_3)(OH)$-*closo*-$MoCB_{10}H_{10}]\cdot CH_2Cl_2$, 2385–90
Polymeric thiolate complexes of Group 12 metals. Crystal and molecular structures of *catena*-[μ-(3-dimethylammonio-1-propanethiolate)]-dichlorocadmium(II) and bis[3-(dimethylammonio)propyl] disulphide tetrabromocadmate(II), 2391–6
Aqua(2,2′-bipyridyl)difluorocopper(II) dihydrate: *X*-ray structure reveals short hydrogen bonds and other unusual features, 2397–400
Synthesis and crystal structure of tris(morpholinecarbodithioato-*S*,*S*′)manganese(IV) pentaiodide–dichloromethane (2/1), 2407–10
Pentamethylcyclopentadienylrhodaborane chemistry. Part 3. The reaction of $[\{Rh(\eta^5\text{-}C_5Me_5)Cl_2\}_2]$ with *closo*-$[B_{10}H_{10}]^{2-}$ and *arachno*-$B_{10}H_{12}$-6,9-$(PMe_2Ph)_2$, and the characterisation of some novel rhodaboranes by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 2417–30
Preparation, crystal and molecular structure, and antimony-121 Mössbauer spectrum of trifluoro(1,10-phenanthroline)(thiourea)-antimony(III), 2451–4
The cyclometallation of benzoic acid to give rhodium, iridium, and osmium *C*,*O*-benzoates. *X*-Ray structure determination of the dibenzoate $[(C_5Me_5)Rh(OOCPh)_2(H_2O)]$, 2459–66
Substitution reactions of the osmium(I) dinuclear compound $[Os_2(MeCO_2)_2(CO)_6]$. *X*-Ray crystal structure of $[N(PPh_3)_2][Os_2(MeCO_2)_2Cl(CO)_5]$, 2473–8
Preparation, nuclear magnetic resonance spectra, and *X*-ray diffraction study of bis(*N*-benzylacetimidic acid ethyl ester)dichloropalladium(II), 2503–4
A realistic model for heme-containing catalases and peroxidases: the *X*-ray structural characterisation of a non-porphyrin iron(III) macrocyclic complex, and the mechanism of its peroxidation of aromatic substrates, 2505–10
Complexes of ligands providing endogenous bridges. Part 4. Copper(II) complexes of macrocyclic Schiff bases derived from 2,6-diacetylpyridine and 1-*n*-diamino-*n*′-hydroxyalkanes (*n*,*n*′ = 3,2; 4,2; and 5,3): synthesis, properties, and structures, 2519–30
Studies involving mixed donor macrocyclic ligands. Comparative *X*-ray diffraction, infrared, and fast atom bombardment mass spectrometric studies of the nickel thiocyanate complexes of three O_2N_3-donor macrocycles, 2537–42
Studies of metal-ion recognition. The interaction of Co^{II}, Ni^{II}, and Cu^{II} with new oxygen–nitrogen donor macrocycles; *X*-ray structures of complexes of Cu^{II} and Ni^{II} with a 15-membered O_2N_3 derivative, 2543–54
Formation of a (4 + 4) Schiff-base macrocyclic ligand by a template rearrangement. Crystal and molecular structures of two tetranuclear manganese(II) complexes, 2555–62
Copper(I) and copper(II) compounds of a dinucleating, octadentate tetrakisbenzimidazole ligand. Crystal and molecular structures of three dihalogeno{1,8-bis[bis(1′-methylbenzimidazol-2′-ylmethyl)amino]-3,6-dioxaoctane}dicopper(II) bis(anion) compounds (halogen = F, Cl, or Br; anion = BF_4, CF_3SO_3, or Cl), 2579–94
Synthetic and equilibrium studies on copper(II), nickel(II), and palladium(II) complexes of *N*,*N*′,*N*″,*N*′-tetrakis(2-hydroxyethyl)-1,4,8,11-tetra-azacyclotetradecane (L); kinetics of the acid dissociation of the copper(II) complex and *X*-ray structural characterisation of the nickel(II) complex $[NiLH_{-1}][ClO_4]$, 2605–14
Subvalent group 14 metal compounds. Part 10. Syntheses and structures of some bis(metallo)plumbylenes $[Pb\{Mo(R)(CO)_3\}_2(OC_4H_8)]$ [R = η-C_5H_5, η-$C_5H_3(SiMe_3)_2$-1,3, or η-C_5Me_5]; *X*-ray crystal structures of $[Pb\{Mo(\eta\text{-}C_5Me_5)(CO)_3\}_2(OC_4H_8)]$ and $[\{Pb[Mo(\eta\text{-}C_5Me_5)(CO)_3][Mo(\eta\text{-}C_5Me_5)(CO)_2(\mu\text{-CO})]\}_2]$, 2635–42
Nickel(II), copper(II), and zinc (II) complexes of penta-azamacrocyclic ligands. Crystal and molecular structure of 3,6,9,12-tetramethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-trienezinc(II) perchlorate, 2643–8

CRYSTAL STRUCTURE (contd)

Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato molybdenum and tungsten nitrosyls. Part 6. Synthesis and electrochemistry of complexes containing saturated heterocyclic amide ligands. *X*-Ray crystal and molecular structure of $[\{Mo(NO)[HB(Me_2pz)_3]Cl\}_2(NC_4H_8N)]$, 2653–60

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 67. Reactions of the salts $[X][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ [X = $P(CH_2Ph)Ph_3$, R = C_6H_4Me-4; X = NEt_4, R = Ph] with bis(cyclo-octa-1,5-diene)platinum, 2661–8

The reaction of $[Mn_2\{\sigma:\eta^2\text{-}C(R)=CHR\}(\mu\text{-}PPh_2)(CO)_7]$ (R = H or Ph) with nucleophiles: *X*-ray crystal structures of $[N(PPh_3)_2][Mn_2(\mu\text{-}H)\{\mu\text{-}C(O)C(Ph)=CHPh\}(\mu\text{-}PPh_2)(CO)_6]$, $[Mn_2\{PPh_2C(Ph)=CHPh\}(CO)_9]$, $[Mn_2(\mu\text{-}PPh_2\text{-}CH=CH_2)(CO)_8]$, $[Mn_2\{\mu\text{-}C(O)C(Ph)=CHPh\}(\mu\text{-}PPh_2)(CO)_6(CNBu^t)_2]$, $[Mn_2\{\mu\text{-}\sigma:\eta^2\text{-}C(Ph)=CHPh\}(\mu\text{-}PPh_2)(CO)_6(CNBu^t)]$, and $[Mn(\mu\text{-}PPh_2CH=CH_2)(CO)_7(PEt_3)]$, 2669–82

Synthesis and study of monomeric and dimeric amidine complexes of nickel(II) and platinum(II). Crystal structure of bis(*N,N'*-diphenylbenzamidino)platinum(II), $[Pt\{PhNC(Ph)NPh\}_2]$, 2687–94

Organic chemistry of dinuclear metal centres. Part 9. Diruthenium complexes of buta-1,3-diene. *X*-Ray crystal structure of $[Ru_2(CO)_2(\mu\text{-}s\text{-}trans\text{-}\eta^{2'}:\eta^{2'}\text{-}CH_2CHCHCH_2)(\eta\text{-}C_5H_5)_2]$, 2703–8

Reactions of tetra-μ-carboxylato-diruthenium(II,II) compounds. *X*-Ray crystal structures of $Ru_2(\mu\text{-}O_2CCF_3)_4(thf)_2$, $Ru_2(\mu\text{-}O_2CR)_4(NO)_2$ (R = Et or CF_3), and $\{Na_3[Ru_2(\mu\text{-}O_2CO)_4]\cdot 6H_2O\}_n$, 2723–36

Heteronuclear complexes containing Group 6 transition metals; chemical, spectroscopic, and theoretical studies on some binuclear complexes and the *X*-ray crystal structure determination of $[WNi(CO)_3(PPh_3)_2(\eta^5\text{-}C_5H_5)]$, 2741–50

Rhodium(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the formation of a tetranuclear cluster from a heterobimetallic precursor: crystal structures of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)RhCl(CO)]$ and $[Fe_2Rh_2(\mu\text{-}Ph_2PCH_2PPh_2)_2(CO)_8]$, 2751–6

Gold–boron chemistry. Part 1. Synthetic, structural, and spectroscopic studies on the compounds $[5,6\text{-}\mu\text{-}(AuPR_3)\text{-}nido\text{-}B_{10}H_{13}]$ (R = cyclo-C_6H_{11} or C_6H_4Me-2), 2761–8

Diplatinum-(II) and -(III) complexes with bridging μ-diphosphonito-*P,P'* or μ-methylenebis(phosphonito)-*P,P'* ligands. Synthesis, structure, and nuclear magnetic resonance analysis of the 'lantern' complexes. Crystal structures of $K_4[Pt_2\{\mu\text{-}CH_2[P(O)OH]_2\}_4]\cdot 6H_2O$ and $K_4[Pt_2\{\mu\text{-}CH_2[P(O)OH]_2\}_4Cl_2]\cdot 8H_2O$, 2769–80

Organoruthenaborane chemistry. Part 5. Products of the reaction between *closo*-$[B_{10}H_{10}]^{2-}$ and $[(\eta^6\text{-}C_6Me_6)ClRuB_3H_8]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[\{(\eta^6\text{-}C_6Me_6)_2Ru_2H_4\}RuB_{10}H_9(OEt)_2]$, 2781–8

Dinitrogen complexes derived from tricarbonyl(η^6-hexaethylbenzene)chromium(0): crystal and molecular structure of μ-dinitrogen-bis[dicarbonyl(η^6-hexaethylbenzene)-chromium(0)]–toluene (1/1), 2789–92

The syntheses and crystal and molecular structures of some ten-vertex (MCB_8; M = Ru or Os) metallacarbaboranes: 9,9,9-(CO)$(PPh_3)_2$-*nido*-9,6-MCB_8H_{10}-5-PPh_3)] (M = Ru or Os) and [9,9,9-(η-C_5H_5)(H)(PPh_3)-*arachno*-9,6-$RuCB_8H_{12}$], 2793–802

Reactions of co-ordinated ligands. Part 42. The synthesis and reactions of cationic η^4-tetraphenylcyclobutadiene-ruthenium complexes; *X*-ray crystal structures of $[Ru(NCMe)(\eta^4\text{-}C_4Ph_4)(\eta\text{-}C_5H_5)][BF_4]$ and $[Ru\{\eta^5\text{-}CH(Ph)C\text{-}(Ph)C(Ph)C(Ph)O\}(\eta\text{-}C_5H_5)]$, 2803–14

The chemistry of tetraphenylcyclopentadienone complexes of ruthenium and rhodium: the *X*-ray crystal structure of $[Ru\{\eta^5\text{-}C_5Ph_4OC(O)CH(OMe)Ph\}(CO)_2Cl]$, 2815–22

Reactions of phosphine, arsine, and stibine with carbonylbis(triethylphosphine)iridium(I) halides. Part 1. Reactions in toluene; *X*-ray crystal structures of $[Ir(CO)ClH(PEt_3)_2(AsH_2)]$ and $[Ir(CO)XH(PEt_3)_2(\mu\text{-}ZH_2)RuCl_2(\eta^6\text{-}MeC_6H_4CHMe_2\text{-}p)]$ (X = Br, Z = P; X = Cl, Z = As), 2831–8

Carbene complexes. Part 21. Synthesis and characterisation of bis(carbene)molybdenum(II) complexes and dimetal(0) complexes of the Group 6 elements containing novel bridging bis(carbene) ligands; *X*-ray structures of $[Mo(CO)_2(L^{Et})_2(OSO_2CF_3)_2]$ [L^{Et} = ≡CN(Et)$(CH_2)_2$NEt] and $[W(CO)_5\{C(OEt)\text{-}CH_2C_6H_4CH_2C(OEt)\text{-}o\}W(CO)_5]$, 2843–52

Mixed-metal phosphinito complexes of platinum(II) and palladium(II) with lanthanide and actinide elements. The single-crystal *X*-ray structure determination of $[UO_2(OH_2)\{(OPPh_2)_2Pd(S_2CNEt_2)\}_2]$, 2853–6

σ-Alkynyl complexes of manganese(I) as η^2-bonding ligands for Group 1B metal–ligand fragments. *X*-Ray crystal structure of $[Mn_2Cu(\mu\text{-}CCBu^t)_2(CO)_6(dppe)_2]PF_6\cdot 0.5CH_2Cl_2$, 2867–74

Manganese(I) complexes with (tricyclohexylphosphonio)dithiocarboxylate as chelate and unidentate ligand. *X*-Ray crystal structure of *fac*-$[Mn(CO)_3\{S_2CP(C_6H_{11})_3\}_2]ClO_4\cdot H_2O$, 2875–80

Crystal structure of a copper complex of 2-carboxypentonic acid; a decomposition product of dehydroascorbic acid, 2905–10

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 68. Reactions between nonacarbonyldi-iron and the salts $[NEt_4][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = Me, Ph, or C_6H_4Me-4); crystal structures of $[NEt_4][FeW\{\mu\text{-}CH(C_6H_4Me\text{-}4)\}(\mu\text{-}\sigma:\eta^5\text{-}C_2B_9H_8Me_2)(\mu\text{-}CO)(CO)_5]$ and $[NEt_4][FeW(\mu_3\text{-}CPh)(\mu\text{-}\sigma:\sigma',\eta^5\text{-}C_2B_9H_7Me_2)(CO)_8]$, 2917–26

Dienophile cycloaddition to cycloheptatriene and related complexes of tricarbonyliron. *X*-Ray crystal structures of $[Fe(CO)_3(\eta^4\text{-}C_7H_7CN)]\cdot C_2(CN)_4$ and $[Fe(CO)_3(\eta^4\text{-}C_7H_7\text{–}C_7H_7)]$, 2927–34

Reactions of allene with triosmium and triruthenium clusters including coupling and oxidative addition: *X*-ray structure of the coupled allene compound $[Os_3(C_6H_8)(CO)_{10}]$, 2935–40

The chemistry of chromium nitrile complexes of 1,2-bis(dimethylphosphino)ethane. *X*-Ray crystal structures of *trans*-$[Cr^{IV}Cl(NEt)(dmpe)_2]CF_3SO_3$, *trans*-$[Cr^{IV}(N{=}CHMe)_2(dmpe)_2][BPh_4]_2$, and *trans*-$[Co^0(H\cdot NCMe)(dmpe)^2][BPh_4]_2$, 2947–54

Binuclear copper(I) complexes of a 24-membered macrocycle (L^1) derived from 2,5-diformylthiophene and 3-thiapentane-1,5-diamine. Crystal and molecular structure of $[Cu_2L^1(NCS)][ClO_4]$; a unique example of a single end-to-end bridging thiocyanate between copper(I) atoms, 2975–80

Reactions of diazo-compounds with the electron-rich germylene $Ge[N(SiMe_3)_2]_2$. Characterisation and some addition reactions of $[(Me_3Si)_2N]_2GeNNC(COOMe)_2$: crystal and molecular structure of $[(Me_3Si)_2N]_2Ge[NHN{=}C(COOMe)_2][C(N_2)COPh]$, 2981–8

Metal–nucleotide interactions: crystal structures of $Ca[Cu(5'\text{-}GMP)_2(en)(H_2O)_2]\cdot 8H_2O$ (5′-GMP = guanosine 5′-monophosphate) and $Ca[Cu(5'\text{-}IMP)_2(H_2O)_4]\cdot 6.7H_2O$ (5′-IMP = inosine 5′-monophosphate), 2999–3002

Dinuclear rhodium and iridium complexes of dicarboxyimidazolates; crystal structure of $[NBu_4][(cod)Rh(dci)Rh(cod)]\cdot 2Pr^iOH$, 3003–8

Mono- and bi-nuclear gold(I) and gold(III) complexes with $S_2C\text{–}PR_3$ ligands. *X*-Ray crystal structures of $[Au(C_6F_5)_3(S_2C\text{–}PEt_3)]$ and $[Au_2(\mu\text{-}S_2CPEt_3)(C_6F_5)_6]\cdot CH_2Cl_2$, 3017–22

Molecular structures of some (dimethylamino)halogenosilanes in the gas phase by electron diffraction and the crystal and molecular structures of mono-and di-chloro(dimethylamino)silane by *X*-ray diffraction at low temperatures, 3035–42

Structure and electrochemical properties of bis(pyridine)cobalt(III) complexes of *N,N'*-ethylenebis(salicylideneimine) homologues with an aryl substituent on the ethylene backbone. Crystal structures of $[Co(salphen)(py)_2]PF_6\cdot 0.5Et_2O$ and $[Co(salcppn)(py)_2]PF_6\cdot 2py$ (py = pyridine), 3055–60

Platinum(II)– and palladium(II)–iron(0) complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand. Crystal structures of $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}CO)PtBr_2]$ and $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}I)PtI]\cdot CH_2Cl_2$, 3079–84

Syntheses and crystal structures of complexes $[M_2R_2]$, [M = Cu, Ag, or Au; R = 2-$C(SiMe_3)_2C_5H_4N$] and $[Cu_4R'_4]$[R' = 2-$CH(SiMe_3)C_5H_4N$]; electrochemical generation of $[Cu_2R_2]^{2+}$, 3085–92

Preparation and structure of nickel(II) complexes of N-substituted tetra-aza macrocycles: molecular structure of [(1*RS*,4*SR*,5*RS*,8*SR*,11*RS*,12*SR*)-1,5,8,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane]nickel(II) diperchlorate, 3093–8

Preparation of two mixed-metal carbonyl clusters based on the platinumtriosmium core. *X*-Ray crystal structure of $[Os_3PtH_2(cod)(CH_2)(CO)_9]$ (cod = cyclo-octa-1,5-diene), 3103–6

CRYSTAL STRUCTURE (contd)
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S{=}C(NHPh){-}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O{-}C(NHPh){=}S\}]$, 3107–8
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$] (1987, 1805), 3107–8

CYANIDE
Tin-119 nuclear magnetic resonance studies of some pseudohalogeno-derivatives of $[SnX_6]^{2-}$ (X = Cl or Br), 315–8
Spectrochemistry of solutions. Part 19. Complexation of gold(I) by cyanide in liquid ammonia: an infrared and Raman spectroscopic study, 673–6
The relative nucleophilic efficiency of water and the azide and thiocyanate ions in the aqueous substitution of chloropentacyanocobaltate(III), 1059–62
Electronic *d–d* spectra and ligand-dissociation rates of pentacyanoruthenate(II) ions, 1267–70
Gold(III) complexes with nitriles containing aromatic nuclei: crystal and molecular structure of trichloro(*p*-toluonitrile)gold(III), 1329–32
The chemistry of vitamin B_{12}. Part 28. Crystal structure of dicyanocobyrinic acid heptamethyl ester and its interaction with alcohols: the effects of hydrogen bonding to co-ordinated cyanide, 1349–58
Nitrogen-14 nuclear quadrupole resonance study of bonding in transition-metal cyanides, 1637–40
Covalence in the $[Cr(CN)_6]^{3-}$ ion studied by *X*-ray diffraction from $Cs_2KCr(CN)_6$, 1747–54
Alkylation of tricarbonyliron complexes by trimethylsilyl cyanide: synthetic and kinetic studies, 2013–6
Crystal structures and properties of $[Au(phen)\{(CN)_{0.92}Br_{0.08}\}_2]Br$ and $[Au(phen)(CN)\{(CN)_{0.82}Br_{0.18}\}]\cdot0.5$*trans*-$[Au(CN)_2Br_2]\cdot0.5Br$-phen (phen = 1,10-phenanthroline) obtained by disproportionation of five-co-ordinate bromodicyano(1,10-phenanthroline)gold(III). Two examples of secondary co-ordination and CN/Br disorder in square-planar gold(III) complexes, 2235–40
The relative nucleophilic efficiency of water and dimethyl sulphoxide in the aqueous substitution of aquapentacyanocobaltate(III) and chloropentacyanocobaltate(III), 2311–6

CYANOGEN
Triosmium clusters containing the ligands CNH_2, NCHCN, and NCMeCN derived from cyanide ion, cyanogen, and acetonitrile, 1457–62

CYCLOBUTADIENE
Isometallocenes. Part 1. Polymethylated isocobaltocenium salts; a general route to $[Co(C_4Me_4)(arene)]$, 2757–60
Reactions of co-ordinated ligands. Part 42. The synthesis and reactions of cationic η^4-tetraphenylcyclobutadiene-ruthenium complexes; *X*-ray crystal structures of $[Ru(NCMe)(\eta^4\text{-}C_4Ph_4)(\eta\text{-}C_5H_5)][BF_4]$ and $[Ru\{\eta^5\text{-}CH(Ph)C(Ph)C(Ph)C(Ph)O\}(\eta\text{-}C_5H_5)]$, 2803–14

CYCLOBUTENEDIONE
Reactions of cyclobutenedione and cyclopentenetrione with zero-valent nickel triad complexes of isocyanide. *X*-Ray crystal structures of (3—4-η-3,4-diphenylcyclobut-3-ene-1,2-dione)- and (4—5-η-4,5-diphenylcyclopent-4-ene-1,2,3-trione)-tris(t-butyl isocyanide)nickel, 853–8

CYCLODEXTRIN
Mössbauer spectroscopic studies on molecular rotation and lattice dynamics of cyclodextrin clathrates of bis(η-cyclopentadienyl)-iron(II) and some of its ring-substituted derivatives, 627–32
On the induced circular dichroism of *d–d* transitions of a 14-membered macrocyclic tetra-amine nickel(II) complex in β-cyclodextrin, 1801–4

CYCLOHEPTATRIENE
Dienophile cycloaddition to cycloheptatriene and related complexes of tricarbonyliron. *X*-Ray crystal structures of $[Fe(CO)_3(\eta^4\text{-}C_7H_7CN)]\cdot C_2(CN)_4$ and $[Fe(CO)_3(\eta^4\text{-}C_7H_7\text{–}C_7H_7)]$, 2927–34

CYCLOHEPTATRIENONE
Triosmium clusters containing ligands derived from tropone (cycloheptatrienone): molecular structure of an oxidative addition product $[Os_3H(\mu_3\text{-}C_7H_5O)(CO)_9]$, 551–6

CYCLOHEPTATRIENYL
The chemistry of cycloheptatrienyl complexes of molybdenum and tungsten: the synthesis and reactions of some vinylidene and alkynyl derivatives, 3049–54

CYCLOHEXADIENYL
Transition-metal mediated asymmetric synthesis. Part 5. Alternative methods for the formation of tricarbonyl(η^5-cyclohexadienyl)-iron(1+) salts, 885–8
Alkylation of tricarbonyliron complexes by trimethylsilyl cyanide: synthetic and kinetic studies, 2013–6

CYCLOHEXANEDIAMINETETRA-ACETATE
Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNN′N′*-tetra-acetate (cdta): structural and magnetic characterization of $[(H_2O)_4Cu(cdta)Ni]\cdot3H_2O$ and $[(H_2O)_5Ni(cdta)Cu]\cdot H_2O$, 1847–52

CYCLOHEXYL
The synthesis and *X*-ray crystal structures of homoleptic tetrahedral aryls of osmium(IV) and of cyclohexyls of ruthenium(IV), osmium(IV), and chromium(IV), 557–62

CYCLOHEXYLIMINIOETHYL
Formation of a methoxycarbonyl group by attack of methanol on an iron(II) carbonyl complex, assisted by intramolecular proton transfer: *X*-ray structure determination of $[Fe(CO)(PMe_3)_2(CNC_6H_{11})\{CO(OCH_3)\}\{C(CH_3){=}NHC_6H_{11}\}]BPh_4$, 473–6

CYCLOMETALLAPHOSPHAZENE
New heterocyclic compounds containing niobium and molybdenum; crystal structure of a cyclomolybdaphosphazene, 847–50

CYCLOMETALLATION
The cyclometallation of benzoic acid to give rhodium, iridium, and osmium *C,O*-benzoates. *X*-Ray structure determination of the dibenzoate $[(C_5Me_5)Rh(OOCPh)_2(H_2O)]$, 2459–66

CYCLO-OCTADIENE
Activation energies for molecular tumbling and cyclopentadienyl rotation in $[M(\eta^5\text{-}C_5H_5)(\eta^4\text{-cod})]$ (M = Rh or Ir; cod = cyclo-octa-1,5-diene) and the *X*-ray crystal structure of and bonding in $[Rh(\eta^5\text{-}C_5H_5)(\eta^4\text{-cod})]$, 1947–52
A reversible metal framework rearrangement in high-nuclearity osmium-platinum cluster compounds. *X*-Ray crystal structures of $[Os_6Pt(CO)_{17}(\mu_3\text{-}NCMe)(C_8H_{12})]$, $[Os_6Pt(CO)_{17}(\mu_4\text{-}NCMe)(C_8H_{12})]$ (C_8H_{12} = cyclo-octa-1,5-diene), and $[Os_6(CO)_{19}(\mu_3\text{-}NCMe)]$: compounds with novel metal–acetonitrile bonding modes, 2245–52
Synthesis, reactivity, and *X*-ray crystal structures of $[Os_6Pt_2H_2(CO)_{16}(\mu\text{-}C_8H_{10})(C_6H_{12})]\cdot CH_2Cl_2$ and $[Os_6H(CO)_{16}(\mu\text{-}C_8H_{11})]$. Model compounds for C–H activation, 2253–60
Dinuclear rhodium and iridium complexes of dicarboxyimidazolates; crystal structure of $[NBu_4][(cod)Rh(dci)Rh(cod)]\cdot2Pr^iOH$, 3003–8
Preparation of two mixed-metal carbonyl clusters based on the platinumtriosmium core. *X*-Ray crystal structure of $[Os_3PtH_2(cod)(CH_2)(CO)_9]$ (cod = cyclo-octa-1,5-diene), 3103–6

CYCLO-OCTATETRAENE
The mechanism of fluxionality of $[Ru(\eta^6\text{-}C_8H_8)(\eta^4\text{-}C_7H_8)]$, $[Ru(\eta^4\text{-}C_8H_8)(\eta^4\text{-}C_7H_8)L]$ [L = PEt_3 or $P(OMe)_3$], and $[Os(\eta^6\text{-}C_8H_8)(\eta^4\text{-}C_8H_{12})]$ (C_8H_8 = cyclo-octatetraene, C_7H_8 = norborna-2,5-diene, and C_8H_{12} = cyclo-octa-1,5-diene), 2649–52

CYCLOPALLADATION
Reactions of planar pentadentate ligands with palladium and platinum chlorides. Crystal structures of *cis*-dichlorobis(*N′*-isopropylidenebenzohydrazide)platinum(II) and the cyclopalladated dianion of *N′,N‴*-(pyridine-2,6-diyldiethylidyne)bis(benzohydrazide), 1175–80
Cyclopalladation to form planar tridentate $[N\text{–}C\text{–}N]^-$ intramolecular co-ordination systems involving pyridine donor groups, including ligand synthesis and *X*-ray structural studies., 1477–84

CYCLOPENTADIENIDE
Preparation and spectroscopic study of functionally substituted cyclopentadienides of thallium(I), potassium, rhodium(I), and iridium(I), 2615–20

CYCLOPENTADIENONE
The chemistry of tetraphenylcyclopentadienone complexes of

CYCLOPENTADIENONE (contd)
ruthenium and rhodium: the *X*-ray crystal structure of [Ru{η^5-$C_5Ph_4OC(O)CH(OMe)Ph$}$(CO)_2Cl$], 2815–22
CYCLOPENTADIENYL
Synthesis of [Mo(η-$C_5H_4Pr^i$)$(PR_3)_2H_3$](R_3 = Me_3 or Me_2Ph), [Mo(η-$C_5H_4Pr^i$)$(Pr^i_2PCH_2CH_2PPr^i_2)H_3$], and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of [{Mo(η-$C_5H_4Pr^i$)(μ-Cl)$_2$}$_2$], 9–20
Application of ultraviolet photoelectron spectroscopy to molecular properties. Part 27. Stereochemistry of ligand exchange reactions of titanocene derivatives, 107–10
The addition of isonitriles to [{M$(CO)_2$(η-C_5H_5)}$_2$] (M = Mo or W). Syntheses and properties of [M_2(μ-η^2-CNR)$(CO)_4$(η-C_5H_5)$_2$] (M = Mo or W, R = Me or Bu^t) and [{Mo(CNR)$(CO)_2$(η-C_5H_5)}$_2$] (R = Me or Bu^t). *X*-Ray crystal structures of [Mo_2(μ-η^2-$CNBu^t$)$(CO)_4$(η-(η-C_5H_5)$_2$] and [{Mo($CNBu^t$)$(CO)_2$(η-C_5H_5)}$_2$], 341–8
η-Benzenebis(trimethylphosphine)iron as a precursor to [Fe(η-C_5R_5)$(PMe_3)_2$] (R = H or Me) derivatives: the equilibrium [Fe](PMe_3)Et ⇌ [Fe](η-C_2H_4)H + PMe_3, where [Fe] = Fe(η-C_5Me_5)(PMe_3), 411–6
Reduction–oxidation properties of organotransition-metal complexes. Part 25. The one-electron reduction of cationic cyclopentadienyl–cobalt and –rhodium nitrosyl derivatives, 467–72
Phosphorus-tellurium compounds. Part 3. Cyclopentadienyliron tellurophosphorane complexes: tellurium-bridged iron phosphine compounds?, 541–4
Mössbauer spectroscopic studies on molecular rotation and lattice dynamics of cyclodextrin clathrates of bis(η-cyclopentadienyl)iron(II) and some of its ring-substituted derivatives, 627–32
Reaction of monoborane with tris(cyclopentadienyl)uranium(IV) derivatives, 691–4
Synthesis and characterization of monocyclopentadienyl niobium derivatives. *X*-Ray crystal structure of di-μ-chloro-μ-oxo-bis[dichloro(η-trimethylsilylcyclopentadienyl)niobium(v)], [{Nb(η-$C_5H_4SiMe_3$)Cl_2}$_2$(μ-O)(μ-Cl)$_2$], 953–6
Photoactivation of methane by η^5-cyclopentadienyl and substituted η^5-cyclopentadienyl Group 8 metal dicarbonyl complexes, [M(η^5-C_5R_5)$(CO)_2$] (M = Rh or Ir, R = H or Me), and dicarbonyl(η^5-indenyl)iridium: a matrix isolation study, 1181–90
Synthesis of (η-cyclopentadienyl)(trimethylphosphine)tungsten derivatives and related chemistry, 1611–8
Preparation and characterisation of dicarboxylatobis(η-cyclopentadienyl)tungsten(IV) complexes: a novel and convenient synthesis from [W(η-C_5H_5)$_2H_2$] using molecular oxygen as a hydrogen acceptor, 1857–60
Activation energies for molecular tumbling and cyclopentadienyl rotation in [M(η^5-C_5H_5)(η^4-cod)] (M = Rh or Ir; cod = cycloocta-1,5-diene) and the *X*-ray crystal structure of and bonding in [Rh(η^5-C_5H_5)(η^4-cod)], 1947–52
A trigonal bipyramidal pentaruthenium cluster, [$Ru_5(CO)_3$(μ$_3$-CO)$_3$(η-C_5H_5)$_4$], 2087–8
Axially asymmetric metal alkyls. Part 5. Synthesis and reduction to Zr^{III} species of Group 4 metallepines [ML(η-C_5H_5)$_2$][L = (2-$CHRC_6H_4$)$_2^{2-}$, R = H, M = Ti, Zr, or Hf), isomers of [ZrL(η-C_5H_5)$_2$] (R = $SiMe_3$), and *meso*-[M{1,8-$(CHSiMe_3)_2C_{10}H_6$}(η-C_5H_5)$_2$] (M = Zr or Hf); *X*-ray crystal structures of [TiL(η-C_5H_5)$_2$] (R = H) and λ-and δ-*RS*-[ZrL(η-C_5H_5)$_2$] (R = $SiMe_3$), 2347–58
Subvalent group 14 metal compounds. Part 10. Syntheses and structures of some bis(metallo)plumbylenes [Pb{Mo(R)-$(CO)_3$}$_2$(OC_4H_8)] [R = η-C_5H_5, η-$C_5H_3(SiMe_3)_2$-1,3, or η-C_5Me_5]; *X*-ray crystal structures of [Pb{Mo(η-C_5Me_5)$(CO)_3$}$_2$(OC_4H_8)] and [{Pb[Mo(η-C_5Me_5)$(CO)_3$][Mo(η-C_5Me_5)$(CO)_2$(μ-CO)]}$_2$], 2635–42
CYCLOPENTENEDITHIOCARBOXYLATE
Chemistry of molybdenum. Part 5. Synthesis, spectroscopic and electrochemical properties of bis(2-aminocyclopent-1-ene-1-dithiocarboxylato)(arylimido)dihalogenomolybdenum(VI) complexes, 3043–8
CYCLOPENTENETRIONE
Reactions of cyclobutenedione and cyclopentenetrione with zero-valent nickel triad complexes of isocyanide. *X*-Ray crystal structures of (3—4-η-3,4-diphenylcyclobut-3-ene-1,2-dione)- and (4—5-η-4,5-diphenylcyclopent-4-ene-1,2,3-trione)-tris(t-butyl isocyanide)nickel, 853–8
CYCLOPROPANATION
Carbene complexes in organic synthesis: a new tungstacarbene-promoted alkyne insertion–olefin cyclopropanation reaction, 1041–50
CYCLOSILAZOXANE
Nitrogen-15 and oxygen-17 nuclear magnetic resonance chemical shifts and ^{15}N-^{29}Si, ^{13}C-^{29}Si, and ^{15}N–^1H coupling constants in cyclosilazoxanes, 1593–6
CYCLOTRI(AZATHIENE)
Reactions of $(NSCl)_3$ with Lewis acids and sulphur: a convenient preparation of [S_2N][$SbCl_6$], 1565–8
CYCLOTRIPHOSPHAZATRIENE
Structure and basicity. Part 10. Conductometric and potentiometric titrations of some cyclotriphosphazatrienes in non-aqueous media. The conductivities of alkali-metal acetates on titration with perchloric acid in acetic acid, 925–30
CYCLOTRIPHOSPHAZENE
Studies of phosphazenes. Part 30. Reactions of hexachlorocyclotriphosphazene with aromatic primary amines: interplay of geminal and non-geminal modes of chlorine replacement, 579–84
Studies of phosphazenes. Part 31. Kinetic studies of the reactions of hexachlorocyclotriphosphazene with aromatic primary amines: evidence for the intermediacy of a reactive three-co-ordinated phosphorus(v) species, 585–90
DECABORANE
Identification of the *endo,exo* isomer of 6,9-$(PMe_2Ph)_2$-*arachno*-$B_{10}H_{12}$ by nuclear magnetic resonance spectroscopy, 1573–6
DECATUNGSTATE
Involvement of hydrogen-bonding protons in delocalization of the paramagnetic electron in a single crystal of photoreduced decatungstate, 1597–604
DECOMPOSITION
New decomposition pathways for formyl complexes of ruthenium(II), 2683–6
DEHYDROASCORBIC ACID
Crystal structure of a copper complex of 2-carboxypentonic acid; a decomposition product of dehydroascorbic acid, 2905–10
DELOCALIZATION
Involvement of hydrogen-bonding protons in delocalization of the paramagnetic electron in a single crystal of photoreduced decatungstate, 1597–604
DEUTERIUM
Determination of the deuterium position in DNb_6I_{11}, 431–4
DIALKYL DITHIOPHOSPHATE
Proton and phosphorus-31 nuclear magnetic resonance study of zinc(II) *O,O'*-dialkyl dithiophosphates in solution, 807–14
Reactions of copper(II) with $[S_2P(OR)_2]^-$ (R = Et or Pr^i) and single-crystal *X*-ray studies of Cu[$S_2P(OEt)_2$]·bipy and Cu[$S_2P(OEt)_2$]·$2PPh_3$, 1027–34
DIAMINE
Co-ordination chemistry of higher oxidation states. Part 22. *trans*-Dioxo-osmium(VI) complexes of diamines, diphosphines, and diarsines, 633–8
The base catalysed hydrolysis of *trans*-dichloro-bis(1,2-diaminoethane)cobalt(III) and -(1,3-diaminopropane)cobalt(III) cations: rate-limiting proton transfer, 1775–8
DIARSENE
Oxidation and reduction of phosphorus–phosphorus and arsenic–arsenic double bonds. An electrochemical study of two diphosphenes and a diarsene, 249–52
DIARSINE
The synthesis of mono- and bi-metallic complexes containing the ligand bis(diphenylarsino)methane, 151–4
Co-ordination chemistry of higher oxidation states. Part 22. *trans*-Dioxo-osmium(VI) complexes of diamines, diphosphines, and diarsines, 633–8
DIAZACYCLOHEPTANE
Thermal investigation and stereochemical studies of some cyclic diamine complexes of nickel(II), zinc(II), and cadmium(II) in the solid state, 2089–94
DIAZAPHOSPHOLE
Isomers in *fac*-[ReBr$(CO)_3(L_c)_2$] (L_c = *cis*-3,4-dihydro-2,3,4,5-tetraphenyl-2*H*-1,2,3-diazaphosphole) arising from chirality of phosphorus ligands: separation and *X*-ray structural studies, 21–6
DIAZENIDE
The chemistry of the hydrazido(1−)-ligand. Preparations and crystal structures of

DIAZENIDE (contd)
[Mo(NHNHCO$_2$Me)(NNCO$_2$Me)(S$_2$CNMe$_2$)$_2$], [Mo(NHNMePh)(NNMePh)(S$_2$CNMe$_2$)$_2$]BPh$_4$, and [ReCl$_2$(NHNHCOPh)-(NNHCOPh)(PPh$_3$)$_2$]. Mechanism of formation of substituted hydrazines from [Mo(NNRPh)$_2$(S$_2$CNMe$_2$)$_2$] (R = Me or Ph), 529–40
DIAZO
Formation of germanium–nitrogen double bonds in reactions of the electron-rich germylene bis[bis(trimethylsilyl)amido]-germanium(II) with a range of diazo-compounds, 501–8
Reactions of diazo-compounds with the electron-rich germylene Ge[N(SiMe$_3$)$_2$]$_2$. Characterisation and some addition reactions of [(Me$_3$Si)$_2$N]$_2$GeNNC(COOMe)$_2$: crystal and molecular structure of [(Me$_3$Si)$_2$N]$_2$Ge-[NHN=C(COOMe)$_2$][C(N$_2$)COPh], 2981–8
DIAZOALKANE
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 58. Some reactions of the compounds [MW(μ-CC$_6$H$_4$Me-4)(CO)$_3$(η^5-C$_9$H$_7$)(η^5-C$_2$B$_9$H$_9$Me$_2$)] (M = Mo or W); crystal structures of the complexes [W$_2$(μ-CC$_6$H$_4$Me-4)(CO)$_2$(L)(η^5-C$_9$H$_7$)(η^5-C$_2$B$_9$H$_9$Me$_2$)]-[L = N$_2$C(C$_6$H$_4$Me-4)$_2$ or O], 1221–8
DIAZONIUM
Chelate ring-opening of rhenium–diphosphine complexes by diazonium salts, 3067–72
DIBENZOTETRATHIAFULVALENIUM
X-Ray crystal structure and properties of dibenzotetrathiafulvalenium trichlorodiethylstannate(IV), 1793–6
DICARBAUNDECABORATE
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 51. Alkylidyne(carbaborane)tungsten-gold and -rhodium complexes; crystal structures of [AuW(μ-CR)(CO)$_2$(PPh$_3$)(η^5-C$_2$B$_9$H$_9$Me$_2$)], [RhW(μ-CR)(CO)$_2$(PPh$_3$)$_2$-(η^5-C$_2$B$_9$H$_9$Me$_2$)], and [RhW(μ-CR)(CO)$_2$(PPh$_3$)$_2$-{η^5-C$_2$B$_9$(C$_7$H$_9$)H$_8$Me$_2$}] (R = C$_6$H$_4$Me-4), 61–72
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 52. Synthesis and reactions of the compounds [RuW(μ-CR)(CO)$_3$(η-C$_5$H$_5$)(η^5-C$_2$B$_9$H$_9$Me$_2$)] (R = C$_6$H$_4$Me-4 or Me); crystal structures of [RuW(μ-CC$_6$H$_4$Me-4)(CO)$_3$(η-C$_5$H$_5$)(η^5-C$_2$B$_9$H$_9$Me$_2$)] and [N(PPh$_3$)$_2$][RuW(μ-CC$_6$H$_4$Me-4)(μ-σ,η^5-C$_2$B$_9$H$_8$Me$_2$)(CO)$_3$(η-C$_5$H$_5$)], 73–80
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 53. Alkylidyne(carbaborane)dimetal complexes with tungsten–molybdenum and –tungsten bonds; crystal structures of the compounds [MW(μ-CC$_6$H$_4$Me-4)(CO)$_3$(η^5-C$_9$H$_7$)(η^5-C$_2$B$_9$H$_9$Me$_2$)] [M = Mo (two isomers) or W], 81–90
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 66. Carbaboranetungsten– platinum complexes. Polyhedral rearrangements of a 12-vertex cage system; crystal structures of [PtW(CO)$_2$(PEt$_3$)$_2${η^6-C$_2$B$_9$H$_8$-(CH$_2$C$_6$H$_4$Me-4)Me$_2$}]·CH$_2$Cl$_2$, [PtW(μ-H){μ-σ:η^5-C$_2$B$_9$H$_7$(CH$_2$C$_6$H$_4$Me-4)Me$_2$}(CO)$_2$(PMe$_3$)(PEt$_3$)$_2$], and related compounds, 2219–34
DIHYDROGEN
Water photolysis. Part 2. An investigation of the relative advantages of various components of the sensitiser–electron relay–metal colloid system for the photoproduction of hydrogen from water, and the use of these systems in the photohydrogenation of unsaturated organic substrates, 1299–304
The mechanism of the photochemical production of hydrogen from aqueous solutions of hydridotris(triethylphosphine)platinum(II), 2941–6
DIMETHYLAMINO
The methyl/trifluoromethyl substitution effect in boranes. A gas-phase electron diffraction and vibrational study of bis(dimethylamino)(trifluoromethyl)borane and dimethylaminobis(trifluoromethyl)borane, 1839–46
DIMETHYLOXAMIDE
Crystal and molecular structure of the six-co-ordinate antimony(III) complex SbCl$_3$L, where L = *N*,*N*′-dimethyloxamide. Unusual O,O ligand binding leading to four-membered Sb(μ-O)$_2$Sb ring formation, 263–6
DIMETHYL SULPHOXIDE
Equilibrium and kinetic study of nitric oxide binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 369–72
Some binuclear molybdenum(VI) complexes obtained from condensation reactions: their crystal and molecular structures, 523–8
Thermodynamics of complex formation of silver with amines in dimethyl sulphoxide, 657–60
The relative nucleophilic efficiency of water and dimethyl sulphoxide in the aqueous substitution of aquapentacyanocobaltate(III) and chloropentacyanocobaltate(III), 2311–6
Reactions in mixed non-aqueous solution containing sulphur dioxide. Part 9. Mechanisms of dissolution of metals, oxides, and sulphites, 2995–8
DINITROGEN
Preparation of isocyanide and mixed dinitrogen–isocyanide complexes of rhenium(I) from reactions of *trans*-[ReCl(N$_2$)(PMe$_2$Ph)$_4$], *mer*-[Re(S$_2$PPh$_2$)(N$_2$)(PMe$_2$Ph)$_3$], or *mer*-[Re(S$_2$CNEt$_2$)(N$_2$)(PMe$_2$Ph)$_3$] with methyl isocyanide; crystal structure of *mer*-[Re(S$_2$PPh$_2$)(N$_2$)(CNMe)(PMe$_2$Ph)$_3$], 319–26
Dinitrogen complexes derived from tricarbonyl(η^6-hexaethylbenzene)chromium(0): crystal and molecular structure of μ-dinitrogen-bis[dicarbonyl(η^6-hexaethylbenzene)-chromium(0)]–toluene (1/1), 2789–92
DIOXYGEN
Molecular orbital study of the bridging dioxygen ligand in [Al$_2$(CH$_3$)$_6$(O$_2$)]$^-$, 355–60
Preparation and characterisation of dicarboxylatobis(η-cyclopentadienyl)tungsten(IV) complexes: a novel and convenient synthesis from [W(η-C$_5$H$_5$)$_2$H$_2$] using molecular oxygen as a hydrogen acceptor, 1857–60
Reverse Bohr effect on the oxygen-binding affinity of heme embedded in a bilayer of liposome as a hemoglobin model: pH-induced oxygen uptake and evolution by aqueous synthetic lipid-heme solution, 1917–20
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 8. Irreversible oxidation of trimethylphosphinemanganese(II) dihalides, [MnX$_2$(PMe$_3$)] (X = Cl, Br, or I), to bis(trimethylphosphine)manganese(III) trihalides, [MnX$_3$(PMe$_3$)$_2$]. *X*-Ray crystal structure of *trans*-triiodobis(trimethylphosphine)manganese(III), the first example of a trigonal-bipyramidal manganese(III) tertiary phosphine complex, 1999–2004
Synthesis and characterisation of some new manganese(II) and manganese(III) complexes of tetradentate Schiff-base ligands and their reaction with molecular oxygen, 2009–12
An electron spin resonance study of the dioxygen adduct of bis[2-(2′-pyridylmethylenehydrazonomethyl)phenolato]cobalt(II) and related materials, 2341–6
Iron–ligand bonding properties of synthetic iron–porphyrin complexes with oxygen transporting ability in aqueous media, 2455–8
Dynamics of oxygen binding to modified iron(II)–porphyrin complexes embedded in the bilayer of phospholipid liposome, 2493–6
Oxygen- and carbon monoxide-binding to a lipophilic diporphyrinatocopperiron complex solubilized in an aqueous medium with a micelle, 2963–8
DIPHENYLKETENE
Co-oligomerization of diphenylketene and ethylene promoted by (η^5-cyclopentadienyl)bis(ethylene)cobalt(I): synthesis and *X*-ray structure of a heterocycle containing cobalt(III), 1789–92
DIPHOSPHATE
Intercalation compounds of (diphosphato)oxotungsten(VI), [WO(P$_2$O$_7$)], 609–12
DIPHOSPHAZANE
Electrophilic attack on a series of dinuclear diphosphazane-bridged derivatives of iron by halogens. *X*-Ray crystal structures of [Fe$_2$I(CO)$_5${μ-(MeO)$_2$PN(Et)P(OMe)$_2$}$_2$]PF$_6$ and [Fe$_2$(μ-Br)(CO)$_4${μ-(PhO)$_2$PN(Et)P(OPh)$_2$}$_2$]PF$_6$, 1933–46
DIPHOSPHENE
Oxidation and reduction of phosphorus–phosphorus and arsenic–arsenic double bonds. An electrochemical study of two diphosphenes and a diarsene, 249–52
DIPHOSPHINE
Synthesis of [Mo(η-C$_5$H$_4$Pr^i)(PR$_3$)$_2$H$_3$](R$_3$ = Me$_3$ or Me$_2$Ph), [Mo(η-C$_5$H$_4$Pr^i)(Pr$^i_2$PCH$_2$CH$_2$PPr$^i_2$)H$_3$], and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of [{Mo(η-C$_5$H$_4$Pr^i)(μ-Cl)$_2$}$_2$], 9–20

DIPHOSPHINE (contd)
High-yield syntheses of sulphido triruthenium carbonyl cluster complexes containing bis(diphenylphosphino)methane, 91–6
Mechanism of the reaction between *trans*-$[W(NNH_2)(p\text{-}MeC_6H_4SO_3)(Ph_2PCH_2CH_2PPh_2)_2]^+$ and NEt_3: the influence of the metal on the rates of proton transfer of hydrazido(2−)-ligands, 197–200
Bis(dialkylphosphino)methane complexes of palladium(II) and the molecular structure of $[Pd_2Cl_3(\mu\text{-}Pr^i_2PCH_2PPr^i_2)(\mu\text{-}Pr^i_2PCH_2PHPr^i)]\cdot 2CHCl_3$ containing an unusual phosphidophosphine bridging ligand, 201–6
Bimetallic systems. Part 16. Formation of Group 6 metal(0)–platinum(II) hydride complexes, by ring opening of *fac*- or *mer*-$[M(CO)_3(dppm\text{-}PP')(dppm\text{-}P)]$ (M = Cr, Mo, or W; dppm = $Ph_2PCH_2PPh_2$). Reactions of $[(CO)_3Mo(\mu\text{-}dppm)_2PtH(Cl)]$, 221–6
Reactions of metal carbonyl cluster complexes with multidentate phosphine ligands; the preparation and thermolysis of bis(dimethylphosphino)methane derivatives of $[Ru_3(CO)_{12}]$, and the *X*-ray crystal structure of $[Ru_3(CO)_9(H)(Me_2PCHPMe_2)]$, 277–80
Reactions between bis(*O*-alkyl dithiocarbonato)nickel(II) complexes and phosphines. Formation of a dithiocarbonate complex of nickel(II): $[Ni(S_2CO)(Ph_2PCH_2CH_2PPh_2)]$, 281–4
Co-ordination chemistry of higher oxidation states. Part 22. *trans*-Dioxo-osmium(VI) complexes of diamines, diphosphines, and diarsines, 633–8
Synthesis and structure of μ-[*trans*-1,2-bis(diphenylphosphino)-ethylene]-bis[chlorogold(I)], 677–80
Bis[bis(diphenylphosphino)methane]platinum(II) salts: evidence for the formation of a five-co-ordinate bis[bis(diphenylphosphino)-methane]halogenoplatinum(II) cation, 705–8
Reactivity of the polyhydride complexes $[ReH_5(PPh_3)_3]$, $[ReH_3(dppe)_2]$ (dppe = $Ph_2PCH_2CH_2PPh_2$), $[ReH_3(PPh_3)_3L]$, and $[ReH_4(PPh_3)_3L]PF_6$ (L = MeCN or Bu^{t}NC) towards electrophiles and nucleophiles, 715–20
Bimetallic systems. Part 17. Synthesis of Group 6 metal(0)–copper(I), –silver(I), or –gold(I) complexes with two $Ph_2PCH_2PPh_2$ (dppm) bridging ligands: crystal structure of $[(OC)_2W(\mu\text{-}CO)(\mu\text{-}dppm)_2(\mu\text{-}Cl)Cu]\cdot CH_2Cl_2$, 769–76
Bimetallic systems. Part 18. Nickel(II)–mercury(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$ ligands, 777–80
Mono- and di-carbonyl complexes of molybdenum(II) with mixed bidentate ligands. *X*-Ray crystal structure of $[MoBr_2(CO)(bipy)(Ph_2PCH_2PPh_2)]$ (bipy = 2,2′-bipyridine): a novel type of $[M(bidentate)_2(unidentate)_3]$ structure, 819–24
Trinuclear osmium clusters, containing bidentate $Ph_2PCH_2PPh_2$ and $Me_2PCH_2PMe_2$ ligands: *X*-ray crystal structures of the complexes $[Os_3H(CO)_9(Me_2PCHPMe_2)]$ and $[Os_3H(OH)(CO)_8(Ph_2PCH_2PPh_2)]$, 931–8
The preparation, structure, and reactions of the mononuclear thiolate–hydride complexes $[MoH(SR)(dppe)_2]$ (R = bulkyl alkyl or aryl group, dppe = $Ph_2PCH_2CH_2PPh_2$); *X*-ray structure determination of *cis*-$[MoH(SC_6H_2Pr^i_3\text{-}2,4,6)(dppe)_2]$, 1115–22
Binuclear copper(I) complexes containing bis(diphenylphosphino)methane bridging ligands: crystal structure of $[Cu_2(\mu\text{-}dppm)_2(MeCN)_4][ClO_4]_2$, 1275–8
Bimetallic systems. Part 19. Chromium(0)–, molybdenum(0)–, or tungsten(0)–iron(0) carbonyl complexes containing one or two bridging $Ph_2PCH_2PPh_2$ ligands: crystal structure of $[(OC)_3Mo(\mu\text{-}Ph_2PCH_2PPh_2)_2Fe(CO)_3]$, 1489–94
Bimetallic systems. Part 20. Synthesis of Group 6 metal(0)–iridium(I) or –rhodium(I) complexes bridged by $Ph_2PCH_2PPh_2$, 1495–500
Bimetallic systems. Part 21. Synthesis of vinylidene- and phenylene-bridged complexes of dinickel(II), nickel(II)–palladium(II), and nickel(II)–platinum(II) containing $Ph_2PCH_2PPh_2$ (dppm): crystal structures of $[(SCN)Ni(\mu\text{-}dppm)_2(\mu\text{-}C{=}CH_2)Ni(NCS)]$ and $[ClPd(\mu\text{-}dppm)_2(\mu\text{-}C{=}CCl_2)NiCl]$, 1501–8
Bimetallic systems. Part 22. Molybdenum(0)– and manganese(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the reversible uptake of carbon monoxide: crystal structure of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)Mo(CO)_4]\cdot 0.5C_6H_5Cl$, 1509–14
Synthesis and substitution reactions of a heterodimetallic molybdenum–manganese complex, $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_6]$; *X*-ray crystal structure of $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_4(dppm\text{-}PP')]$, 1557–64
Unsymmetrical bridging of an acetylide ligand in the cation $[Pt_2(C{\equiv}CBu^t)_3(\mu\text{-}dppm)_2]^+$ (dppm = $Ph_2PCH_2PPh_2$): *X*-ray structure determination of the tetraphenylborate salt, 1659–64
Synthesis and reactions of $[(OC)_2Re(\mu\text{-}dppm)_2(\mu\text{-}Cl)_2Rh(\eta^2\text{-}C_2H_4)]$ (dppm = $Ph_2PCH_2PPh_2$): crystal structure of the 1:2 chlorobenzene solvate, 1763–8
The reversible one-electron oxidation of *trans*-$[Mn^IL(CO)(Ph_2PCH_2PPh_2)_2]^{z+}$: synthesis and characterisation of stable manganese(II) derivatives, 1769–74
The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of $[M_2Ru_3(\mu_3\text{-}S)(\mu\text{-}Ph_2PCH_2PPh_2)\text{-}(CO)_9]$ (M = Cu or Au), 1967–76
The synthesis and reactions of palladium–iron carbonyl complexes containing bridging $Ph_2PCH_2PPh_2$ ligands, 2005–8
Pentafluorophenyl complexes of palladium and platinum containing chelating, unidentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 2077–82
Platinum(II) complexes with ligands $(RO)_2PCH_2P(OR)_2$ (R = Me, Et, Ph, or C_6H_4Me-4); crystal structure of *cis,cis*-$[Pt_2Me_4\{\mu\text{-}(EtO)_2PCH_2P(OEt)_2\}_2]$, 2117–24
Some chemistry of the heterobimetallic complexes $[(OC)_3Mo(\mu\text{-}dppm)_2Rh(CO)Cl]$ and $[(OC)_3Mo(\mu\text{-}dppm)_2\text{-}M(CO)_2]PF_6$ (M = Rh or Ir, dppm = $Ph_2PCH_2PPh_2$). Structure of $[(OC)_3Mo(\mu\text{-}dppm)_2(\mu\text{-}C{\equiv}CMe)Rh(CO)]$, 2171–6
Gold-197 Mössbauer spectra of some gold–ruthenium cluster compounds, 2333–6
Determination of the relative signs of $^2J(^{31}P\text{–}^{31}P)$ in complexes of tungsten(0) and molybdenum(0) using two-dimensional [^{31}P–^{31}P]-COSY-45 nuclear magnetic resonance chemical shift correlation, 2401–6
New decomposition pathways for formyl complexes of ruthenium(II), 2683–6
Rhodium(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the formation of a tetranuclear cluster from a heterobimetallic precursor: crystal structures of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)RhCl(CO)]$ and $[Fe_2Rh_2(\mu\text{-}Ph_2PCH_2PPh_2)_2(CO)_8]$, 2751–6
σ-Alkynyl complexes of manganese(I) as η^2-bonding ligands for Group 1B metal–ligand fragments. *X*-Ray crystal structure of $[Mn_2Cu(\mu\text{-}CCBu^t)_2(CO)_6(dppe)_2]PF_6\cdot 0.5CH_2Cl_2$, 2867–74
Electronic structure and redox reactions of vanadium(II) polypyridine, isocyanide, and phosphine complexes, 2881–4
The chemistry of chromium nitrile complexes of 1,2-bis(dimethylphosphino)ethane. *X*-Ray crystal structures of *trans*-$[Cr^{IV}Cl(NEt)(dmpe)_2]CF_3SO_3$, *trans*-$[Cr^{IV}(N{=}CHMe)_2(dmpe)_2][BPh_4]_2$, and *trans*-$[Co^0(H{\cdot}NCMe)(dmpe)^2][BPh_4]_2$, 2947–54
The chemistry of cycloheptatrienyl complexes of molybdenum and tungsten: the synthesis and reactions of some vinylidene and alkynyl derivatives, 3049–54
Chelate ring-opening of rhenium–diphosphine complexes by diazonium salts, 3067–72
Platinum(II)– and palladium(II)–iron(0) complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand. Crystal structures of $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}CO)PtBr_2]$ and $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}I)PtI]\cdot CH_2Cl_2$, 3079–84

DIPHOSPHITE
Preparation and reactivity of halide-bridged binuclear carbonyl manganese and rhenium anions, $[M_2(\mu\text{-}X)(CO)_{8-2n}\{(EtO)_2POP(OEt)_2\}_n]^-$ (X = halide; M = Mn, *n* = 1 or 2; M = Re, *n* = 1). *X*-Ray crystal structure of $[Mn_2Au(\mu\text{-}Br)(CO)_6(\mu\text{-}tedip)(PPh_3)]$, 1551–6

DIPHOSPHONITE
Diplatinum-(II) and -(III) complexes with bridging μ-diphosphonito-*P,P′* or μ-methylenebis(phosphonito)-*P,P′* ligands. Synthesis, structure, and nuclear magnetic resonance analysis of the 'lantern' complexes. Crystal structures of $K_4[Pt_2\{\mu\text{-}CH_2[P(O)OH]_2\}_4]\cdot 6H_2O$ and $K_4[Pt_2\{\mu\text{-}CH_2[P(O)OH]_2\}_4Cl_2]\cdot 8H_2O$, 2769–80

DIPYRIDYL KETONE
Di-2-pyridyl ketone complexes of copper(I): efficient photocatalysts for norbornadiene–quadricyclane conversion, 1797–800
The reaction of di-2-pyridyl ketone with antimony(III) fluoride. Preparation, properties, Mössbauer spectrum and crystal structure of difluoro(methoxydi-2-pyridylmethoxo-*NN′O*)antimony(III), 2337–40

DISELENATE
Preparation, structure, and properties of $[Ni\{S_2C_2(CF_3)_2\}_2]^-$ and $[Ni\{Se_2C_2(CF_3)_2\}_2]^-$ salts with small cations, 1683–6
DISSOCIATION
Electronic *d–d* spectra and ligand-dissociation rates of pentacyanoruthenate(II) ions, 1267–70
DISSOLUTION
Reactions in mixed non-aqueous solution containing sulphur dioxide. Part 9. Mechanisms of dissolution of metals, oxides, and sulphites, 2995–8
DISULPHIDE
Polymeric thiolate complexes of Group 12 metals. Crystal and molecular structures of *catena*-[μ-(3-dimethylammonio-1-propanethiolate)]-dichlorocadmium(II) and bis[3-(dimethylammonio)propyl] disulphide tetrabromocadmate(II), 2391–6
DITHIADIAZATRICYCLODOCOSAHEPTAENE
Copper(II) complexes of the 14-membered *cis*-N_2S_2 macrocycle 14,14,16-trimethyl-2,6-dithia-13,17-diazatricyclo[16.4.0.0^{7,12}]docosa-1(18),7(12),8,10,16,19,21-heptaene, 963–4
DITHIOCARBAMATE
The chemistry of the hydrazido(1−)-ligand. Preparations and crystal structures of $[Mo(NHNHCO_2Me)(NNCO_2Me)(S_2CNMe_2)_2]$, $[Mo(NHNMePh)(NNMePh)(S_2CNMe_2)_2]BPh_4$, and $[ReCl_2(NHNHCOPh)(NNHCOPh)(PPh_3)_2]$. Mechanism of formation of substituted hydrazines from $[Mo(NNRPh)_2(S_2CNMe_2)_2]$ (R = Me or Ph), 529–40
Dimeric tungsten(V) compounds containing $[(S)W(\mu\text{-}S)_2W(S)]^{2+}$. Syntheses and structures of di-μ-sulphido-bis[(diethyl dithiophosphato-*SS'*)sulphidotungsten(V)] and di-μ-sulphido-bis[(*N,N*-diethyldithiocarbamato-*SS'*)sulphidotungsten(V)], 1163–8
Mixed chloro-, *NN*-dialkyldithiocarbamato-, and *S*-alkyl *NN*-dialkyldithiocarbamate complexes of platinum(II). Crystal structure of $[PtCl(S_2CNEt_2)(MeS_2CNEt_2)]$, 1823–6
Synthesis and crystal structure of tris(morpholinecarbodithioato-*S,S'*)manganese(IV) pentaiodide–dichloromethane (2/1), 2407–10
Dithiocarbamates of tungsten(III) and tungsten(IV), 2531–6
Mixed-metal phosphinito complexes of platinum(II) and palladium(II) with lanthanide and actinide elements. The single-crystal *X*-ray structure determination of $[UO_2(OH_2)\{(OPPh_2)_2Pd(S_2CNEt_2)\}_2]$, 2853–6
Mean phosphorus– and arsenic–sulphur bond enthalpies in diethyldithiocarbamate chelate complexes, 2955–8
DITHIOCARBAZATE
Electron-transfer platinum complexes of esters of dithiocarbazic acid. The crystal and molecular structure of $[Pt\{N(CH_2Ph)NC(S)SMe\}_2]$, 1035–40
DITHIOCARBONATE
Reactions between bis(*O*-alkyl dithiocarbonato)nickel(II) complexes and phosphines. Formation of a dithiocarbonate complex of nickel(II): $[Ni(S_2CO)(Ph_2PCH_2CH_2PPh_2)]$, 281–4
Preparation of isocyanide and mixed dinitrogen–isocyanide complexes of rhenium(I) from reactions of *trans*-$[ReCl(N_2)(PMe_2Ph)_4]$, *mer*-$[Re(S_2PPh_2)(N_2)(PMe_2Ph)_3]$, or *mer*-$[Re(S_2CNEt_2)(N_2)(PMe_2Ph)_3]$ with methyl isocyanide; crystal structure of *mer*-$[Re(S_2PPh_2)(N_2)(CNMe)(PMe_2Ph)_3]$, 319–26
DITHIOCARBOXYLATE
Linear-chain platinum(II) dithiocarboxylates: crystal structure of tetrakis(phenyldithioacetato)diplatinum(II), 35–40
DITHIOLATE
Preparation, structure, and properties of $[Ni\{S_2C_2(CF_3)_2\}_2]^-$ and $[Ni\{Se_2C_2(CF_3)_2\}_2]^-$ salts with small cations, 1683–6
Preparation and crystal structures of bis(tetraphenylarsonium) tris(oxalato)technetate(IV), and tetraphenylarsonium tris(benzene-1,2-dithiolato)technetate(V): octahedral *versus* trigonal-prismatic geometry for tris-bidentate complexes of technetium, 2305–10
DITHIONITRYL
Reactions of $(NSCl)_3$ with Lewis acids and sulphur: a convenient preparation of $[S_2N][SbCl_6]$, 1565–8
DITHIO-OXAMIDE
Dithio-oxamides as ligands: crystal structures and vibrational analyses of bis(*N,N'*-dicyclohexyldithio-oxamidato)palladium(II), bis(*N,N'*-dibenzyldithio-oxamide)copper(II) diperchlorate, and dichloro(*N,N'*-dimethyldithio-oxamide)zinc(II) and the carbon-13 nuclear magnetic resonance spectra of related compounds, 1921–8
DITHIOPHOSPHATE
Dimeric tungsten(V) compounds containing $[(S)W(\mu\text{-}S)_2W(S)]^{2+}$. Syntheses and structures of di-μ-sulphido-bis[(diethyl dithiophosphato-*SS'*)sulphidotungsten(V)] and di-μ-sulphido-bis[(*N,N*-diethyldithiocarbamato-*SS'*)sulphidotungsten(V)], 1163–8
Crystal and molecular structure of $[\{Ag[S_2P(OEt)_2]\cdot PPh_3\}_2]$, 1569–72
DITHIOPHOSPHINATE
The crystal structures of the diphenyldithiophosphinates of antimony(III) and bismuth(III): $M(S_2PPh_2)_3$ (M = Sb or Bi), 145–50
The crystal structure of bismuth diethyldithiophosphinate–benzene (1/1), $Bi(S_2PEt_2)_3\cdot C_6H_6$, 1257–60
DODECABORANE
Reactions of heteroboranes. Synthesis of $[2,2\text{-}(PPh_3)_2\text{-}1,2\text{-}SePtB_{10}H_{10}]\cdot CH_2Cl_2$, its crystal and molecular structure and that of $SeB_{11}H_{11}$, 699–704
DODECABORATE
Ruthenium(II) complexes of *closo* dodecaboranyl anions and the molecular structure of the *pileo* thirteen-vertex ruthenaborane $[(PPh_3)_2ClRuB_{12}H_{11}(NEt_3)]$, 451–6
ELECTROCHEMISTRY
Magnetic and electrochemical studies on the series $[Cr_nFe_{3-n}(\mu_3\text{-}O)(gly)_6(H_2O)_3][NO_3]_7\cdot xH_2O$(gly = glycine; n = 0, 1, 2, or 3) and on the mixed-valence complex $[Fe^{II}Fe^{III}_2(\mu_3\text{-}O)(gly)_6(H_2O)_3]Cl_6$, 97–100
Preparation and magnetism of the binuclear iron(II) complexes $[\{Fe(C_9H_{21}N_3)X_2\}_2]$ (X = NCS, NCO, or N_3) and their reaction with NO. Crystal structures of $[\{Fe(C_9H_{21}N_3)(NCS)_2\}_2]$ and $[Fe(C_9H_{21}N_3)(NO)(N_3)_2]$, 187–92
Syntheses, spectroscopic, and electrochemical properties of dinuclear copper(II) complexes bridged by a single thiolate sulphur atom; *X*-ray crystal structures of $[Cu_2(L^1)_2(SC_6H_4Cl\text{-}4)]\text{-}[PF_6]$ and $[Cu_2(L^2)_2(SC_6H_4CH_3\text{-}4)][ClO_4]$, 241–8
Oxidation and reduction of phosphorus–phosphorus and arsenic–arsenic double bonds. An electrochemical study of two diphosphenes and a diarsene, 249–52
Structure and properties of tris(quinoxaline-2,3-dithiolato)-molybdate(IV), 297–302
Studies on transition-metal macrocyclic complexes. Single-crystal *X*-ray structure and electrochemistry of the bis-macrocyclic complex $[Cu(L)_2][ClO_4]_2\cdot 2H_2O$ (L = 1,4,7-triazacyclononane), 373–8
Reduction–oxidation properties of organotransition-metal complexes. Part 25. The one-electron reduction of cationic cyclopentadienyl–cobalt and –rhodium nitrosyl derivatives, 467–72
Redox behaviour of the iron–sulphur cluster $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{2+}$. Synthesis and crystal structure of the new paramagnetic monopositive species $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^+$ as its $[PF_6]^-$ salt, 831–6
Some investigations of $S_5N_5^+$ compounds; *X*-ray crystal structure of $[S_5N_5]Cl$, 915–24
New dicyclopentadienyl phosphine, phosphite, and acetylene niobium(III) complexes. Crystal structure of $[Nb(\eta\text{-}C_5H_4SiMe_3)_2\text{-}Cl(PhC{\equiv}CPh)]$, 975–80
High nuclearity osmium cluster radicals, 1051–4
A structural comparison of bis(quinoxaline-2,3-dithiolato)cuprate complexes, 1083–8
Synthesis, spectroscopic, and electrochemical properties of bis(2,2'-bipyridyl)ruthenium compounds of some pyridyl-1,2,4-triazoles, 1389–96
Reduction–oxidation properties of organotransition-metal complexes. Part 26. The synthesis of paramagnetic triazenido-and aryl(1-aryliminoethyl)amido-bridged dirhodium carbonyl derivatives containing the $[Rh_2]^{3+}$ core, 1403–8
Reactions of chromium(III) complexes of 1,10-phenanthroline, 2,2'-bipyridyl, and oxalate with the pulse radiolytically generated aquated electron, zinc(I), and cadmium(I), 1425–30
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato-molybdenum and -tungsten nitrosyls. Part 4. The effects of ligating atom type on the reduction potentials of monometallic complexes, 1733–6
The reversible one-electron oxidation of *trans*-$[Mn^IL(CO)(Ph_2PCH_2PPh_2)_2]^{2+}$: synthesis and characterisation of stable manganese(II) derivatives, 1769–74

ELECTROCHEMISTRY (contd)
Electrochemical reactions of dichlorobis(aryl isocyanide)platinum(II) complexes, 1833–8
Co-ordination chemistry of higher oxidation states. Part 23. Synthesis and properties of tetrahalogenoiridium(IV) complexes, $[IrL_2X_4]$ [X = Cl or Br; L = pyridine, PR_3, AsR_3, SbR_3, SR_2, or SeR_2]. Crystal and molecular structure of *trans*-$[Ir(AsEt_3)_2Br_4]$, 1901–10
Disproportionation, electrochemistry, and electronic coupling involving mononuclear and binuclear cobalt phthalocyanine derivatives, 2511–8
Dithiocarbamates of tungsten(III) and tungsten(IV), 2531–6
Bicyclam [6,6′-bi(1,4,8,11-tetra-azacyclotetradecane)]: a ditopic receptor for homo- and hetero-bimetallic complexes, 2631–4
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato molybdenum and tungsten nitrosyls. Part 6. Synthesis and electrochemistry of complexes containing saturated heterocyclic amide ligands. *X*-Ray crystal and molecular structure of $[\{Mo(NO)[HB(Me_2pz)_3]Cl\}_2(NC_4H_8N)]$, 2653–60
Electronic structure and redox reactions of vanadium(II) polypyridine, isocyanide, and phosphine complexes, 2881–4
Chemistry of molybdenum. Part 5. Synthesis, spectroscopic and electrochemical properties of bis(2-aminocyclopent-1-ene-1-dithiocarboxylato)(arylimido)dihalogenomolybdenum(VI) complexes, 3043–8
Structure and electrochemical properties of bis(pyridine)cobalt(III) complexes of *N*,*N*′-ethylenebis(salicylideneimine) homologues with an aryl substituent on the ethylene backbone. Crystal structures of $[Co(salphen)(py)_2]PF_6{\cdot}0.5Et_2O$ and $[Co(salcppn)(py)_2]PF_6{\cdot}2py$ (py = pyridine), 3055–60
Synthesis, characterisation, and photochemical properties of a series of ruthenium containing metallopolymers based on poly-*N*-vinylimidazole, 3073–8
Syntheses and crystal structures of complexes $[M_2R_2]$, [M = Cu, Ag, or Au; R = 2-$C(SiMe_3)_2C_5H_4N$] and $[Cu_4R'_4]$[R′ = 2-$CH(SiMe_3)C_5H_4N$]; electrochemical generation of $[Cu_2R_2]^{2+}$, 3085–92

ELECTRON DIFFRACTION
Electron diffraction study of the molecular structure of *N*-methylbis(trimethylsilyl)amine in the gas phase, 785–8
The methyl/trifluoromethyl substitution effect in boranes. A gas-phase electron diffraction and vibrational study of bis(dimethylamino)(trifluoromethyl)borane and dimethylaminobis(trifluoromethyl)borane, 1839–46
Molecular structures of some (dimethylamino)halogenosilanes in the gas phase by electron diffraction and the crystal and molecular structures of mono-and dichloro(dimethylamino)-silane by *X*-ray diffraction at low temperatures, 3035–42
The structure of bis(dimethylamino)silane in the gas phase, determined by electron diffraction, 3061–6

ELECTRONIC
Linear-chain platinum(II) dithiocarboxylates: crystal structure of tetrakis(phenyldithioacetato)diplatinum(II), 35–40
Electronic absorption spectra and photochemical reactivity of Group 5 metal alkyl compounds: photochemical α-hydrogen abstraction, 163–8
Electron spin resonance and electronic spectra and crystal and molecular structures of copper(II) amino acid complexes, 457–66
Electron spin resonance and optical spectroscopic studies on Cu^{2+} and Co^{2+} in $Cs_2[Zn(N_3)_4]$, 939–42
Electronic *d*–*d* spectra and ligand-dissociation rates of pentacyanoruthenate(II) ions, 1267–70
Concentrated aqueous solutions and liquid hydrates of zinc chloride: a study of their solvent behaviour using intershell, charge-transfer, and *d*–*d* adsorption spectra, 1485–8
On the induced circular dichroism of *d*–*d* transitions of a 14-membered macrocyclic tetra-amine nickel(II) complex in β-cyclodextrin, 1801–4
Crystal and molecular structure of chromium(II) sulphate pentahydrate and single-crystal electronic spectra and bonding of $CrSO_4{\cdot}5H_2O$, $CuSO_4{\cdot}5H_2O$, and $CuSO_4{\cdot}5D_2O$, 1817–22
Molten lithium carbonate–sodium carbonate–potassium carbonate eutectic: electronic spectroscopy of first-row transition metal compounds, 2329–32
Characterizations of the redox orbitals of mixed-ligand 2,2′-bipyridine–2-(2′-pyridyl)quinoline ruthenium(II) complexes, 2467–72
Disproportionation, electrochemistry, and electronic coupling involving mononuclear and binuclear cobalt phthalocyanine derivatives, 2511–8
New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states, 2621–30
Electronic structure and redox reactions of vanadium(II) polypyridine, isocyanide, and phosphine complexes, 2881–4

ELECTRON TRANSFER
Oxidation and reduction of phosphorus–phosphorus and arsenic–arsenic double bonds. An electrochemical study of two diphosphenes and a diarsene, 249–52
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid, 573–8
A kinetic and mechanistic study of the formation of a diaryl ditelluride in solution by the sulphite reduction of the aryltellurium trichloride, 985–90
Electron-transfer platinum complexes of esters of dithiocarbazic acid. The crystal and molecular structure of $[Pt\{N(CH_2Ph)NC(S)SMe\}_2]$, 1035–40
Reaction paths and kinetics in the Cr^{2+} reduction of the Mo^{V}_2 and Mo^{IV}_3 ions $[Mo_2O_4(H_2O)_6]^{2+}$ and $[Mo_3O_4(H_2O)_9]^{4+}$, 1397–402
Reactions of chromium(III) complexes of 1,10-phenanthroline, 2,2′-bipyridyl, and oxalate with the pulse radiolytically generated aquated electron, zinc(I), and cadmium(I), 1425–30
Kinetic studies on the oxidation of ethylenediaminebis(pyridine)- and bis(ethylenediamine)-platinum(II) complexes by hexachloroiridate(IV), 1473–6
The reversible one-electron oxidation of *trans*-$[Mn^{I}L(CO)(Ph_2PCH_2PPh_2)_2]^{z+}$: synthesis and characterisation of stable manganese(II) derivatives, 1769–74
Electrochemical reactions of dichlorobis(aryl isocyanide)platinum(II) complexes, 1833–8
Photochemical reduction of carbon dioxide to carbon monoxide in water using a nickel(II) tetra-azamacrocycle complex as catalyst, 2105–10
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid (1987, 573), 2291–2
Kinetics of reduction of tris(ethylenediamine)cobalt(III) by ethylenediamine-*NNN*′*N*′-tetra-acetatochromate(II), 2485–8
Kinetic studies on 1:1 electron-transfer reactions involving blue copper proteins. Part 14. Reactions of poplar plastocyanin with inorganic complexes, 2573–8
Reduction–oxidation properties of organotransition-metal complexes. Part 27. Mixed-valence triazenido-bridged di-iridium compounds, 2737–40
Electronic structure and redox reactions of vanadium(II) polypyridine, isocyanide, and phosphine complexes, 2881–4

ENERGY STORAGE
Di-2-pyridyl ketone complexes of copper(I): efficient photocatalysts for norbornadiene–quadricyclane conversion, 1797–800

EQUILIBRIUM
Equilibrium and kinetic study of nitric oxide binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 369–72
Tungsten(VI) equilibria: a potentiometric and calorimetric investigation, 1701–6
Studies on transition-metal–peptide complexes. Part 13. Copper(II) and nickel(II) complexes of amino acids and peptides containing a thioether group, 1717–20
An electron spin resonance study of the equilibrium between tetrahalogeno-and pentahalogeno-nitridotechnetate(VI) ions in solution, 1721–6
Complex formation of beryllium(II) with salicylate and hydroxide ions in 1 mol dm^{-3} sodium perchlorate aqueous solution at 25 °C, 1853–6
Beryllium(II) hydrolysis in 3.0 mol dm^{-3} perchlorate, 2431–8
Studies of metal carbonate equilibria. Part 15. The beryllium(II)–water–carbon dioxide(g) system in acidic 3.0 mol dm^{-3} perchlorate media, 2439–44
Studies of metal carbonate equilibria. Part 16. The beryllium(II)–water–carbon dioxide(g) system in neutral-to-alkaline 3.0 mol dm^{-3} perchlorate media at 25 °C, 2445–50

E.S.R.
The single-crystal electron spin resonance spectrum of oxobis(2-methylquinolin-8-olato)vanadium(IV) as a pure compound and diluted in chlorobis(2-methylquinolin-8-olato)gallium(III), 111–8
Hydrogen halide radical anions, 421–6
Electron spin resonance and electronic spectra and crystal and

E.S.R. (contd)
molecular structures of copper(II) amino acid complexes, 457–66
The radical cations of sulphur ($S_8^{\bullet+}$) and tetrasulphur tetranitride ($S_4N_4^{\bullet+}$): a radiation–electron spin resonance study, 729–32
Dicopper(II) complexes of a 28-membered N_8 macrocycle. Evidence for proton transfer from co-ordinated secondary amine to thiocyanate, leading to formation of an isothiocyanic acid complex, 873–80
Electron spin resonance and optical spectroscopic studies on Cu^{2+} and Co^{2+} in $Cs_2[Zn(N_3)_4]$, 939–42
Involvement of hydrogen-bonding protons in delocalization of the paramagnetic electron in a single crystal of photoreduced decatungstate, 1597–604
An electron spin resonance study of the equilibrium between tetrahalogeno-and pentahalogeno-nitridotechnetate(VI) ions in solution, 1721–6
An electron spin resonance study of the copper(II) chelate of 2-(2′-pyridylmethylenehydrazonomethyl)phenol and related compounds, 1755–62
An electron spin resonance study of the dioxygen adduct of bis[2-(2′-pyridylmethylenehydrazonomethyl)phenolato]cobalt(II) and related materials, 2341–6
Characterizations of the redox orbitals of mixed-ligand 2,2′-bipyridine–2-(2′-pyridyl)quinoline ruthenium(II) complexes, 2467–72
An electron spin resonance study of frozen aqueous solutions containing 5,10,15,20-tetrakis(*N*-methyl-4′-pyridinio)-porphyrinatocobalt(II), 3099–102

ESTER
Studies on borate esters. Part 8. Interactions of cations with oxyacid anion-bridged esters of D-glucarate in alkaline media, 2051–8

ETHANOIC ACID
Fluorine-19 nuclear magnetic resonance studies of tetraethylammonium fluoride in glacial ethanoic acid, 1271–4

ETHENYL
Synthesis and characterization of new heterobimetallic iron–cobalt complexes with ethenyl bridges $CR^2{=}CHR^1$ [R^1 = H or C(O)OMe, R^2 = C(O)OMe or C(O)OEt]. *X*-Ray structure of $[FeCo(CO)_6\{\mu\text{-}MeOC(O)C{=}C(H)C(O)OMe\}]$, 1619–22

ETHOXIDE
Some binuclear molybdenum(VI) complexes obtained from condensation reactions: their crystal and molecular structures, 523–8

ETHYL
η-Benzenebis(trimethylphosphine)iron as a precursor to $[Fe(\eta\text{-}C_5R_5)(PMe_3)_2]$ (R = H or Me) derivatives: the equilibrium $[Fe](PMe_3)Et \rightleftharpoons [Fe](\eta\text{-}C_2H_4)H + PMe_3$, where $[Fe] = Fe(\eta\text{-}C_5Me_5)(PMe_3)$, 411–6
X-Ray crystal structure and properties of dibenzotetrathiafulvalenium trichlorodiethylstannate(IV), 1793–6

ETHYLENE
Nucleophilic attack upon ethylene co-ordinated to mercury(II): a molecular-orbital study concerning the origin of the acceleration by Hg^{II}, 361–8
η-Benzenebis(trimethylphosphine)iron as a precursor to $[Fe(\eta\text{-}C_5R_5)(PMe_3)_2]$ (R = H or Me) derivatives: the equilibrium $[Fe](PMe_3)Et \rightleftharpoons [Fe](\eta\text{-}C_2H_4)H + PMe_3$, where $[Fe] = Fe(\eta\text{-}C_5Me_5)(PMe_3)$, 411–6
Synthesis and reactions of $[(OC)_2Re(\mu\text{-}dppm)_2(\mu\text{-}Cl)_2Rh(\eta^2\text{-}C_2H_4)]$ (dppm = $Ph_2PCH_2PPh_2$): crystal structure of the 1:2 chlorobenzene solvate, 1763–8
Co-oligomerization of diphenylketene and ethylene promoted by (η^5-cyclopentadienyl)bis(ethylene)cobalt(I): synthesis and *X*-ray structure of a heterocycle containing cobalt(III), 1789–92
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 9. The reaction of ethylene with $[MnX_2L]$ (X = Cl, Br, or I; L = PPr^n_3, PBu^n_3, $PPhMe_2$, $PPhEt_2$, or PPh_3) in tetrahydrofuran solution, 2241–4

ETHYLENEDIAMINE
Metal–ligand bonding parameters and magnetic properties of some previously reported tetragonal nickel(II) complexes, 825–30
Thermally induced phase transition and *X*-ray crystal structure of bis(*N*,*N*′-dimethyl-1,2-ethanediamine)di-isothiocyanatonickel(II), 997–1000
Kinetic studies on the oxidation of ethylenediaminebis(pyridine)- and bis(ethylenediamine)-platinum(II) complexes by hexachloroiridate(IV), 1473–6
Chiral recognition in ion pairing of an optically active tris(ethylenediamine)cobalt(III) cation and application to chromatographic resolution of metal complex anions, 1977–82

ETHYLENEDIAMINETETRA-ACETATE
Kinetics of reduction of tris(ethylenediamine)cobalt(III) by ethylenediamine-*NNN*′*N*′-tetra-acetatochromate(II), 2485–8

ETHYLPYRIDINE
Studies in Werner clathrates. Part 4. Structures of tetrakis(4-ethylpyridine)di-isothiocyanatonickel(II) and its clathrates with *p*-, *m*-, and *o*-xylene, carbon disulphide, and carbon tetrachloride, 2125–40

EXAFS
A structural comparison of the two polymorphs of $Sr_2Co_2O_5$ by cobalt *K*-edge extended *X*-ray absorption fine structure spectroscopy, 2359–64

EXCHANGE
Synthesis of $[Mo(\eta\text{-}C_5H_4Pr^i)(PR_3)_2H_3]$($R_3$ = Me_3 or Me_2Ph), $[Mo(\eta\text{-}C_5H_4Pr^i)(Pr^i{}_2PCH_2CH_2PPr^i{}_2)H_3]$, and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of $[\{Mo(\eta\text{-}C_5H_4Pr^i)(\mu\text{-}Cl)_2\}_2]$, 9–20
Application of ultraviolet photoelectron spectroscopy to molecular properties. Part 27. Stereochemistry of ligand exchange reactions of titanocene derivatives, 107–10
Thermal and photochemical behaviour of organotetraruthenium clusters: solution structures and dynamics of phosphine-substituted derivatives, 349–54
Menthyl-substituted organophosphorus compounds. Part 6. Preparation of P(L-men)RCl and P(L-men)R(S)Cl (R = Me, Et, Pr^i Bu^t, or Ph; men = *cyclo*-C_6H_9-2-Pr^i-5-Me) and nuclear magnetic resonance studies of configuration and halogen exchange, 795–806
Exchange of cadmium into the sodium and ammonium forms of the natural zeolites clinoptilolite, mordenite, and ferrierite, 1911–6
Lamellar inorganic ion exchangers. Proton–copper(II) ion exchange in γ-titanium bis(hydrogenphosphate), 2045–50
Ligand intra- and inter-molecular exchange processes in dioxopentakis(trimethylurea)uranium(VI). A proton nuclear magnetic resonance study, 2885–8

EXTRACTION
Synthesis of *SS*′-bis(4′-benzo[15-crown-5])dithioglyoxime and its complexes with copper(II), nickel(II), cobalt(II), cobalt(III), palladium(II), platinum(II), and platinum(IV), 1827–32

FILM
Synthesis, characterisation, and photochemical properties of a series of ruthenium containing metallopolymers based on poly-*N*-vinylimidazole, 3073–8

FLUORIDE
Preparation of the nitrenes $ReF_5(NF)$, $ReF_5(NCl)$, and $ReF_5(NBr)$. *X*-Ray crystal structures of $ReF_5(NF)$ and $ReF_5(NCl)$, 567–72
Rhenium carbonyl fluorides: preparation of $[Re(CO)_6][ReF_6]$ and some reactions of $[Re(CO)_4F{\cdot}ReF_5]$, 741–6
Direct synthesis of alkali trifluorocobaltate(II) monohydrates, $A[CoF_3]{\cdot}H_2O$ (A = NH_4, Na, or K). First synthesis and structural assessment of mixed-ligand fluoro complexes of cobalt(II), $[Co(NH_3)_4F_2]$ and $A[CoF(SO_4)(H_2O)_2]$ (A = NH_4, Na, or K), 1055–8
Fluorine-19 nuclear magnetic resonance studies of tetraethylammonium fluoride in glacial ethanoic acid, 1271–4
A quantitative approach to host–guest interactions for matrix-isolated alkali-metal salts of hexafluoroanions and perchlorates, 1521–8
The preparation and crystal structure of $5SbF_3{\cdot}3SbF_5$, 1989–98
Attempted synthesis of osmium(VI) and iridium(VI) thiofluorides; the preparation of $OsF_5{\cdot}SF_4$ and $IrF_5{\cdot}SF_4$, 2285–8
Metal(II) hexafluoroantimonates: preparation and characterization of $MF_2{\cdot}2SbF_5$ (M = Mg, Ni, Zn, Fe, Co, Cu, Cr, Ag, Cd, or Pb) and the *X*-ray structure determination of $AgF_2{\cdot}2SbF_5$, 2379–84
Aqua(2,2′-bipyridyl)difluorocopper(II) dihydrate: *X*-ray structure reveals short hydrogen bonds and other unusual features, 2397–400
Preparation, crystal and molecular structure, and antimony-121 Mössbauer spectrum of trifluoro(1,10-phenanthroline)(thiourea)-antimony(III), 2451–4

FLUORINE-19
Fluorine-19 nuclear magnetic resonance studies of tetraethylammonium fluoride in glacial ethanoic acid, 1271–4

FLUOROPHOSPHATE
Direct synthesis of ammonium monofluorophosphate monohydrate,

FLUOROPHOSPHATE (contd)
$[NH_4]_2[PO_3F]\cdot H_2O$, and potassium monofluorophosphate, $K_2[PO_3F]$, 477–80
FLUOROPHOSPHINE
Instant ligands. Part 2. Preparation of some bidentate ligands containing PF_2O Groups from aromatic hydroxy compounds and their reactions to form molybdenum and tungsten complexes, 3023–8
FLUXIONALITY
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au), 379–90
Heteronuclear 1,3-intramolecular metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of 2-thia-4-selenapentane. A dynamic nuclear magnetic resonance investigation, 757–60
Carbon-13 and phosphorus-31 nuclear magnetic resonance studies of the fluxional behaviour of the complexes $[Fe_3(CO)_{10}\{P(OMe)_3\}_2]$ and $[Fe_3(CO)_{10}\{P(OMe)_3\}\{P(OPh)_3\}]$, 1249–52
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au) (1987, 379), 1283–4
The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of $[M_2Ru_3(\mu_3\text{-}S)(\mu\text{-}Ph_2PCH_2PPh_2)(CO)_9]$ (M = Cu or Au), 1967–76
Preparation and spectroscopic study of functionally substituted cyclopentadienides of thallium(I), potassium, rhodium(I), and iridium(I), 2615–20
The mechanism of fluxionality of $[Ru(\eta^6\text{-}C_8H_8)(\eta^4\text{-}C_7H_8)]$, $[Ru(\eta^4\text{-}C_8H_8)(\eta^4\text{-}C_7H_8)L]$ [L = PEt_3 or $P(OMe)_3$], and $[Os(\eta^6\text{-}C_8H_8)(\eta^4\text{-}C_8H_{12})]$ (C_8H_8 = cyclo-octatetraene, C_7H_8 = norborna-2,5-diene, and C_8H_{12} = cyclo-octa-1,5-diene), 2649–52
A two-dimensional nuclear magnetic resonance exchange study of the fluxionality of $[PtIMe_3(MeSCH_2CH_2SEt)]$, 2695–702
FORMYL
New decomposition pathways for formyl complexes of ruthenium(II), 2683–6
Hydride addition reactions of the olefin complexes $[Fe(\eta^5\text{-}C_5H_5)(CO)_2(olefin)][BF_4]$; formyl formation as the kinetically preferred process; regio- and stereo-selectivity during addition to the co-ordinated olefins, 2857–62
FORMYLMETHYLIDENE
Decarboxylation of vinylene carbonate at triosmium clusters; *X*-ray crystal structure of the μ_3-formylmethylidene compound $[Os_3(CHCHO)(CO)_{10}]$, 1861–4
FREE RADICAL
New decomposition pathways for formyl complexes of ruthenium(II), 2683–6
GALLIUM
The single-crystal electron spin resonance spectrum of oxobis(2-methylquinolin-8-olato)vanadium(IV) as a pure compound and diluted in chlorobis(2-methylquinolin-8-olato)gallium(III), 111–8
Synthesis and properties of gallium tetramethylammonium sodalite, $NMe_4GaSi_5O_{12}$, 623–6
The chemistry of trichloro[tris(trimethylsilyl)methyl] and trichloro[tris(dimethylphenylsilyl)methyl] complexes of gallium, indium, and thallium. Crystal and molecular structures of $[Li(thf)_2(\mu\text{-}Cl)_2GaCl\{C(SiMe_2Ph)_3\}]$·thf, $[Li(thf)_3(\mu\text{-}Cl)InCl_2\{C(SiMe_3)_3\}]$, and $[\{(Me_3Si)_3C\}In(\mu\text{-}Cl)_2\{\mu\text{-}Fe(CO)_4\}In\{C(SiMe_3)_3\}]$ (thf = tetrahydrofuran), 747–56
GERMANE
Reactions of diazo-compounds with the electron-rich germylene $Ge[N(SiMe_3)_2]_2$. Characterisation and some addition reactions of $[(Me_3Si)_2N]_2GeNNC(COOMe)_2$: crystal and molecular structure of $[(Me_3Si)_2N]_2Ge[NHN{=}C(COOMe)_2][C(N_2)COPh]$, 2981–8
GERMANIUM
Formation of germanium–nitrogen double bonds in reactions of the electron-rich germylene bis[bis(trimethylsilyl)amido]-germanium(II) with a range of diazo-compounds, 501–8
Chemistry of 2,3-dihydro-2-metallaphenalenes. Part 2. Formation of $\overline{CHMMe_2C(MMe_3)C_{10}H_6}]^x$ *via* base decomposition of $[1,8\text{-}(Me_3MCH_2)_2C_{10}H_6]^y$ (M = Ge, $x = 2-$, $y = 0$; M = P, $x = 0$, $y = 2+$); crystal structures of the novel bis-ylide $\overline{CHPMe_2C(PMe_3)C_{10}H_6}$ and of a distorted *peri*-ylide, 1-Me-8-$(Me_3PCH)C_{10}H_6$, 1647–54
Aspects of germanium-73 nuclear magnetic resonance spectroscopy, 2365–72
Reactions of diazo-compounds with the electron-rich germylene $Ge[N(SiMe_3)_2]_2$. Characterisation and some addition reactions of $[(Me_3Si)_2N]_2GeNNC(COOMe)_2$: crystal and molecular structure of $[(Me_3Si)_2N]_2Ge[NHN{=}C(COOMe)_2][C(N_2)COPh]$, 2981–8
GERMANIUM-73
Aspects of germanium-73 nuclear magnetic resonance spectroscopy, 2365–72
GLUCARATE
Studies on borate esters. Part 8. Interactions of cations with oxyacid anion-bridged esters of D-glucarate in alkaline media, 2051–8
GLYCINATE
Synthesis, characterisation, and aquation kinetics of [*N*-(*o*-carboxyphenyl)iminodiacetato](glycinato)chromate(III) complex ion in acidic media, 2095–8
GLYCINE
Magnetic and electrochemical studies on the series $[Cr_nFe_{3-n}(\mu_3\text{-}O)(gly)_6(H_2O)_3][NO_3]_7\cdot xH_2O$(gly = glycine; n = 0, 1, 2, or 3) and on the mixed-valence complex $[Fe^{II}Fe^{III}_2(\mu_3\text{-}O)(gly)_6(H_2O)_3]Cl_6$, 97–100
GLYCOLATE
Studies on transition-metal peroxo complexes. Part 7. Molybdenum(VI) and tungsten(VI) carboxylato peroxo complexes, and the *X*-ray crystal structure of $K_2[MoO(O_2)_2(glyc)]\cdot 2H_2O$, 991–6
GOLD
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 51. Alkylidyne(carbaborane)tungsten-gold and -rhodium complexes; crystal structures of $[AuW(\mu\text{-}CR)(CO)_2(PPh_3)(\eta^5\text{-}C_2B_9H_9Me_2)]$, $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$, and $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2\{\eta^5\text{-}C_2B_9(C_7H_9)H_8Me_2\}]$ (R = C_6H_4Me-4), 61–72
Preparations, crystal structures, and molybdenum-95 nuclear magnetic resonance spectroscopic studies of triphenylphosphine–gold–tetrathiomolybdate(VI) clusters, 303–6
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au), 379–90
Spectrochemistry of solutions. Part 19. Complexation of gold(I) by cyanide in liquid ammonia: an infrared and Raman spectroscopic study, 673–6
Synthesis and structure of μ-[*trans*-1,2-bis(diphenylphosphino)-ethylene]-bis[chlorogold(I)], 677–80
Bimetallic systems. Part 17. Synthesis of Group 6 metal(0)–copper(I), –silver(I), or –gold(I) complexes with two $Ph_2PCH_2PPh_2$ (dppm) bridging ligands: crystal structure of $[(OC)_2W(\mu\text{-}CO)(\mu\text{-}dppm)_2(\mu\text{-}Cl)Cu]\cdot CH_2Cl_2$, 769–76
Bimetallic systems. Part 18. Nickel(II)–mercury(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$ ligands, 777–80
The heteronuclear cluster chemistry of the Group 1B metals. Part 3. Synthesis, structures, and chemical properties of the trimetallic hexanuclear Group 1B metal cluster compounds $[MM'Ru_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]$ (M = Cu, M′ = Ag or Au; M = Ag, M′ = Au). *X*-Ray crystal structure of the complex $[AgCuRu_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]\cdot CH_2Cl_2$, 1001–8
Lewis-base adducts of Group 11 metal(I) compounds. Part 27. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance, far-infrared, and structural studies on the mononuclear 2:1 adducts of triphenylphosphine with copper(I) and gold(I) halides, 1089–98
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au) (1987, 379), 1283–4
Gold(III) complexes with nitriles containing aromatic nuclei: crystal and molecular structure of trichloro(*p*-toluonitrile)gold(III), 1329–32

GOLD (contd)
Preparation and reactivity of halide-bridged binuclear carbonyl manganese and rhenium anions, $[M_2(\mu\text{-}X)(CO)_{8-2n}\{(EtO)_2POP(OEt)_2\}_n]^-$ (X = halide; M = Mn, n = 1 or 2; M = Re, n = 1). *X*-Ray crystal structure of $[Mn_2Au(\mu\text{-}Br)(CO)_6(\mu\text{-tedip})(PPh_3)]$, 1551–6
The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of $[M_2Ru_3(\mu_3\text{-}S)(\mu\text{-}Ph_2PCH_2PPh_2)(CO)_9]$ (M = Cu or Au), 1967–76
Crystal structures and properties of $[Au(phen)\{(CN)_{0.92}Br_{0.08}\}_2]Br$ and $[Au(phen)(CN)\{(CN)_{0.82}Br_{0.18}\}]\cdot 0.5$*trans*-$[Au(CN)_2Br_2]\cdot 0.5Br$·phen (phen = 1,10-phenanthroline) obtained by disproportionation of five-co-ordinate bromodicyano(1,10-phenanthroline)gold(III). Two examples of secondary co-ordination and CN/Br disorder in square-planar gold(III) complexes, 2235–40
Gold-197 Mössbauer spectra of some gold–ruthenium cluster compounds, 2333–6
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 67. Reactions of the salts $[X][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ [X = $P(CH_2Ph)Ph_3$, R = C_6H_4Me-4; X = NEt_4, R = Ph] with bis(cyclo-octa-1,5-diene)platinum, 2661–8
Gold–boron chemistry. Part 1. Synthetic, structural, and spectroscopic studies on the compounds $[5,6\text{-}\mu\text{-}(AuPR_3)\text{-}nido\text{-}B_{10}H_{13}]$ (R = cyclo-C_6H_{11} or C_6H_4Me-2), 2761–8
Reactivity of a cluster-co-ordinated acyl. Reduction of a μ-acetyl ligand to acetaldehyde, 2839–42
σ-Alkynyl complexes of manganese(I) as η^2-bonding ligands for Group 1B metal–ligand fragments. *X*-Ray crystal structure of $[Mn_2Cu(\mu\text{-}CCBu^t)_2(CO)_6(dppe)_2]PF_6\cdot 0.5CH_2Cl_2$, 2867–74
Mono- and bi-nuclear gold(I) and gold(III) complexes with $S_2C\text{–}PR_3$ ligands. *X*-Ray crystal structures of $[Au(C_6F_5)_3(S_2C\text{–}PEt_3)]$ and $[Au_2(\mu\text{-}S_2CPEt_3)(C_6F_5)_6]\cdot CH_2Cl_2$, 3017–22
Syntheses and crystal structures of complexes $[M_2R_2]$, [M = Cu, Ag, or Au; R = 2-$C(SiMe_3)_2C_5H_4N$] and $[Cu_4R'_4]$[R′ = 2-$CH(SiMe_3)C_5H_4N$]; electrochemical generation of $[Cu_2R_2]^{2+}$, 3085–92
GROUP 6
Bimetallic systems. Part 17. Synthesis of Group 6 metal(0)–copper(I), –silver(I), or –gold(I) complexes with two $Ph_2PCH_2PPh_2$ (dppm) bridging ligands: crystal structure of $[(OC)_2W(\mu\text{-}CO)(\mu\text{-dppm})_2(\mu\text{-}Cl)Cu]\cdot CH_2Cl_2$, 769–76
GROUP THEORY
A group theoretical paradigm for describing the skeletal molecular orbitals of cluster compounds. Part 2. Bispherical clusters, 1445–56
HAEMOGLOBIN
Reverse Bohr effect on the oxygen-binding affinity of heme embedded in a bilayer of liposome as a hemoglobin model: pH-induced oxygen uptake and evolution by aqueous synthetic lipid-heme solution, 1917–20
HAFNIUM
Crystal and molecular structures of $[MCl(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf) and $[TiCl_{0.5}Br_{0.5}(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$: an unusual π-interaction with the metals and molecular orbital calculations on model compounds, 1641–6
Axially asymmetric metal alkyls. Part 5. Synthesis and reduction to Zr^{III} species of Group 4 metallepines $[ML(\eta\text{-}C_5H_5)_2]$[L = $(2\text{-}CHRC_6H_4)_2^{2-}$, R = H, M = Ti, Zr, or Hf), isomers of $[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), and *meso*-[M{1,8-$(CHSiMe_3)_2C_{10}H_6$}$(\eta\text{-}C_5H_5)_2$] (M = Zr or Hf); *X*-ray crystal structures of $[TiL(\eta\text{-}C_5H_5)_2]$ (R = H) and λ-and δ-*RS*-$[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), 2347–58
HALIDE
Secondary bonding. Part 14. Structural isomerism in diaryliodonium halides and the structure of di(*p*-tolyl)iodonium bromide, 193–6
Tin-119 nuclear magnetic resonance studies of some pseudohalogeno-derivatives of $[SnX_6]^{2-}$ (X = Cl or Br), 315–8
Ferromagnetic tetrahalogenochromates(II) containing mixed halides (Cl and Br), 965–8
Reaction of silicon difluoride with halogens: a reinvestigation, 1123–6
Metallocene derivatives of early transition elements. Part 4. Synthesis and crystal structures of a series of zirconocene(IV) halides $[Zr(\eta\text{-}C_5H_4SiMe_3)_2X_2]$ (X = Cl or Br) and $[Zr\{\eta\text{-}C_5H_3(SiMe_3)_2\text{-}1,3\}_2X_2]$ (X = F, Br, or I), 1463–72
Preparation and reactivity of halide-bridged binuclear carbonyl manganese and rhenium anions, $[M_2(\mu\text{-}X)(CO)_{8-2n}\{(EtO)_2POP(OEt)_2\}_n]^-$ (X = halide; M = Mn, n = 1 or 2; M = Re, n = 1). *X*-Ray crystal structure of $[Mn_2Au(\mu\text{-}Br)(CO)_6(\mu\text{-tedip})(PPh_3)]$, 1551–6
Comments on the attempts by Wickens and Abrams to synthesize manganese(II) phosphine complexes: a reply, 1581–2
An electron spin resonance study of the equilibrium between tetrahalogeno-and pentahalogeno-nitridotechnetate(VI) ions in solution, 1721–6
Charge density in the $CoCl_4^{2-}$ ion: a comparison with spin density and theoretical calculations, 1737–46
Co-ordination chemistry of higher oxidation states. Part 23. Synthesis and properties of tetrahalogenoiridium(IV) complexes, $[IrL_2X_4]$ [X = Cl or Br; L = pyridine, PR_3, AsR_3, SbR_3, SR_2, or SeR_2]. Crystal and molecular structure of *trans*-$[Ir(AsEt_3)_2Br_4]$, 1901–10
Substitution reactions and luminescence spectra of the mixed complexes $[ReCl_xBr_{6-x}]^{2-}$, 1929–32
Structure and solvation of mercury(II) iodide and bromide in tetrahydrothiophene solution, 2411–6
HETEROPOLYANION
Triaquahenicosatungstodiphosphate(6−) heteropolyanion, $[P_2W_{21}O_{71}(OH_2)_3]^{6-}$: *X*-ray crystallographic and ^{183}W nuclear magnetic resonance structural studies (1986, 2237), 1283–4
HEXA-ALKYLBENZENE
The triphenylphosphine cone angle and restricted rotation about the chromium–phosphorus bond in dicarbonyl(η^6-hexa-alkylbenzene)(triphenylphosphine)chromium(0) complexes. Crystal and molecular structure of dicarbonyl(η^6-hexa-n-propylbenzene)(triphenylphosphine)chromium(0), 1545–50
HEXAFLUOROBUTYNE
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η^2-vinyl complexes of molybdenum and tungsten, $[M\{\eta^3\text{-}C(CF_3)C(CF_3)(L\text{–}L)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (L–L = $MeCO_2^-$, $Me_2PS_2^-$, $Et_2NCS_2^-$, or $Me_2NCS_2^-$), 2715–22
HISTIDINE
A thermodynamic and spectroscopic study of the proton and copper(II) complexes of L-prolyl-L-histidine, D-prolyl-L-histidine, L-histidyl-L-histidine, and D-histidyl-L-histidine, 661–6
HYDRATE
Concentrated aqueous solutions and liquid hydrates of zinc chloride: a study of their solvent behaviour using intershell, charge-transfer, and *d*–*d* adsorption spectra, 1485–8
HYDRATION
Studies on alkaline-earth metal sulphites. Part 9. Solid solutions of $Ca_3(SO_4)_2SO_3\cdot 12H_2O$ in $Ca_3(SO_3)_2SO_4\cdot 12H_2O$: structure, stability, and dehydration products, 235–40
HYDRAZIDE
Mechanism of the reaction between *trans*-$[W(NNH_2)(p\text{-}MeC_6H_4SO_3)(Ph_2PCH_2CH_2PPh_2)_2]^+$ and NEt_3: the influence of the metal on the rates of proton transfer of hydrazido(2−)-ligands, 197–200
The chemistry of the hydrazido(1−)-ligand. Preparations and crystal structures of $[Mo(NHNHCO_2Me)(NNCO_2Me)(S_2CNMe_2)_2]$, $[Mo(NHNMePh)(NNMePh)(S_2CNMe_2)_2]BPh_4$, and $[ReCl_2(NHNHCOPh)\text{-}(NNHCOPh)(PPh_3)_2]$. Mechanism of formation of substituted hydrazines from $[Mo(NNRPh)_2(S_2CNMe_2)_2]$ (R = Me or Ph), 529–40
HYDRAZINE
The chemistry of the hydrazido(1−)-ligand. Preparations and crystal structures of $[Mo(NHNHCO_2Me)(NNCO_2Me)(S_2CNMe_2)_2]$, $[Mo(NHNMePh)(NNMePh)(S_2CNMe_2)_2]BPh_4$, and $[ReCl_2(NHNHCOPh)\text{-}(NNHCOPh)(PPh_3)_2]$. Mechanism of formation of substituted hydrazines from $[Mo(NNRPh)_2(S_2CNMe_2)_2]$ (R = Me or Ph), 529–40
HYDRAZONE
Investigation into aroylhydrazones as chelating agents. Part 8. Synthesis and spectroscopic characterization of complexes of Co, Ni, Cu, Zn, and Cd with 2,6-diacetylpyridine bis(salicyloylhydrazone); *X*-ray crystal structure of dichloro[2,6-diacetylpyridine bis(salicyloylhydrazone)]cadmium(II)–chloroform–methanol (1/1/1), 177–82

HYDRAZONE (contd)
Reactions of planar pentadentate ligands with palladium and platinum chlorides. Crystal structures of *cis*-dichlorobis(*N'*-isopropylidenebenzohydrazide)platinum(II) and the cyclopalladated dianion of *N',N'''*-(pyridine-2,6-diyldiethylidyne)bis(benzohydrazide), 1175–80
An electron spin resonance study of the dioxygen adduct of bis[2-(2'-pyridylmethylenehydrazonomethyl)phenolato]cobalt(II) and related materials, 2341–6
HYDRIDE
Synthesis of $[Mo(\eta\text{-}C_5H_4Pr^i)(PR_3)_2H_3]$($R_3$ = Me_3 or Me_2Ph), $[Mo(\eta\text{-}C_5H_4Pr^i)(Pr^i_2PCH_2CH_2PPr^i_2)H_3]$, and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of $[\{Mo(\eta\text{-}C_5H_4Pr^i)(\mu\text{-}Cl)_2\}_2]$, 9–20
High-yield syntheses of sulphido triruthenium carbonyl cluster complexes containing bis(diphenylphosphino)methane, 91–6
Reactivity of ruthenium(II) complexes with 1-hydroxymethyl- or 1-benzyl-3,5-dimethylpyrazole and similar functionalized bipyrazoles. *X*-Ray crystal structure of carbonylchlorohydrido(3,5-dimethylpyrazole-N^2)bis(triphenylphosphine)ruthenium(II), 183–6
Bimetallic systems. Part 16. Formation of Group 6 metal(0)–platinum(II) hydride complexes, by ring opening of *fac*- or *mer*-$[M(CO)_3(dppm\text{-}PP')(dppm\text{-}P)]$ (M = Cr, Mo, or W; dppm = $Ph_2PCH_2PPh_2$). Reactions of $[(CO)_3Mo(\mu\text{-}dppm)_2PtH(Cl)]$, 221–6
Synthesis of the alkoxo(hydrido)-clusters $[Ru_3(\mu\text{-}H)(CO)_{10}(\mu\text{-}OR)]$ [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4
Synthesis and *X*-ray crystal structure of a cuboctahedral platinum cluster, 259–62
Reactions of metal carbonyl cluster complexes with multidentate phosphine ligands; the preparation and thermolysis of bis(dimethylphosphino)methane derivatives of $[Ru_3(CO)_{12}]$, and the *X*-ray crystal structure of $[Ru_3(CO)_9(H)(Me_2PCHPMe_2)]$, 277–80
Reaction of $[Os_5H_2(CO)_{15}]$ with pyridine: crystal and molecular structures of $[Os_5H_3(CO)_{14}(C_5H_4N)]$ and $[Os_5H_2(CO)_{14}(C_5H_5N)]$, 327–34
Nucleophilic attack upon ethylene co-ordinated to mercury(II): a molecular-orbital study concerning the origin of the acceleration by Hg^{II}, 361–8
Activation of alkanes involving rhenium atoms: synthesis and electronic structure of binuclear μ-alkylidene derivatives; molecular structure of $[\{Re(\eta\text{-}C_6H_6)\}_2(\mu\text{-}CHBu^t)(\mu\text{-}H)_2]$, 391–402
Synthesis of η-arene rhenium compounds using rhenium atoms. Crystal structures of $[Re_2(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)_2(\mu\text{-}H)_2]$ and $[Re(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)H]$, 403–10
Triosmium clusters containing ligands derived from tropone (cycloheptatrienone): molecular structure of an oxidative addition product $[Os_3H(\mu_3\text{-}C_7H_5O)(CO)_9]$, 551–6
Binuclear rhodium and iridium complexes containing pentamethylcyclopentadienyl and pyrazolate ligands, 639–46
Vibrational studies of interstitial hydrogen in metal carbonyl clusters, 685–90
Reactivity of the polyhydride complexes $[ReH_5(PPh_3)_3]$, $[ReH_3(dppe)_2]$ (dppe = $Ph_2PCH_2CH_2PPh_2$), $[ReH_3(PPh_3)_3L]$, and $[ReH_4(PPh_3)_3L]PF_6$ (L = MeCN or Bu^tNC) towards electrophiles and nucleophiles, 715–20
Reactions of aluminopolyhydride complexes of tungsten. *X*-Ray crystal structures of $[(Me_3P)_3H_3W(\mu\text{-}H)_2AlCl(\mu\text{-}N{=}CHEt)]_2$, $\{[(Me_3P)_3WH_5]Li\}_4$, $WH_2I(PMe_3)_3(SiMe_3)$, and $(Me_3P)_3H_2W(\mu\text{-}H)_3AlCl_2(NMe_3)$, 837–46
Trinuclear osmium clusters, containing bidentate $Ph_2PCH_2PPh_2$ and $Me_2PCH_2PMe_2$ ligands: *X*-ray crystal structures of the complexes $[Os_3H(CO)_9(Me_2PCHPMe_2)]$ and $[Os_3H(OH)(CO)_8(Ph_2PCH_2PPh_2)]$, 931–8
Complexes of the platinum metals. Part 32. Synthesis and chemistry of some iridium sulphonate derivatives, 947–52
Extended Hückel molecular orbital study of the effects of edge-bridging hydrogen atoms on the lengths of boron–boron and metal–metal bonds in cluster compounds, and the crystal structure of benzyltrimethylammonium octahydrotriborate(1−), 1017–26
High nuclearity osmium cluster radicals, 1051–4
The preparation, structure, and reactions of the mononuclear thiolate–hydride complexes $[MoH(SR)(dppe)_2]$ (R = bulkyl alkyl or aryl group, dppe = $Ph_2PCH_2CH_2PPh_2$); *X*-ray structure determination of *cis*-$[MoH(SC_6H_2Pr^i_3\text{-}2,4,6)(dppe)_2]$, 1115–22
The preparation, characterisation, and some reactions of $[Os_3H(X)(CO)_9(NMe_3)]$ (X = Cl, Br, or I), 1285–8
Systematic cluster build up by the reaction of $[Os_3H(X)(CO)_9(MeCN)]$ (X = SR or NHMe; R = Me, Et, or Ph) with $[OsH_2(CO)_4]$, 1289–92
Substitution reactions on triosmium clusters of the type $[Os_3(\mu\text{-}H)(\mu\text{-}X)(CO)_{10}]$ [X = NHMe, O_2CR, C_5H_4N, or OC(NHMe); R = H, Me, or Ph], 1293–8
Triosmium clusters containing the ligands CNH_2, NCHCN, and NCMeCN derived from cyanide ion, cyanogen, and acetonitrile, 1457–62
Intermediates in the conversion of $[Os(CO)_{11}(PRPh_2)]$ (R = Me or Ph) into $[Os_3(\mu_3\text{-}C_6H_4)(\mu_3\text{-}PR)(CO)_9]$: crystal and molecular structures of $[Os_3(\mu\text{-}H)(\mu_3\text{-}C_6H_4PMePh\text{-}o)(CO)_9]$ and $[Os_3(\mu_3\text{-}C_6H_4PMe\text{-}o)(CO)_{10}]$, 1529–34
Synthesis and substitution reactions of a heterodimetallic molybdenum–manganese complex, $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_6]$; *X*-ray crystal structure of $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_4(dppm\text{-}PP')]$, 1557–64
Synthesis of (η-cyclopentadienyl)(trimethylphosphine)tungsten derivatives and related chemistry, 1611–8
Substituted osmium clusters containing labile ligands. The preparation, characterisation, and some reactions of $[Os_3H(OR)(CO)_9(MeCN)]$ R = H, Me, Et, or Ph) and the isomers of $[Os_3H(SR)(CO)_9(MeCN)]$ (R = Me, Et, or Ph); the crystal and molecular structure of $[Os_3H(SPh)(CO)_9(PEt_3)]$, 1623–30
Preparation and characterisation of dicarboxylatobis(η-cyclopentadienyl)tungsten(IV) complexes: a novel and convenient synthesis from $[W(\eta\text{-}C_5H_5)_2H_2]$ using molecular oxygen as a hydrogen acceptor, 1857–60
Decarboxylation of vinylene carbonate at triosmium clusters; *X*-ray crystal structure of the μ_3-formylmethylidene compound $[Os_3(CHCHO)(CO)_{10}]$, 1861–4
Three isomeric hydride compounds formed by protonation of decacarbonylbis(dimethylphenylphosphine)triosmium and the *X*-ray structures of two of these: 1,1- and 1,2-$[Os_3H(CO)_{10}(PMe_2Ph)_2][PF_6]$, 1869–76
Reactions of rhodium(V) hydrido–silyl complexes with butyl-lithium and with lithium triethylhydridoborate, 1963–6
Carbon-13 nuclear magnetic resonance study of the properties of the unsaturated anion $[Re_3(\mu\text{-}H)_4(CO)_9(NCMe)]^-$ at low temperature, 2083–6
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 66. Carbaboranetungsten– platinum complexes. Polyhedral rearrangements of a 12-vertex cage system; crystal structures of $[PtW(CO)_2(PEt_3)_2\{\eta^6\text{-}C_2B_9H_9\text{-}(CH_2C_6H_4Me\text{-}4)Me_2\}]\cdot CH_2Cl_2$, $[PtW(\mu\text{-}H)\{\mu\text{-}\sigma{:}\eta^5\text{-}C_2B_9H_7(CH_2C_6H_4Me\text{-}4)Me_2\}(CO)_2(PMe_3)(PEt_3)_2]$, and related compounds, 2219–34
Synthesis, reactivity, and *X*-ray crystal structures of $[Os_6Pt_2H_2(CO)_{16}(\mu\text{-}C_8H_{10})(C_6H_{12})]\cdot CH_2Cl_2$ and $[Os_6H(CO)_{16}(\mu\text{-}C_8H_{11})]$. Model compounds for C–H activation, 2253–60
Synthesis using metal atoms of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[OsH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[MnH(\eta\text{-}C_6H_6)(PMe_3)_2]$, and $[Mn(\eta\text{-}C_5H_5)(PMe_3)_3]$: arene ligand-exchange reactions of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, 2823–30
Reactions of phosphine, arsine, and stibine with carbonylbis(triethylphosphine)iridium(I) halides. Part 1. Reactions in toluene; *X*-ray crystal structures of $[Ir(CO)ClH(PEt_3)_2(AsH_2)]$ and $[Ir(CO)XH(PEt_3)_2(\mu\text{-}ZH_2)RuCl_2(\eta^6\text{-}MeC_6H_4CHMe_2\text{-}p)]$ (X = Br, Z = P; X = Cl, Z = As), 2831–8
Reactivity of a cluster-co-ordinated acyl. Reduction of a μ-acetyl ligand to acetaldehyde, 2839–42
Hydride addition reactions of the olefin complexes $[Fe(\eta^5\text{-}C_5H_5)(CO)_2(olefin)][BF_4]$; formyl formation as the kinetically preferred process; regio- and stereo-selectivity during addition to the co-ordinated olefins, 2857–62
The mechanism of the photochemical production of hydrogen from aqueous solutions of hydridotris(triethylphosphine)platinum(II), 2941–6
Preparation, isomerization, and reactions of hydride complexes of ruthenium(II), 2989–94

HYDRIDE (contd)
Preparation of two mixed-metal carbonyl clusters based on the platinumtriosmium core. *X*-Ray crystal structure of $[Os_3PtH_2(cod)(CH_2)(CO)_9]$ (cod = cyclo-octa-1,5-diene), 3103–6
HYDRIDE ABSTRACTION
Transition-metal mediated asymmetric synthesis. Part 5. Alternative methods for the formation of tricarbonyl(η^5-cyclohexadienyl)-iron(1+) salts, 885–8
HYDRIDO
η-Benzenebis(trimethylphosphine)iron as a precursor to $[Fe(\eta\text{-}C_5R_5)(PMe_3)_2]$ (R = H or Me) derivatives: the equilibrium $[Fe](PMe_3)Et \rightleftharpoons [Fe](\eta\text{-}C_2H_4)H + PMe_3$, where $[Fe] = Fe(\eta\text{-}C_5Me_5)(PMe_3)$, 411–6
HYDROFORMYLATION
Transformation of $[PPh_4][Fe_5RhC(CO)_{16}]$ under hydroformylation conditions: synthesis and crystal structure of $[PPh_4][Fe_3Rh_3C(CO)_{15}]$, 1983–8
HYDROGENATION
Water photolysis. Part 2. An investigation of the relative advantages of various components of the sensitiser–electron relay–metal colloid system for the photoproduction of hydrogen from water, and the use of these systems in the photohydrogenation of unsaturated organic substrates, 1299–304
Preparation, isomerization, and reactions of hydride complexes of ruthenium(II), 2989–94
HYDROGEN BONDING
Crystal structure and mass spectrometry of dichlorodimethylbis[2(1*H*)-pyridinethione-*S*]tin(IV), 815–8
The chemistry of vitamin B_{12}. Part 28. Crystal structure of dicyanocobyrinic acid heptamethyl ester and its interaction with alcohols: the effects of hydrogen bonding to co-ordinated cyanide, 1349–58
Aqua(2,2′-bipyridyl)difluorocopper(II) dihydrate: *X*-ray structure reveals short hydrogen bonds and other unusual features, 2397–400
HYDROGEN HALIDE
Hydrogen halide radical anions, 421–6
HYDROLYSIS
Aluminium speciation studies in biological fluids. A new investigation of aluminium hydroxide equilibria under physiological conditions, 1145–8
The base catalysed hydrolysis of *trans*-dichloro-bis(1,2-diaminoethane)cobalt(III) and -(1,3-diaminopropane)cobalt(III) cations: rate-limiting proton transfer, 1775–8
Complex formation of beryllium(II) with salicylate and hydroxide ions in 1 mol dm^{-3} sodium perchlorate aqueous solution at 25 °C, 1853–6
The hydrolysis of metal ions. Part 10. Kinetic and equilibrium measurements of molybdenum(VI), 2149–58
Oxygen-17 and aluminium-27 nuclear magnetic resonance spectroscopic investigations of aluminium(III) hydrolysis products, 2317–22
Beryllium(II) hydrolysis in 3.0 mol dm^{-3} perchlorate, 2431–8
HYDROSILYLATION
The formation and characterisation of (η^2-ethene)hydrido(η^5-pentamethylcyclopentadienyl)(trisubstituted-silyl)rhodium complexes; intermediates in catalytic dehydrogenative silylation reactions, 2709–14
HYDROXIDE
Binuclear rhodium and iridium complexes containing pentamethylcyclopentadienyl and pyrazolate ligands, 639–46
Trinuclear osmium clusters, containing bidentate $Ph_2PCH_2PPh_2$ and $Me_2PCH_2PMe_2$ ligands: *X*-ray crystal structures of the complexes $[Os_3H(CO)_9(Me_2PCHPMe_2)]$ and $[Os_3H(OH)(CO)_8(Ph_2PCH_2PPh_2)]$, 931–8
Aluminium speciation studies in biological fluids. A new investigation of aluminium hydroxide equilibria under physiological conditions, 1145–8
HYPOHALOGENITE
Stability constants for hypochloritoborate and hypobromitoborate complex ions in aqueous solution, 943–6
IMIDAZOLE
Metal–ligand bonding parameters and magnetic properties of some previously reported tetragonal nickel(II) complexes, 825–30
Crystal structures of low- and high-spin iron(III) complexes with quadridentate Schiff bases, 1157–62
IMIDAZOLEDICARBOXYLATE
Dinuclear rhodium and iridium complexes of dicarboxyimidazolates; crystal structure of $[NBu_4][(cod)Rh(dci)Rh(cod)]\cdot 2Pr^iOH$, 3003–8
IMIDAZOLEPROPANAMIDE
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution, 435–42
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), and hydrogen ions in aqueous solution (1987, 435), 1583–4
IMIDE
Preparation of the nitrenes $ReF_5(NF)$, $ReF_5(NCl)$, and $ReF_5(NBr)$. *X*-Ray crystal structures of $ReF_5(NF)$ and $ReF_5(NCl)$, 567–72
Bis(organoimido)-complexes of tungsten(VI): the crystal and molecular structures of tetrachlorobis(t-butylamine)bis(t-butylimido)bis(μ-*p*-tolylimido)-ditungsten(VI) and tetrachlorobis(μ-phenylimido)-bis(t-butylamine)bis(t-butylimido)ditungsten(VI), 2059–66
Preparation and characterisation of mononuclear organoimido complexes of tungsten: *X*-ray crystal structures of $[WCl_4(NC_6H_4Me\text{-}p)(thf)]$, $[P(CH_2Ph)Ph_3][WCl_5(NC_6H_4Me\text{-}p)]$, and $[WCl_2(NPh)_2(bipy)]$, 2067–76
IMINOMETHYL
Triosmium clusters containing the ligands CNH_2, NCHCN, and NCMeCN derived from cyanide ion, cyanogen, and acetonitrile, 1457–62
INDENYL
Preparation and reactions of bis(isocyanide)(η^5-indenyl)rhodium(I) complexes, 851–2
INDIUM
The chemistry of trichloro[tris(trimethylsilyl)methyl] and trichloro[tris(dimethylphenylsilyl)methyl] complexes of gallium, indium, and thallium. Crystal and molecular structures of $[Li(thf)_2(\mu\text{-}Cl)_2GaCl\{C(SiMe_2Ph)_3\}]\cdot thf$, $[Li(thf)_3(\mu\text{-}Cl)InCl_2\{C(SiMe_3)_3\}]$, and $[\{(Me_3Si)_3C\}In(\mu\text{-}Cl)_2\{\mu\text{-}Fe(CO)_4\}In\{C(SiMe_3)_3\}]$ (thf = tetrahydrofuran), 747–56
Methylindium dialkylamido compounds: mono- and bis-dialkylamides involving four- and five-co-ordinated indium(III). *X*-Ray crystal structures of $\{Me_2In[\overline{N(CH_2)_2N(Me)CH_2CH_2}]\}_2$, $MeIn[\overline{MeNC(CH)_4N}]_2$, and $MeIn\{[MeN(CH_2)_2NMe]\text{-}InMe_2\}_2$, 2159–64
INSERTION
Solvent effect in the insertion reaction of iron(II) carbonyls, 519–22
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\overline{\{ON(Ph)C(NPh)S\}}(PPh_3)_2]$ and $[Pt\overline{\{ON(Ph)C(O)NPh\}}(PPh_3)_2]$, 605–8
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\overline{\{ON(Ph)C(NPh)S\}}(PPh_3)_2]$ and $[Pt\overline{\{ON(Ph)C(O)NPh\}}(PPh_3)_2]$ (1987, 605), 1583–4
Insertion of 2,6-dimethylphenyl isocyanide into the U–N bond of $[U(\eta\text{-}C_5H_5)_3(NEt_2)]$. Crystal structure of $[U(\eta\text{-}C_5H_5)_3\{C(NEt_2)=NC_6H_3Me_2\text{-}2,6\}]$, 2039–44
INTERCALATION
Intercalation compounds of (diphosphato)oxotungsten(VI), $[WO(P_2O_7)]$, 609–12
INTERMEDIATE
The formation and characterisation of (η^2-ethene)hydrido(η^5-pentamethylcyclopentadienyl)(trisubstituted-silyl)rhodium complexes; intermediates in catalytic dehydrogenative silylation reactions, 2709–14
IODIDE
Determination of the deuterium position in DNb_6I_{11}, 431–4
IODINE
Secondary bonding. Part 14. Structural isomerism in diaryliodonium halides and the structure of di(*p*-tolyl)iodonium bromide, 193–6
IODONIUM
Secondary bonding. Part 14. Structural isomerism in diaryliodonium halides and the structure of di(*p*-tolyl)iodonium bromide, 193–6
ION PAIR
Chiral recognition in ion pairing of an optically active

ION PAIR (contd)
tris(ethylenediamine)cobalt(III) cation and application to chromatographic resolution of metal complex anions, 1977–82
I.R.
Matrix isolation studies of monomeric boron trisulphide, 267–70
Spectrochemistry of solutions. Part 19. Complexation of gold(I) by cyanide in liquid ammonia: an infrared and Raman spectroscopic study, 673–6
Synthesis and structure of μ-[*trans*-1,2-bis(diphenylphosphino)-ethylene]-bis[chlorogold(I)], 677–80
The vibrational spectrum of tricarbonyl(trichlorostibine)nickel(0), $[Ni(CO)_3(SbCl_3)]$, 681–2
Vibrational studies of interstitial hydrogen in metal carbonyl clusters, 685–90
The configuration and lattice dynamics of complexes of dialkyltin(IV) with adenosine 5′-monophosphate and phenyl phosphates, 789–94
Matrix isolation and mass spectrometric studies on the vaporisation of chromium(III) chloride. The characterisation of molecular $CrCl_4$ and $CrCl_3$, 859–66
Photoactivation of methane by η^5-cyclopentadienyl and substituted η^5-cyclopentadienyl Group 8 metal dicarbonyl complexes, $[M(\eta^5\text{-}C_5R_5)(CO)_2]$ (M = Rh or Ir, R = H or Me), and dicarbonyl(η^5-indenyl)iridium: a matrix isolation study, 1181–90
A quantitative approach to host–guest interactions for matrix-isolated alkali-metal salts of hexafluoroanions and perchlorates, 1521–8
Infrared spectroscopic studies of uranyl(VI) species adsorbed from aqueous $[UO_2(CO_3)_3]^{4-}$ solutions on to a polymer bearing amidoxime groups, 2489–92
Surface characterisation of α-, β-, γ-, and δ-UO_3 using *X*-ray photoelectron spectroscopy, 3009–16
Preparations, properties, and vibrational spectra of some (dimethylamino)halogenosilanes, 3029–34
IRIDIUM
Binuclear rhodium and iridium complexes containing pentamethylcyclopentadienyl and pyrazolate ligands, 639–46
Complexes of the platinum metals. Part 32. Synthesis and chemistry of some iridium sulphonate derivatives, 947–52
Photoactivation of methane by η^5-cyclopentadienyl and substituted η^5-cyclopentadienyl Group 8 metal dicarbonyl complexes, $[M(\eta^5\text{-}C_5R_5)(CO)_2]$ (M = Rh or Ir, R = H or Me), and dicarbonyl(η^5-indenyl)iridium: a matrix isolation study, 1181–90
Kinetic studies on the oxidation of ethylenediaminebis(pyridine)- and bis(ethylenediamine)-platinum(II) complexes by hexachloroiridate(IV), 1473–6
Bimetallic systems. Part 20. Synthesis of Group 6 metal(0)–iridium(I) or –rhodium(I) complexes bridged by $Ph_2PCH_2PPh_2$, 1495–500
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$], 1805–12
Co-ordination chemistry of higher oxidation states. Part 23. Synthesis and properties of tetrahalogenoiridium(IV) complexes, $[IrL_2X_4]$ [X = Cl or Br; L = pyridine, PR_3, AsR_3, SbR_3, SR_2, or SeR_2]. Crystal and molecular structure of *trans*-$[Ir(AsEt_3)_2Br_4]$, 1901–10
Activation energies for molecular tumbling and cyclopentadienyl rotation in $[M(\eta^5\text{-}C_5H_5)(\eta^4\text{-cod})]$ (M = Rh or Ir; cod = cycloocta-1,5-diene) and the *X*-ray crystal structure of and bonding in $[Rh(\eta^5\text{-}C_5H_5)(\eta^4\text{-cod})]$, 1947–52
Some chemistry of the heterobimetallic complexes $[(OC)_3Mo(\mu\text{-dppm})_2Rh(CO)Cl]$ and $[(OC)_3Mo(\mu\text{-dppm})_2\text{-}M(CO)_2]PF_6$ (M = Rh or Ir, dppm = $Ph_2PCH_2PPh_2$). Structure of $[(OC)_3Mo(\mu\text{-dppm})_2(\mu\text{-}C{\equiv}CMe)Rh(CO)]$, 2171–6
Attempted synthesis of osmium(VI) and iridium(VI) thiofluorides; the preparation of $OsF_5{\cdot}SF_4$ and $IrF_5{\cdot}SF_4$, 2285–8
The cyclometallation of benzoic acid to give rhodium, iridium, and osmium *C,O*-benzoates. *X*-Ray structure determination of the dibenzoate $[(C_5Me_5)Rh(OOCPh)_2(H_2O)]$, 2459–66
Preparation and spectroscopic study of functionally substituted cyclopentadienides of thallium(I), potassium, rhodium(I), and iridium(I), 2615–20
Reduction–oxidation properties of organotransition-metal complexes. Part 27. Mixed-valence triazenido-bridged di-iridium compounds, 2737–40
Reactions of phosphine, arsine, and stibine with carbonylbis(triethylphosphine)iridium(I) halides. Part 1. Reactions in toluene; *X*-ray crystal structures of $[Ir(CO)ClH(PEt_3)_2(AsH_2)]$ and $[Ir(CO)XH(PEt_3)_2(\mu\text{-}ZH_2)RuCl_2(\eta^6\text{-}MeC_6H_4CHMe_2\text{-}p)]$ (X = Br, Z = P; X = Cl, Z = As), 2831–8
Dinuclear rhodium and iridium complexes of dicarboxyimidazolates; crystal structure of $[NBu_4][(cod)Rh(dci)Rh(cod)]{\cdot}2Pr^iOH$, 3003–8
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$] (1987, 1805), 3107–8
IRON
Magnetic and electrochemical studies on the series $[Cr_nFe_{3-n}(\mu_3\text{-}O)(gly)_6(H_2O)_3][NO_3]_7{\cdot}xH_2O$(gly = glycine; n = 0, 1, 2, or 3) and on the mixed-valence complex $[Fe^{II}Fe^{III}{}_2(\mu_3\text{-}O)(gly)_6\text{-}(H_2O)_3]Cl_6$, 97–100
Preparation and magnetism of the binuclear iron(II) complexes $[\{Fe(C_9H_{21}N_3)X_2\}_2]$ (X = NCS, NCO, or N_3) and their reaction with NO. Crystal structures of $[\{Fe(C_9H_{21}N_3)(NCS)_2\}_2]$ and $[Fe(C_9H_{21}N_3)(NO)(N_3)_2]$, 187–92
Synthesis of the alkoxo(hydrido)-clusters $[Ru_3(\mu\text{-}H)(CO)_{10}(\mu\text{-}OR)]$ [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4
Equilibria in aqueous solutions of some iron(II) complexes, 255–8
Equilibrium and kinetic study of nitric oxide binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 369–72
η-Benzenebis(trimethylphosphine)iron as a precursor to $[Fe(\eta\text{-}C_5R_5)(PMe_3)_2]$ (R = H or Me) derivatives: the equilibrium $[Fe](PMe_3)Et \rightleftharpoons [Fe](\eta\text{-}C_2H_4)H + PMe_3$, where $[Fe] = Fe(\eta\text{-}C_5Me_5)(PMe_3)$, 411–6
Formation of a methoxycarbonyl group by attack of methanol on an iron(II) carbonyl complex, assisted by intramolecular proton transfer: *X*-ray structure determination of $[Fe(CO)(PMe_3)_2(CNC_6H_{11})\{CO(OCH_3)\}\{C(CH_3){=}NHC_6H_{11}\}]\text{-}BPh_4$, 473–6
Solvent effect in the insertion reaction of iron(II) carbonyls, 519–22
Phosphorus-tellurium compounds. Part 3. Cyclopentadienyliron tellurophosphorane complexes: tellurium-bridged iron phosphine compounds?, 541–4
Mössbauer spectroscopic studies on molecular rotation and lattice dynamics of cyclodextrin clathrates of bis(η-cyclopentadienyl)-iron(II) and some of its ring-substituted derivatives, 627–32
A comparative synthetic and structural study of μ-oxo and μ-sulphido derivatives of *N,N′*-ethylenebis(acetylacetoneiminato)-iron(III), including the synthesis and structural characterization of the parent compound, 709–14
The chemistry of trichloro[tris(trimethylsilyl)methyl] and trichloro[tris(dimethylphenylsilyl)methyl] complexes of gallium, indium, and thallium. Crystal and molecular structures of $[Li(thf)_2(\mu\text{-}Cl)_2GaCl\{C(SiMe_2Ph)_3\}]{\cdot}thf$, $[Li(thf)_3(\mu\text{-}Cl)InCl_2\{C(SiMe_3)_3\}]$, and $[\{(Me_3Si)_3C\}In(\mu\text{-}Cl)_2\{\mu\text{-}Fe(CO)_4\}In\{C(SiMe_3)_3\}]$ (thf = tetrahydrofuran), 747–56
Redox behaviour of the iron–sulphur cluster $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{2+}$. Synthesis and crystal structure of the new paramagnetic monopositive species $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{+}$ as its $[PF_6]^-$ salt, 831–6
Transition-metal mediated asymmetric synthesis. Part 5. Alternative methods for the formation of tricarbonyl(η^5-cyclohexadienyl)iron(1+) salts, 885–8
Chemistry of metallacyclic complexes containing the FeCSCS ring obtained by metal-promoted CS_2–CS coupling. *X*-Ray crystal structures of $[(CO)(cp)FeC\{SW(CO)_5\}SC(FeL_n)S]$ and $[(CO)(cp)FeC(SEt)SC(FeL_n)S]SO_3CF_3$ [FeL_n = Fe(cp)$(CO)_2$, cp = η-C_5H_5], 1133–44
Crystal structures of low- and high-spin iron(III) complexes with quadridentate Schiff bases, 1157–62
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 54. Compounds derived from $[W(\equiv CC_6H_4Me\text{-}4)(CO)_2(\eta\text{-}C_5Me_5)]$ with bonds between tungsten and iron, cobalt, or nickel; crystal structure of $[FeW\{\mu\text{-}C(C_6H_4Me\text{-}4)C(O)O\}(CO)_5(\eta\text{-}C_5Me_5)]$, 1191–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 56. Synthesis of iron–molybdenum compounds; crystal structures of $[FeMo(\mu\text{-}CR)(CO)_6(\eta\text{-}C_5H_5)]$ and $[FeMo_2(\mu_3\text{-}RC_2R)(CO)_6(\eta\text{-}C_5H_5)_2]$ (R = C_6H_4Me-4), 1209–14

IRON (contd)
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 57. Reactions of iron– molybdenum complexes with oxygen and sulphur; crystal structure of $[FeMo(\mu\text{-}\eta^2\text{-}SCC_6H_4Me\text{-}4)(CO)_5(\eta\text{-}C_5H_5)]$, 1215–20
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 61. Reactions of iron– molybdenum compounds with tertiary phosphines, 1243–8
Carbon-13 and phosphorus-31 nuclear magnetic resonance studies of the fluxional behaviour of the complexes $[Fe_3(CO)_{10}\{P(OMe)_3\}_2]$ and $[Fe_3(CO)_{10}\{P(OMe)_3\}\{P(OPh)_3\}]$, 1249–52
New spin-crossover iron(III) complexes with large hysteresis effects and time dependence of their magnetism, 1341–8
Magnetic exchange interactions in perovskite solid solutions. Part 8. A study of the solid solution $SrFe_{1-x}Co_xO_{3-y}$ by iron-57 Mössbauer spectroscopy, 1419–24
Bimetallic systems. Part 19. Chromium(0)–, molybdenum(0)–, or tungsten(0)–iron(0) carbonyl complexes containing one or two bridging $Ph_2PCH_2PPh_2$ ligands: crystal structure of $[(OC)_3Mo(\mu\text{-}Ph_2PCH_2PPh_2)_2Fe(CO)_3]$, 1489–94
Bimetallic systems. Part 22. Molybdenum(0)– and manganese(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the reversible uptake of carbon monoxide: crystal structure of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)Mo(CO)_4]\cdot 0.5C_6H_5Cl$, 1509–14
Synthesis and characterization of new heterobimetallic iron–cobalt complexes with ethenyl bridges $CR^2{=}CHR^1$ [R^1 = H or C(O)OMe, R^2 = C(O)OMe or C(O)OEt]. *X*-Ray structure of $[FeCo(CO)_6\{\mu\text{-}MeOC(O)C{=}C(H)C(O)OMe\}]$, 1619–22
Iron antimonate, 1727–32
Reverse Bohr effect on the oxygen-binding affinity of heme embedded in a bilayer of liposome as a hemoglobin model: pH-induced oxygen uptake and evolution by aqueous synthetic lipid-heme solution, 1917–20
Electrophilic attack on a series of dinuclear diphosphazane-bridged derivatives of iron by halogens. *X*-Ray crystal structures of $[Fe_2I(CO)_5\{\mu\text{-}(MeO)_2PN(Et)P(OMe)_2\}_2]PF_6$ and $[Fe_2(\mu\text{-}Br)(CO)_4\{\mu\text{-}(PhO)_2PN(Et)P(OPh)_2\}_2]PF_6$, 1933–46
X-Ray structural study of [Fe(saltrien)]X (X = Br·2H_2O, BPh_4, or PF_6). Origin of unusual magnetic behaviour of the spin-crossover complex $[Fe(saltrien)]PF_6$, 1957–62
Transformation of $[PPh_4][Fe_3RhC(CO)_{16}]$ under hydroformylation conditions: synthesis and crystal structure of $[PPh_4][Fe_3Rh_3C(CO)_{15}]$, 1983–8
The synthesis and reactions of palladium–iron carbonyl complexes containing bridging $Ph_2PCH_2PPh_2$ ligands, 2005–8
Alkylation of tricarbonyliron complexes by trimethylsilyl cyanide: synthetic and kinetic studies, 2013–6
(η-Benzene)bis(trimethylphosphine)iron: a useful precursor to polyene iron derivatives. Crystal structure of 2—4-η:2′— 4′-η-1,1′-bi(cyclohex-3-en-2-yl)bis(trimethylphosphine)iron(II), 2031–8
Reactions of main group homopolynuclear cations with transition-metal carbonyls. Stabilisation of $\mu\text{-}Se_2^{2+}$ and $\eta^3\text{-}Te_3^{2+}$ dicationic ligands and crystal structures of $[Mo(Te_3)(CO)_4][SbF_6]_2$ and $[FeW(Se_2)(CO)_8][SbF_6]_2$, 2111–6
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 64. Addition of methylene groups to iron–molybdenum complexes; crystal structures of $[FeMo(\mu\text{-}CH_2)\{\mu\text{-}\sigma{:}\eta\text{-}C(C_6H_4Me\text{-}4){=}CH_2\}(CO)_5(\eta\text{-}C_5H_5)]$ and $[FeMo\{\mu\text{-}C(C_6H_4Me\text{-}4)C(OMe)C(H)\}(CO)_5(\eta\text{-}C_5H_5)]$, 2201–10
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 65. Reactions of the complexes $[FeW(\mu\text{-}CR)(CO)_5(\eta\text{-}C_5Me_5)]$ and $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5Me_5)]$ (R = C_6H_4Me-4) with alkynes $R'C_2R'$ (R′ = Me or Ph); crystal structures of $[FeW\{\mu\text{-}C(R)C(O)C(Me)C(Me)\}(CO)_5(\eta\text{-}C_5Me_5)]\cdot CH_2Cl_2$ and $[FeW\{\mu\text{-}C(R)C(Et)C(H)C(Me)C(Me)\}(\mu\text{-}CO)(CO)_3(\eta\text{-}C_5Me_5)]$, 2211–8
Iron–ligand bonding properties of synthetic iron–porphyrin complexes with oxygen transporting ability in aqueous media, 2455–8
Dynamics of oxygen binding to modified iron(II)–porphyrin complexes embedded in the bilayer of phospholipid liposome, 2493–6
A realistic model for heme-containing catalases and peroxidases: the *X*-ray structural characterisation of a non-porphyrin iron(III) macrocyclic complex, and the mechanism of its peroxidation of aromatic substrates, 2505–10
Kinetic studies on 1:1 electron-transfer reactions involving blue copper proteins. Part 14. Reactions of poplar plastocyanin with inorganic complexes, 2573–8
New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states, 2621–30
Rhodium(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the formation of a tetranuclear cluster from a heterobimetallic precursor: crystal structures of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)RhCl(CO)]$ and $[Fe_2Rh_2(\mu\text{-}Ph_2PCH_2PPh_2)_2(CO)_8]$, 2751–6
Hydride addition reactions of the olefin complexes $[Fe(\eta^5\text{-}C_5H_5)(CO)_2(olefin)][BF_4]$; formyl formation as the kinetically preferred process; regio- and stereo-selectivity during addition to the co-ordinated olefins, 2857–62
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 68. Reactions between nonacarbonyldi-iron and the salts $[NEt_4][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = Me, Ph, or C_6H_4Me-4); crystal structures of $[NEt_4][FeW\{\mu\text{-}CH(C_6H_4Me\text{-}4)\}(\mu\text{-}\sigma{:}\eta^5\text{-}C_2B_9H_8Me_2)(\mu\text{-}CO)(CO)_5]$ and $[NEt_4][FeW(\mu_3\text{-}CPh)(\mu\text{-}\sigma{:}\sigma',\eta^5\text{-}C_2B_9H_7Me_2)(CO)_8]$, 2917–26
Dienophile cycloaddition to cycloheptatriene and related complexes of tricarbonyliron. *X*-Ray crystal structures of $[Fe(CO)_3(\eta^4\text{-}C_7H_7CN)]\cdot C_2(CN)_4$ and $[Fe(CO)_3(\eta^4\text{-}C_7H_7\text{–}C_7H_7)]$, 2927–34
Oxygen- and carbon monoxide-binding to a lipophilic diporphyrinatocopperiron complex solubilized in an aqueous medium with a micelle, 2963–8
Platinum(II)– and palladium(II)–iron(0) complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand. Crystal structures of $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}CO)PtBr_2]$ and $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}I)PtI]\cdot CH_2Cl_2$, 3079–84
IRON-57
Magnetic exchange interactions in perovskite solid solutions. Part 8. A study of the solid solution $SrFe_{1-x}Co_xO_{3-y}$ by iron-57 Mössbauer spectroscopy, 1419–24
ISOCYANATE
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\{\overline{ON(Ph)C(NPh)S}\}(PPh_3)_2]$ and $[Pt\{\overline{ON(Ph)C(O)NPh}\}(PPh_3)_2]$, 605–8
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\{\overline{ON(Ph)C(NPh)S}\}(PPh_3)_2]$ and $[Pt\{\overline{ON(Ph)C(O)NPh}\}(PPh_3)_2]$ (1987, 605), 1583–4
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S{=}C(NHPh)\text{-}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O\text{–}C(NHPh){=}S\}]$, 2299–304
Reactivity of rhenium(I) pseudo-allyl complexes with hererocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S{=}C(NHPh)\text{-}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O\text{–}C(NHPh){=}S\}]$ (1987, 2299), 3107–8
ISOCYANIDE
Phenyl complexes of ruthenium(II): competition between carbonyl substitution and formation of benzoyl complexes, and an *X*-ray study of a benzoyl complex, 27–34
Preparation of isocyanide and mixed dinitrogen–isocyanide complexes of rhenium(I) from reactions of *trans*-$[ReCl(N_2)(PMe_2Ph)_4]$, *mer*-$[Re(S_2PPh_2)(N_2)(PMe_2Ph)_3]$, or *mer*-$[Re(S_2CNEt_2)(N_2)(PMe_2Ph)_3]$ with methyl isocyanide; crystal structure of *mer*-$[Re(S_2PPh_2)(N_2)(CNMe)(PMe_2Ph)_3]$, 319–26
Chemistry of platinum sulphido-complexes. Part 6. Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with the rhodium complexes $[(RhL_2Cl)_2]$ (L = CO or C_2H_4) and $[Rh(2,6\text{-}Me_2C_6H_3NC)_4]PF_6$. Crystal and molecular structures of $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(CO)_2]\cdot[Rh(CO)_2Cl_2]\cdot C_3H_6O$ and $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4\cdot(2,6\text{-}Me_2C_6H_3NC)_2]PF_6$, 335–40
The addition of isonitriles to $[\{M(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (M = Mo or W). Syntheses and properties of $[M_2(\mu\text{-}\eta^2\text{-}CNR)(CO)_4(\eta\text{-}C_5H_5)_2]$ (M = Mo or W, R = Me or Bu^t) and $[\{Mo(CNR)(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (R = Me or Bu^t). *X*-Ray crystal

ISOCYANIDE (contd)
structures of $[Mo_2(\mu\text{-}\eta^2\text{-}CNBu^t)(CO)_4(\eta\text{-}(\eta\text{-}C_5H_5)_2]$ and $[\{Mo(CNBu^t)(CO)_2(\eta\text{-}C_5H_5)\}_2]$, 341–8
Syntheses and reactions of acyclic and cyclic carbene complexes *cis*-$[MI(=CMeNHMe)(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo or W) and *cis*-$[MI\{=\overline{C(CH_2)_nNMe}\}(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo, n = 3 or 4; M = W, n = 3) obtained through alkyl migration to methyl isocyanide co-ordinated to molybdenum or tungsten. *X*-Ray structure of *cis*-$[MI\{=\overline{C(CH_2)_3NMe}\}(CO)_2(\eta\text{-}C_5H_5)]$ (1986, 2127), 479–80
Reactivity of the polyhydride complexes $[ReH_5(PPh_3)_3]$, $[ReH_3(dppe)_2]$ (dppe = $Ph_2PCH_2CH_2PPh_2$), $[ReH_3(PPh_3)_3L]$, and $[ReH_4(PPh_3)_3L]PF_6$ (L = MeCN or Bu^tNC) towards electrophiles and nucleophiles, 715–20
Preparation and reactions of bis(isocyanide)(η^5-indenyl)rhodium(I) complexes, 851–2
Reactions of cyclobutenedione and cyclopentenetrione with zero-valent nickel triad complexes of isocyanide. *X*-Ray crystal structures of (3—4-η-3,4-diphenylcyclobut-3-ene-1,2-dione)- and (4—5-η-4,5-diphenylcyclopent-4-ene-1,2,3-trione)-tris(t-butyl isocyanide)nickel, 853–8
Conversion of uncharged methyl complexes of ruthenium(II) into a cationic acyl complex: a mechanistic study, 1713–6
Cationic ruthenium(II) phosphite complexes. Preparation and properties of monocarbonyl and isocyanide derivatives, 1813–6
Electrochemical reactions of dichlorobis(aryl isocyanide)platinum(II) complexes, 1833–8
Insertion of 2,6-dimethylphenyl isocyanide into the U–N bond of $[U(\eta\text{-}C_5H_5)_3(NEt_2)]$. Crystal structure of $[U(\eta\text{-}C_5H_5)_3\{C(NEt_2)=NC_6H_3Me_2\text{-}2,6\}]$, 2039–44

ISOMERISATION
trans to *cis* isomerisations of platinum(II) olefin phosphine complexes, $[PtX_2(ol)(PR_3)]$ (X = Cl or Br; ol = ethene, propene, or 1-heptene; PR_3 = PBu_3 or PMe_2Ph), 139–14
Preparation, isomerization, and reactions of hydride complexes of ruthenium(II), 2989–94

ISOMERISM
Isomers in *fac*-$[ReBr(CO)_3(L_c)_2]$ (L_c = *cis*-3,4-dihydro-2,3,4,5-tetraphenyl-2*H*-1,2,3-diazaphosphole) arising from chirality of phosphorus ligands: separation and *X*-ray structural studies, 21–6
The first authenticated example of geometrical isomers in organotin(IV) co-ordination chemistry. The crystal and molecular structures of octahedral *cis*- and *trans*-SnR_2 isomeric adducts of dichlorobis(4-chlorophenyl)tin(IV) with 4,4′-dimethyl-2,2′-bipyridyl, 129–38

ISOTHIOCYANATE
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\{ON(Ph)C(NPh)S\}(PPh_3)_2]$ and $[Pt\{ON(Ph)C(O)NPh\}(PPh_3)_2]$, 605–8
Thermally induced phase transition and *X*-ray crystal structure of bis(*N*,*N*′-dimethyl-1,2-ethanediamine)di-isothiocyanatonickel(II), 997–1000
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\{ON(Ph)C(NPh)S\}(PPh_3)_2]$ and $[Pt\{ON(Ph)C(O)NPh\}(PPh_3)_2]$ (1987, 605), 1583–4
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S=C(NHPh)\text{-}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O\text{–}C(NHPh)=S\}]$, 2299–304
Reactivity of rhenium(I) pseudo-allyl complexes with hererocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S=C(NHPh)\text{-}S\}]\cdot[Re(CO)_2(PPh_3)_2\{O\text{–}C(NHPh)=S\}]$ (1987, 2299), 3107–8

ISOTHIOCYANIC ACID
Dicopper(II) complexes of a 28-membered N_8 macrocycle. Evidence for proton transfer from co-ordinated secondary amine to thiocyanate, leading to formation of an isothiocyanic acid complex, 873–80

KINETICS
Mechanism of the reaction between *trans*-$[W(NNH_2)(p\text{-}MeC_6H_4SO_3)(Ph_2PCH_2CH_2PPh_2)_2]^+$ and NEt_3: the influence of the metal on the rates of proton transfer of hydrazido(2−)-ligands, 197–200
Equilibrium and kinetic study of nitric oxide binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 369–72
A structural study of the complexation of the lithium ion by the cryptand 4,7,13-trioxa-1,10-diazabicyclo[8.5.5]icosane, 489–92
Kinetics and mechanism of ring closure for a series of photoproduced $[M(CO)_5L]$ intermediates (M = Cr, Mo, or W; L = pyridine 2-carbaldehyde imine ligand), 513–8
Solvent effect in the insertion reaction of iron(II) carbonyls, 519–22
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid, 573–8
Studies of phosphazenes. Part 31. Kinetic studies of the reactions of hexachlorocyclotriphosphazene with aromatic primary amines: evidence for the intermediacy of a reactive three-co-ordinated phosphorus(v) species, 585–90
Substitution reactions between penta-ammineaquaruthenium(II) and thiocyanato, protonated acetato, cyano, and oxalato ligands in aqueous acidic solutions, 957–60
A kinetic and mechanistic study of the formation of a diaryl ditelluride in solution by the sulphite reduction of the aryltellurium trichloride, 985–90
The relative nucleophilic efficiency of water and the azide and thiocyanate ions in the aqueous substitution of chloropentacyanocobaltate(III), 1059–62
Electronic *d*–*d* spectra and ligand-dissociation rates of pentacyanoruthenate(II) ions, 1267–70
Reaction paths and kinetics in the Cr^{2+} reduction of the Mo^V_2 and Mo^{IV}_3 ions $[Mo_2O_4(H_2O)_6]^{2+}$ and $[Mo_3O_4(H_2O)_9]^{4+}$, 1397–402
Reactions of chromium(III) complexes of 1,10-phenanthroline, 2,2′-bipyridyl, and oxalate with the pulse radiolytically generated aquated electron, zinc(I), and cadmium(I), 1425–30
Kinetic studies on the oxidation of ethylenediaminebis(pyridine)- and bis(ethylenediamine)-platinum(II) complexes by hexachloroiridate(IV), 1473–6
The base catalysed hydrolysis of *trans*-dichloro-bis(1,2-diaminoethane)cobalt(III) and -(1,3-diaminopropane)cobalt(III) cations: rate-limiting proton transfer, 1775–8
Kinetic study of the homogeneous methanolysis of tetraphosphorus decasulphide, 1897–900
Alkylation of tricarbonyliron complexes by trimethylsilyl cyanide: synthetic and kinetic studies, 2013–6
Synthesis, characterisation, and aquation kinetics of [*N*-(*o*-carboxyphenyl)iminodiacetato](glycinato)chromate(III) complex ion in acidic media, 2095–8
The hydrolysis of metal ions. Part 10. Kinetic and equilibrium measurements of molybdenum(VI), 2149–58
Chromium(III)-catalysed cerium(IV) oxidation of arsenic(III) in aqueous sulphuric acid (1987, 573), 2291–2
Preparation and properties of the tungsten(V) aqua dimer $[W_2O_4(H_2O)_6]^{2+}$: kinetic studies on the 1:1 reaction with NCS^-, 2293–8
The relative nucleophilic efficiency of water and dimethyl sulphoxide in the aqueous substitution of aquapentacyanocobaltate(III) and chloropentacyanocobaltate(III), 2311–6
Kinetics of reduction of tris(ethylenediamine)cobalt(III) by ethylenediamine-*NNN*′*N*′-tetra-acetatochromate(II), 2485–8
Dynamics of oxygen binding to modified iron(II)–porphyrin complexes embedded in the bilayer of phospholipid liposome, 2493–6
Kinetic studies on 1:1 electron-transfer reactions involving blue copper proteins. Part 14. Reactions of poplar plastocyanin with inorganic complexes, 2573–8
Synthetic and equilibrium studies on copper(II), nickel(II), and palladium(II) complexes of *N*,*N*′,*N*″,*N*‴-tetrakis(2-hydroxyethyl)-1,4,8,11-tetra-azacyclotetradecane (L); kinetics of the acid dissociation of the copper(II) complex and *X*-ray structural characterisation of the nickel(II) complex $[NiLH_{-1}][ClO_4]$, 2605–14
Ligand intra- and inter-molecular exchange processes in dioxopentakis(trimethylurea)uranium(VI). A proton nuclear magnetic resonance study, 2885–8

LANTHANIDE
Syntheses, *X*-ray structures, and properties of complexes of

LANTHANIDE (contd)
macrocyclic hexaimines with lanthanide nitrates, 1665–74
Syntheses and properties of complexes of a conjugated macrocyclic hexamine with lanthanide nitrates, 2289–90
Mixed-metal phosphinito complexes of platinum(II) and palladium(II) with lanthanide and actinide elements. The single-crystal *X*-ray structure determination of $[UO_2(OH_2)\{(OPPh_2)_2Pd(S_2CNEt_2)\}_2]$, 2853–6
LATTICE DYNAMICS
Mössbauer spectroscopic studies on molecular rotation and lattice dynamics of cyclodextrin clathrates of bis(η-cyclopentadienyl)-iron(II) and some of its ring-substituted derivatives, 627–32
The configuration and lattice dynamics of complexes of dialkyltin(IV) with adenosine 5′-monophosphate and phenyl phosphates, 789–94
LEAD
Binucleating N_6 24- and 26-membered macrocyclic ligands. Part 1. Dilead complexes: *X*-ray crystal structure determination of a macrocyclic dilead complex containing nitrogen-only bridging thiocyanate, 123–8
Complexes of ligands providing endogenous bridges. Part 1. The syntheses and crystal structures of barium and lead(II) complexes of macrocyclic Schiff bases derived from heterocyclic dicarbonyls and 1,*n*-diamino-*n*′-hydroxyalkanes (*n*,*n*′ = 3,2; 4,2; or 5,3), 207–18
Subvalent group 14 metal compounds. Part 10. Syntheses and structures of some bis(metallo)plumbylenes $[Pb\{Mo(R)(CO)_3\}_2(OC_4H_8)]$ [R = η-C_5H_5, η-$C_5H_3(SiMe_3)_2$-1,3, or η-C_5Me_5]; *X*-ray crystal structures of $[Pb\{Mo(\eta\text{-}C_5Me_5)(CO)_3\}_2(OC_4H_8)]$ and $[\{Pb[Mo(\eta\text{-}C_5Me_5)(CO)_3][Mo(\eta\text{-}C_5Me_5)(CO)_2(\mu\text{-}CO)]\}_2]$, 2635–42
LITHIUM
Nucleophilic attack upon ethylene co-ordinated to mercury(II): a molecular-orbital study concerning the origin of the acceleration by Hg^{II}, 361–8
A structural study of the complexation of the lithium ion by the cryptand 4,7,13-trioxa-1,10-diazabicyclo[8.5.5]icosane, 489–92
Reactions of aluminopolyhydride complexes of tungsten. *X*-Ray crystal structures of $[(Me_3P)_3H_3W(\mu\text{-}H)_2AlCl(\mu\text{-}N{=}CHEt)]_2$, $\{[(Me_3P)_3WH_5]Li\}_4$, $WH_2I(PMe_3)_3(SiMe_3)$, and $(Me_3P)_3H_2W(\mu\text{-}H)_3AlCl_2(NMe_3)$, 837–46
The ring-stacking principle in organolithium chemistry: its development through the isolation and crystal structures of hexameric iminolithium clusters (RR′C=NLi)$_6$ (R′ = Ph, R = Bu^t or Me_2N; R = R′ = Me_2N or Bu^t), 1071–82
The syntheses and crystal structures of two iminolithium adducts: $(Ph_2C{=}NLi{\cdot}NC_5H_5)_4$, a tetrameric pseudo-cubane formed by stacking two $(LiN)_2$ four-membered rings, and $[Bu^t_2C{=}NLi{\cdot}OP(NMe_2)_3]_2$, a dimer prevented sterically from stacking, 2141–8
LUMINESCENCE
Substitution reactions and luminescence spectra of the mixed complexes $[ReCl_xBr_{6-x}]^{2-}$, 1929–32
MACROCYCLE
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 5. Unsubstituted fifteen-membered-ring macrocycles containing 2,2′:6′,2″-terpyridine, 5–8
Binucleating N_6 24- and 26-membered macrocyclic ligands. Part 1. Dilead complexes: *X*-ray crystal structure determination of a macrocyclic dilead complex containing nitrogen-only bridging thiocyanate, 123–8
Preparation and magnetism of the binuclear iron(II) complexes $[\{Fe(C_9H_{21}N_3)X\}_2]$ (X = NCS, NCO, or N_3) and their reaction with NO. Crystal structures of $[\{Fe(C_9H_{21}N_3)(NCS)\}_2]$ and $[Fe(C_9H_{21}N_3)(NO)(N_3)_2]$, 187–92
Complexes of ligands providing endogenous bridges. Part 1. The syntheses and crystal structures of barium and lead(II) complexes of macrocyclic Schiff bases derived from heterocyclic dicarbonyls and 1,*n*-diamino-*n*′-hydroxyalkanes (*n*,*n*′ = 3,2; 4,2; or 5,3), 207–18
Complexes of ligands providing endogenous bridges. Part 2. The non-template synthesis of tetraimine Schiff base macrocycles derived from 1,3-diamino-2-hydroxypropane, 219–20
Studies on transition-metal macrocyclic complexes. Single-crystal *X*-ray structure and electrochemistry of the bis-macrocyclic complex $[Cu(L)_2][ClO_4]_2{\cdot}2H_2O$ (L = 1,4,7-triazacyclononane), 373–8
Stereochemical role of lone pairs in main-group elements. Part 3. Structure and bonding in trichloro(1,4,7,10,13-pentaoxacyclopentadecane)antimony(III) studied by means of *X*-ray crystallography at 120 K, 427–30
A structural study of the complexation of the lithium ion by the cryptand 4,7,13-trioxa-1,10-diazabicyclo[8.5.5]icosane, 489–92
Studies of pendant-arm macrocyclic ligands. Part 5. Synthesis of two pyridine-containing penta-aza macrocycles with single pyrrolidinyl pendant arms and characterisation of their nickel(II) and copper(II) complexes. Crystal structure of perchlorato{7-[2′-(1″-pyrrolidinyl)ethyl]-3,7,11,17-tetra-azabicyclo[11.3.1]-heptadeca-1(17),13,15-triene}nickel(II) perchlorate, 545–50
Amino-acid complexation to a cobalt(III) complex of optically-active (2*R*,5*R*,8*R*,11*R*)-2,5,8,11-tetraethyl-1,4,7,10-tetra-azacyclododecane (L). Preparation, characterization, and structure determinations of $[Co^{III}\{(R)\text{- or }(S)\text{-alaO}\}L]Br{\cdot}[ClO_4]{\cdot}H_2O$, 721–8
The chemistry of vitamin B_{12}. Part 27. The cage phosphite ester 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane and the anionic dimethyl phosphite as ligands for cobalt(III) corrinoids, 761–8
Dicopper(II) complexes of a 28-membered N_8 macrocycle. Evidence for proton transfer from co-ordinated secondary amine to thiocyanate, leading to formation of an isothiocyanic acid complex, 873–80
Synthesis of pyridine-containing tetra-aza macrocycles: 3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), its 3,11-dibenzyl (L^2) and 3,7,11-tribenzyl (L^3) derivatives, and their nickel(II), copper(II), and zinc(II) complexes: crystal structures of $L^2{\cdot}HCl$ and $[Ni(L^2)Cl]ClO_4{\cdot}H_2O$, 889–94
Copper(II) complexes of the 14-membered *cis*-N_2S_2 macrocycle 14,14,16-trimethyl-2,6-dithia-13,17-diazatricyclo[16.4.0.0^{7,12}]-docosa-1(18),7(12),8,10,16,19,21-heptaene, 963–4
Studies of the pyridine-containing tetra-azamacrocycle 3,7,11-trimethyl-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1) and its complexes with Ni^{2+}, Cu^{2+}, and Zn^{2+}. Crystal structures of the unsymmetric isomer of $[Ni(L^1)][ClO_4]_2$, and the five-co-ordinate complexes $[Ni(L^1)(dmso)][ClO_4]_2$ (dmso = dimethyl sulphoxide) and $[Ni(L^1)Cl]ClO_4$, 1107–14
Reactions of copper(II) complexes of optically active *N*-substituted diamines with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of optically active macrocycles, 1127–32
Crystal and molecular structure of a novel mixed-spin binuclear nickel(II) complex with *N*,*N*′,*N*″,*N*‴-tetrakis(2-aminoethyl)-1,4,8,11-tetra-azacyclotetradecane, 1261–2
The chemistry of vitamin B_{12}. Part 28. Crystal structure of dicyanocobyrinic acid heptamethyl ester and its interaction with alcohols: the effects of hydrogen bonding to co-ordinated cyanide, 1349–58
Ring-size and substituent effects in relatively rigid macrocyclic quinquedentate ligands. Structure determinations of $[ZnL^4(H_2O)_2][ClO_4]_2{\cdot}MeCN$ and $[ZnL^3(ClO_4)_2]$, 1631–6
Syntheses, *X*-ray structures, and properties of complexes of macrocyclic hexaimines with lanthanide nitrates, 1665–74
Correlation between thermal stability and molecular and crystal structure in a series of potassium–dibenzo-18-crown-6 solid compounds, 1779–84
On the induced circular dichroism of *d*–*d* transitions of a 14-membered macrocyclic tetra-amine nickel(II) complex in β-cyclodextrin, 1801–4
Synthesis of *SS*′-bis(4′-benzo[15-crown-5])dithioglyoxime and its complexes with copper(II), nickel(II), cobalt(II), cobalt(III), palladium(II), platinum(II), and platinum(IV), 1827–32
Complexes of ligands providing endogenous bridges. Part 3. Non-transition metal complexes of macrocycles derived from 2,6-diacetyl-4-methylphenol; the crystal structure of a mononuclear barium complex, 1865–8
Photochemical reduction of carbon dioxide to carbon monoxide in water using a nickel(II) tetra-azamacrocycle complex as catalyst, 2105–10
Syntheses and properties of complexes of a conjugated macrocyclic hexamine with lanthanide nitrates, 2289–90
Iron–ligand bonding properties of synthetic iron–porphyrin complexes with oxygen transporting ability in aqueous media, 2455–8
Dynamics of oxygen binding to modified iron(II)–porphyrin complexes embedded in the bilayer of phospholipid liposome, 2493–6
A realistic model for heme-containing catalases and peroxidases: the *X*-ray structural characterisation of a non-porphyrin iron(III) macrocyclic complex, and the mechanism of its peroxidation of

MACROCYCLE (contd)
aromatic substrates, 2505–10
Disproportionation, electrochemistry, and electronic coupling involving mononuclear and binuclear cobalt phthalocyanine derivatives, 2511–8
Complexes of ligands providing endogenous bridges. Part 4. Copper(II) complexes of macrocyclic Schiff bases derived from 2,6-diacetylpyridine and 1-*n*-diamino-*n'*-hydroxyalkanes (*n,n'* = 3,2; 4,2; and 5,3): synthesis, properties, and structures, 2519–30
Studies involving mixed donor macrocyclic ligands. Comparative *X*-ray diffraction, infrared, and fast atom bombardment mass spectrometric studies of the nickel thiocyanate complexes of three O_2N_3-donor macrocycles, 2537–42
Studies of metal-ion recognition. The interaction of Co^{II}, Ni^{II}, and Cu^{II} with new oxygen–nitrogen donor macrocycles; *X*-ray structures of complexes of Cu^{II} and Ni^{II} with a 15-membered O_2N_3 derivative, 2543–54
Formation of a (4 + 4) Schiff-base macrocyclic ligand by a template rearrangement. Crystal and molecular structures of two tetranuclear manganese(II) complexes, 2555–62
Molecular mechanics studies of a series of dicopper complexes of a 20-membered macrocycle containing differing types of bridges between the metal centres, 2563–72
Synthetic and equilibrium studies on copper(II), nickel(II), and palladium(II) complexes of *N,N',N'',N'''*-tetrakis(2-hydroxyethyl)-1,4,8,11-tetra-azacyclotetradecane (L); kinetics of the acid dissociation of the copper(II) complex and *X*-ray structural characterisation of the nickel(II) complex $[NiLH_{-1}][ClO_4]$, 2605–14
Bicyclam [6,6'-bi(1,4,8,11-tetra-azacyclotetradecane)]: a ditopic receptor for homo- and hetero-bimetallic complexes, 2631–4
Nickel(II), copper(II), and zinc (II) complexes of penta-azamacrocyclic ligands. Crystal and molecular structure of 3,6,9,12-tetramethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-trienezinc(II) perchlorate, 2643–8
The effect of increase in chelate ring size beyond six-membered on the metal ion size selectivity patterns of tetra-aza macrocycles, 2911–6
Oxygen- and carbon monoxide-binding to a lipophilic diporphyrinatocopperiron complex solubilized in an aqueous medium with a micelle, 2963–8
Binuclear copper(I) complexes of a 24-membered macrocycle (L^1) derived from 2,5-diformylthiophene and 3-thiapentane-1,5-diamine. Crystal and molecular structure of $[Cu_2L^1(NCS)][ClO_4]$; a unique example of a single end-to-end bridging thiocyanate between copper(I) atoms, 2975–80
Preparation and structure of nickel(II) complexes of N-substituted tetra-aza macrocycles: molecular structure of [(1*RS*,4*SR*,5*RS*,8*SR*,11*RS*,12*SR*)-1,5,8,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane]nickel(II) diperchlorate, 3093–8
An electron spin resonance study of frozen aqueous solutions containing 5,10,15,20-tetrakis(*N*-methyl-4'-pyridinio)-porphyrinatocobalt(II), 3099–102

MAGNETISM
Magnetic and electrochemical studies on the series $[Cr_nFe_{3-n}(\mu_3\text{-}O)(gly)_6(H_2O)_3][NO_3]_7\cdot xH_2O$(gly = glycine; n = 0, 1, 2, or 3) and on the mixed-valence complex $[Fe^{II}Fe^{III}_2(\mu_3\text{-}O)(gly)_6(H_2O)_3]Cl_6$, 97–100
Preparation and magnetism of the binuclear iron(II) complexes $[\{Fe(C_9H_{21}N_3)X_2\}_2]$ (X = NCS, NCO, or N_3) and their reaction with NO. Crystal structures of $[\{Fe(C_9H_{21}N_3)(NCS)_2\}_2]$ and $[Fe(C_9H_{21}N_3)(NO)(N_3)_2]$, 187–92
Magnetostructural correlations in parallel square-planar chloride bridged copper(II) dimers: structure, dynamic nuclear magnetic resonance study, and magnetic properties of $[Cu_2(terpy)_2Cl_2][PF_6]_2$, 285–92
Magnetic exchange interactions in perovskite solid solutions. Part 7. A study of the non-stoicheiometric phase $SrCoO_{3-y}$ by ^{57}Fe Mössbauer spectroscopy, 667–72
Metal–ligand bonding parameters and magnetic properties of some previously reported tetragonal nickel(II) complexes, 825–30
Ferromagnetic tetrahalogenochromates(II) containing mixed halides (Cl and Br), 965–8
Mixed oxidation state trinuclear cobalt complexes with bridging sulphito and Schiff-base ligands. Part 1. Preparation of the complexes $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}L)_2Co^{III}_2(ROH)_2]$ (L = Schiff base anion, R = alkyl) and structure determination of $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}\alpha,\alpha'\text{-}Me_2\text{-}salpd)_2Co^{III}_2(Pr^nOH)_2]\cdot 2Pr^nOH$, 1305–12
Spectroscopic, magnetic, and structural properties of diaquadimethanolbis(*N*-tosylvalinato)copper(II) and its 2,2'-bipyridine adduct, 1333–40
New spin-crossover iron(III) complexes with large hysteresis effects and time dependence of their magnetism, 1341–8
Magnetic exchange interactions in perovskite solid solutions. Part 8. A study of the solid solution $SrFe_{1-x}Co_xO_{3-y}$ by iron-57 Mössbauer spectroscopy, 1419–24
Structure and magnetism of analogous *O*-bridged (phenolato) and *S*-bridged (thiophenolato) binuclear copper(II) complexes. Crystal structure of the binuclear *S*-bridged complex μ-[2,6-bis(4'-cyclohexyl-4'-hydroxy-2',3'-diazabuta-1',3'-dien-1'-yl)-4-methylthiophenolato(3−)-*S,N*^{2},*O*(Cu^{1,2})]-methanol-μ-(pyrazolyl-*N,N'*)-dicopper(II), 1585–92
Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNN'N'*-tetra-acetate (cdta): structural and magnetic characterization of $[(H_2O)_4Cu(cdta)Ni]\cdot 3H_2O$ and $[(H_2O)_5Ni(cdta)Cu]\cdot H_2O$, 1847–52
X-Ray structural study of [Fe(saltrien)]X (X = Br·2H_2O, BPh_4, or PF_6). Origin of unusual magnetic behaviour of the spin-crossover complex $[Fe(saltrien)]PF_6$, 1957–62
New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states, 2621–30

MANGANESE
Thiosemicarbazones as co-ordinating agents. Part 2. Synthesis, spectroscopic characterization, and *X*-ray structure of aquachloro(pyridoxal thiosemicarbazone)manganese(II) chloride and aqua(pyridoxal thiosemicarbazonato)copper(II) chloride monohydrate, 227–34
The synthesis of silica-supported phosphine and phosphine oxide complexes of manganese(II) iodide and their reaction with sulphur dioxide, 1415–8
Bimetallic systems. Part 22. Molybdenum(0)– and manganese(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the reversible uptake of carbon monoxide: crystal structure of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)\text{-}Mo(CO)_4]\cdot 0.5C_6H_5Cl$, 1509–14
Preparation and reactivity of halide-bridged binuclear carbonyl manganese and rhenium anions, $[M_2(\mu\text{-}X)(CO)_{8-2n}\{(EtO)_2POP(OEt)_2\}_n]^-$ (X = halide; M = Mn, n = 1 or 2; M = Re, n = 1). *X*-Ray crystal structure of $[Mn_2Au(\mu\text{-}Br)(CO)_6(\mu\text{-}tedip)(PPh_3)]$, 1551–6
Synthesis and substitution reactions of a heterodimetallic molybdenum–manganese complex, $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_6]$; *X*-ray crystal structure of $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_4(dppm\text{-}PP')]$, 1557–64
Comments on the attempts by Wickens and Abrams to synthesize manganese(II) phosphine complexes: a reply, 1581–2
The reversible one-electron oxidation of *trans*-$[Mn^IL(CO)(Ph_2PCH_2PPh_2)_2]^{+}$: synthesis and characterisation of stable manganese(II) derivatives, 1769–74
The co-ordination of small molecules by manganese(II) phosphine complexes. The reaction of some $[MnX_2L]$ (X = Cl, Br, or I; L = PPr^n_3, PMe_2Ph, or PPh_3) complexes with tetracyanoethylene, 1785–8
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 8. Irreversible oxidation of trimethylphosphinemanganese(II) dihalides, $[MnX_2(PMe_3)]$ (X = Cl, Br, or I), to bis(trimethylphosphine)manganese(III) trihalides, $[MnX_3(PMe_3)_2]$. *X*-Ray crystal structure of *trans*-triiodobis(trimethylphosphine)manganese(III), the first example of a trigonal-bipyramidal manganese(III) tertiary phosphine complex, 1999–2004
Synthesis and characterisation of some new manganese(II) and manganese(III) complexes of tetradentate Schiff-base ligands and their reaction with molecular oxygen, 2009–12
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 9. The reaction of ethylene with $[MnX_2L]$ (X = Cl, Br, or I; L = PPr^n_3, PBu^n_3, $PPhMe_2$, $PPhEt_2$, or PPh_3) in tetrahydrofuran solution, 2241–4
Synthesis and crystal structure of tris(morpholinecarbodithioato-*S,S'*)manganese(IV) pentaiodide–dichloromethane (2/1), 2407–10
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 10. The reversible binding of nitric oxide by $[MnX_2(PR_3)]$ (X = Cl, Br, I; R_3 = Pr^n_3, Bu^n_3, $PhMe_2$, $PhEt_2$) in the solid state and in tetrahydrofuran solution. The effect of

MANGANESE (contd)
the nature of the halogen and the phosphine on stability and reversibility, 2497–502
Formation of a (4 + 4) Schiff-base macrocyclic ligand by a template rearrangement. Crystal and molecular structures of two tetranuclear manganese(II) complexes, 2555–62
Metallo-ligands containing terminal oxygen donors bonded to a manganese–carbonyl moiety, 2599–604
The reaction of $[Mn_2\{\sigma:\eta^2\text{-}C(R)=CHR\}(\mu\text{-}PPh_2)(CO)_7]$ (R = H or Ph) with nucleophiles: *X*-ray crystal structures of $[N(PPh_3)_2][Mn_2(\mu\text{-}H)\{\mu\text{-}C(O)C(Ph)=CHPh\}(\mu\text{-}PPh_2)(CO)_6]$, $[Mn_2\{PPh_2C(Ph)=CHPh\}(CO)_9]$, $[Mn_2(\mu\text{-}PPh_2CH=CH_2)(CO)_8]$, $[Mn_2\{\mu\text{-}C(O)C(Ph)=CHPh\}(\mu\text{-}PPh_2)(CO)_6(CNBu^t)_2]$, $[Mn_2\{\mu\text{-}\sigma:\eta^2\text{-}C(Ph)=CHPh\}(\mu\text{-}PPh_2)(CO)_6(CNBu^t)]$, and $[Mn(\mu\text{-}PPh_2CH=CH_2)(CO)_7(PEt_3)]$, 2669–82
Synthesis using metal atoms of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[OsH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[MnH(\eta\text{-}C_6H_6)\text{-}(PMe_3)_2]$, and $[Mn(\eta\text{-}C_5H_5)(PMe_3)_3]$: arene ligand-exchange reactions of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, 2823–30
Electrolytic behaviour of transition metal bromides in acetonitrile solution, 2863–6
σ-Alkynyl complexes of manganese(I) as η^2-bonding ligands for Group 1B metal–ligand fragments. *X*-Ray crystal structure of $[Mn_2Cu(\mu\text{-}CCBu^t)_2(CO)_6(dppe)_2]PF_6\cdot 0.5CH_2Cl_2$, 2867–74
Manganese(I) complexes with (tricyclohexylphosphonio)dithiocarboxylate as chelate and unidentate ligand. *X*-Ray crystal structure of *fac*-$[Mn(CO)_3\{S_2CP(C_6H_{11})_3\}_2]ClO_4\cdot H_2O$, 2875–80
A comparison of the efficiencies of polymer-supported phosphine and phosphine oxide complexes of manganese(II), $MnX_2(C_6H_4PPh_2)$ and $MnX_2\{C_6H_4P(O)Ph_2\}_n$ (*n* = 2 or 4; X = Cl, Br, or I), for the absorption of sulphur dioxide, 2959–62
MASS SPECTROMETRY
Bonding in and gas-phase pyrolysis of the arsoranes and stibanes EMe_3X_2 (E = As or Sb, X = F or Cl) and $SbMe_4F$: ultraviolet photoelectron and field-ionization mass spectrometric studies, 481–8
Matrix isolation and mass spectrometric studies on the vaporisation of chromium(III) chloride. The characterisation of molecular $CrCl_4$ and $CrCl_3$, 859–66
Studies involving mixed donor macrocyclic ligands. Comparative *X*-ray diffraction, infrared, and fast atom bombardment mass spectrometric studies of the nickel thiocyanate complexes of three O_2N_3-donor macrocycles, 2537–42
MATRIX ISOLATION
Matrix isolation studies of monomeric boron trisulphide, 267–70
Matrix isolation and mass spectrometric studies on the vaporisation of chromium(III) chloride. The characterisation of molecular $CrCl_4$ and $CrCl_3$, 859–66
Photoactivation of methane by η^5-cyclopentadienyl and substituted η^5-cyclopentadienyl Group 8 metal dicarbonyl complexes, $[M(\eta^5\text{-}C_5R_5)(CO)_2]$ (M = Rh or Ir, R = H or Me), and dicarbonyl(η^5-indenyl)iridium: a matrix isolation study, 1181–90
A quantitative approach to host–guest interactions for matrix-isolated alkali-metal salts of hexafluoroanions and perchlorates, 1521–8
MECHANISM
Mechanism of the reaction between *trans*-$[W(NNH_2)(p\text{-}MeC_6H_4SO_3)(Ph_2PCH_2CH_2PPh_2)_2]^+$ and NEt_3: the influence of the metal on the rates of proton transfer of hydrazido(2−)-ligands, 197–200
Synthesis of the alkoxo(hydrido)-clusters $[Ru_3(\mu\text{-}H)(CO)_{10}(\mu\text{-}OR)]$ [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4
Kinetics and mechanism of ring closure for a series of photoproduced $[M(CO)_5L]$ intermediates (M = Cr, Mo, or W; L = pyridine 2-carbaldehyde imine ligand), 513–8
The chemistry of the hydrazido(1−)-ligand. Preparations and crystal structures of $[Mo(NHNHCO_2Me)(NNCO_2Me)(S_2CNMe_2)_2]$, $[Mo(NHNMePh)(NNMePh)(S_2CNMe_2)_2]BPh_4$, and $[ReCl_2(NHNHCOPh)\text{-}(NNHCOPh)(PPh_3)_2]$. Mechanism of formation of substituted hydrazines from $[Mo(NNRPh)_2(S_2CNMe_2)_2]$ (R = Me or Ph), 529–40
Crystal structure of *trans*-dicarbonylbis(dimethylphenylphosphine)-(*N*-phenylsalicylideneaminato)rhenium(I) and *mer*-trichloro-(dimethylphenylphosphine) (*N*-phenylsalicylideneaminato)-rhenium(IV). Comparison of Re^I, Re^{III}, Re^{IV}, and Re^V complexes with the salicylideneaminato ligand and reaction mechanisms, 613–8
Reaction of monoborane with tris(cyclopentadienyl)uranium(IV) derivatives, 691–4
Substitution reactions between penta-ammineaquaruthenium(II) and thiocyanato, protonated acetato, cyano, and oxalato ligands in aqueous acidic solutions, 957–60
The relative nucleophilic efficiency of water and the azide and thiocyanate ions in the aqueous substitution of chloropentacyanocobaltate(III), 1059–62
Conversion of uncharged methyl complexes of ruthenium(II) into a cationic acyl complex: a mechanistic study, 1713–6
Synthesis, characterisation, and aquation kinetics of [*N*-(*o*-carboxyphenyl)iminodiacetato](glycinato)chromate(III) complex ion in acidic media, 2095–8
The relative nucleophilic efficiency of water and dimethyl sulphoxide in the aqueous substitution of aquapentacyanocobaltate(III) and chloropentacyanocobaltate(III), 2311–6
A realistic model for heme-containing catalases and peroxidases: the *X*-ray structural characterisation of a non-porphyrin iron(III) macrocyclic complex, and the mechanism of its peroxidation of aromatic substrates, 2505–10
New decomposition pathways for formyl complexes of ruthenium(II), 2683–6
Hydride addition reactions of the olefin complexes $[Fe(\eta^5\text{-}C_5H_5)\text{-}(CO)_2(olefin)][BF_4]$; formyl formation as the kinetically preferred process; regio- and stereo-selectivity during addition to the co-ordinated olefins, 2857–62
Ligand intra- and inter-molecular exchange processes in dioxopentakis(trimethylurea)uranium(VI). A proton nuclear magnetic resonance study, 2885–8
The mechanism of the photochemical production of hydrogen from aqueous solutions of hydridotris(triethylphos-phine)platinum(II), 2941–6
Reactions in mixed non-aqueous solution containing sulphur dioxide. Part 9. Mechanisms of dissolution of metals, oxides, and sulphites, 2995–8
MELT
Molten lithium carbonate–sodium carbonate–potassium carbonate eutectic: electronic spectroscopy of first-row transition metal compounds, 2329–32
MENTHYL
Menthyl-substituted organophosphorus compounds. Part 6. Preparation of P(L-men)RCl and P(L-men)R(S)Cl (R = Me, Et, Pr^i Bu^t, or Ph; men = *cyclo*-C_6H_9-2-Pr^i-5-Me) and nuclear magnetic resonance studies of configuration and halogen exchange, 795–806
MERCURY
Synthesis and properties of heterotrimetallic complexes containing M–Pt–Hg (M = Mo, W, Fe, Mn, or Co) arrays, 119–22
Nucleophilic attack upon ethylene co-ordinated to mercury(II): a molecular-orbital study concerning the origin of the acceleration by Hg^{II}, 361–8
Methyl- and phenyl-mercury(II) derivatives of 2-mercaptopyridine. Crystal and molecular structure of methyl(pyridine-2-thiolato)-mercury(II) (1986, 1945), 683–4
Bimetallic systems. Part 18. Nickel(II)–mercury(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$ ligands, 777–80
Neutral and anionic 2-methyl-6-nitrophenyl palladium(II) complexes. The *X*-ray crystal structure of bis(benzyltriphenylphosphonium) *trans*-di-μ-chloro-bis[chloro(2-methyl-6-nitrophenyl)palladate(II)]–acetone (1/2), 1655–8
Polymeric thiolate complexes of Group 12 metals. Crystal and molecular structures of *catena*-[μ-(3-dimethylammonio-1-propanethiolate)]-dichlorocadmium(II) and bis[3-(dimethylammonio)propyl] disulphide tetrabromocadmate(II), 2391–6
Structure and solvation of mercury(II) iodide and bromide in tetrahydrothiophene solution, 2411–6
METALLABORANE
Reactions of heteroboranes. Synthesis of $[2,2\text{-}(PPh_3)_2\text{-}1,2\text{-}SePtB_{10}H_{10}]\cdot CH_2Cl_2$, its crystal and molecular structure and that of $SeB_{11}H_{11}$, 699–704
Pentamethylcyclopentadienylrhodaborane chemistry. Part 3. The reaction of $[\{Rh(\eta^5\text{-}C_5Me_5)Cl_2\}_2]$ with *closo*-$[B_{10}H_{10}]^{2-}$ and *arachno*-$B_{10}H_{12}$-6,9-$(PMe_2Ph)_2$, and the characterisation of some

METALLABORANE (contd)
novel rhodaboranes by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 2417–30
Gold–boron chemistry. Part 1. Synthetic, structural, and spectroscopic studies on the compounds [5,6-μ-($AuPR_3$)-*nido*-$B_{10}H_{13}$] (R = cyclo-C_6H_{11} or C_6H_4Me-2), 2761–8
Organoruthenaborane chemistry. Part 5. Products of the reaction between *closo*-$[B_{10}H_{10}]^{2-}$ and $[(\eta^6\text{-}C_6Me_6)ClRuB_3H_8]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[\{(\eta^6\text{-}C_6Me_6)_2Ru_2H_4\}RuB_{10}H_8(OEt)_2]$, 2781–8
METALLACYCLOSULPHONE
Metallacyclic complexes. Synthesis of metallathietane-3,3-dioxide (metallacyclosulphone) complexes of platinum(II), palladium(II), and nickel(II); crystal structure of *trans*-2,4-diphenyl-1,1-bis(triphenylarsine)platinathietane-3,3-dioxide, 733–40
METALLAPHENALENE
Chemistry of 2,3-dihydro-2-metallaphenalenes. Part 2. Formation of $\overline{CHMMe_2C(MMe_3)C_{10}H_6}]^x$ *via* base decomposition of [1,8-$(Me_3MCH_2)_2C_{10}H_6]^y$ (M = Ge, $x = 2-$, $y = 0$; M = P, $x = 0$, $y = 2+$); crystal structures of the novel bis-ylide $\overline{CHPMe_2C(PMe_3)C_{10}H_6}$ and of a distorted *peri*-ylide, 1-Me-8-$(Me_3PCH)C_{10}H_6$, 1647–54
METALLOBORANE
Synthesis, molecular structure, and nuclear magnetic resonance spectra of the anionic monocarbametallaborane $[NEt_4][(CO)_3(PPh_3)(OH)$-*closo*-$MoCB_{10}H_{10}]\cdot CH_2Cl_2$, 2385–90
METHANE
Photoactivation of methane by η^5-cyclopentadienyl and substituted η^5-cyclopentadienyl Group 8 metal dicarbonyl complexes, $[M(\eta^5\text{-}C_5R_5)(CO)_2]$ (M = Rh or Ir, R = H or Me), and dicarbonyl(η^5-indenyl)iridium: a matrix isolation study, 1181–90
METHANOL
Formation of a methoxycarbonyl group by attack of methanol on an iron(II) carbonyl complex, assisted by intramolecular proton transfer: *X*-ray structure determination of $[Fe(CO)(PMe_3)_2(CNC_6H_{11})\{CO(OCH_3)\}\{C(CH_3)=NHC_6H_{11}\}]\text{-}BPh_4$, 473–6
Electrocatalytic reduction of carbon dioxide to methanol. Part 5. Relationship between the ability of metal complexes to engage in homogeneous catalysis and their co-ordination chemistry, 1377–80
METHANOLYSIS
Kinetic study of the homogeneous methanolysis of tetraphosphorus decasulphide, 1897–900
METHOXIDE
Binuclear rhodium and iridium complexes containing pentamethylcyclopentadienyl and pyrazolate ligands, 639–46
METHOXYCARBONYL
Formation of a methoxycarbonyl group by attack of methanol on an iron(II) carbonyl complex, assisted by intramolecular proton transfer: *X*-ray structure determination of $[Fe(CO)(PMe_3)_2(CNC_6H_{11})\{CO(OCH_3)\}\{C(CH_3)=NHC_6H_{11}\}]\text{-}BPh_4$, 473–6
METHYL
The synthesis of mono- and bi-metallic complexes containing the ligand bis(diphenylarsino)methane, 151–4
Molecular orbital study of the bridging dioxygen ligand in $[Al_2(CH_3)_6(O_2)]^-$, 355–60
Sulphur–arsenic ligands. Part 1. Dimethyl(methylthio)arsine complexes of halogenotrimethylplatinum(IV) and tricarbonylhalogenorhenium(I), 417–20
Bonding in and gas-phase pyrolysis of the arsoranes and stibanes EMe_3X_2 (E = As or Sb, X = F or Cl) and $SbMe_4F$: ultraviolet photoelectron and field-ionization mass spectrometric studies, 481–8
Interactions of the trimethyltin(IV) cation with carboxylic acids, amino acids, and related ligands, 563–6
Methyl- and phenyl-mercury(II) derivatives of 2-mercaptopyridine. Crystal and molecular structure of methyl(pyridine-2-thiolato)mercury(II) (1986, 1945), 683–4
Electron diffraction study of the molecular structure of *N*-methylbis(trimethylsilyl)amine in the gas phase, 785–8
Crystal structure and mass spectrometry of dichlorodimethylbis[2(1*H*)-pyridinethione-*S*]tin(IV), 815–8
Conversion of uncharged methyl complexes of ruthenium(II) into a cationic acyl complex: a mechanistic study, 1713–6
Platinum(II) complexes with ligands $(RO)_2PCH_2P(OR)_2$ (R = Me, Et, Ph, or C_6H_4Me-4); crystal structure of *cis,cis*-$[Pt_2Me_4\{\mu\text{-}(EtO)_2PCH_2P(OEt)_2\}_2]$, 2117–24
Methylindium dialkylamido compounds: mono- and bis-dialkylamides involving four- and five-co-ordinated indium(III). *X*-Ray crystal structures of $\{Me_2In[\overline{N(CH_2)_2N(Me)CH_2CH_2}]\}_2$, $MeIn[MeN\overline{C(CH)_4N}]_2$, and $MeIn\{[Me_2N(CH_2)_2NMe]\text{-}InMe_2\}_2$, 2159–64
A two-dimensional nuclear magnetic resonance exchange study of the fluxionality of $[PtIMe_3(MeSCH_2CH_2SEt)]$, 2695–702
METHYLENE
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 64. Addition of methylene groups to iron–molybdenum complexes; crystal structures of $[FeMo(\mu\text{-}CH_2)\{\mu\text{-}\sigma{:}\eta\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(CO)_5(\eta\text{-}C_5H_5)]$ and $[FeMo\{\mu\text{-}C(C_6H_4Me\text{-}4)C(OMe)C(H)\}(CO)_5(\eta\text{-}C_5H_5)]$, 2201–10
Preparation of two mixed-metal carbonyl clusters based on the platinumtriosmium core. *X*-Ray crystal structure of $[Os_3PtH_2(cod)(CH_2)(CO)_9]$ (cod = cyclo-octa-1,5-diene), 3103–6
METHYLENEAMIDE
Triosmium clusters containing the ligands CNH_2, NCHCN, and NCMeCN derived from cyanide ion, cyanogen, and acetonitrile, 1457–62
The syntheses and crystal structures of two iminolithium adducts: $(Ph_2C{=}NLi{\cdot}NC_5H_5)_4$, a tetrameric pseudo-cubane formed by stacking two $(LiN)_2$ four-membered rings, and $[Bu^t{}_2C{=}NLi{\cdot}OP(NMe_2)_3]_2$, a dimer prevented sterically from stacking, 2141–8
METHYLENEAMINE
The ring-stacking principle in organolithium chemistry: its development through the isolation and crystal structures of hexameric iminolithium clusters $(RR'C{=}NLi)_6$ (R′ = Ph, R = Bu^t or Me_2N; R = R′ = Me_2N or Bu^t), 1071–82
METHYLIDYNE
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 49. Synthesis of mixed-metal complexes with bonds between Cr, Mo or W and Co, Rh, Ir or Re; crystal structure of $[CrReRh_2(\mu_3\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}CO)(CO)_9(\mu\text{-}C_9H_7)_2]$, 41–50
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 50. Synthesis and crystal structures of the complexes $[RuW_3(\mu\text{-}CO)(\mu_3\text{-}\eta\text{-}CO)(\mu\text{-}CPh)_2(\mu_3\text{-}CPh)(CO)_2(\eta\text{-}C_5H_5)_3]$ and $[RuW_3(\mu\text{-}CO)(\mu_3\text{-}\eta\text{-}CO)\{\mu\text{-}C(Ph)C(O)\}(\mu\text{-}CPh)(\mu_3\text{-}CPh)(CO)(PMePh_2)(\eta\text{-}C_5H_5)_3]$, 51–60
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 51. Alkylidyne(carbaborane)tungsten-gold and -rhodium complexes; crystal structures of $[AuW(\mu\text{-}CR)(CO)_2(PPh_3)(\eta^5\text{-}C_2B_9H_9Me_2)]$, $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$, and $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2\{\eta^5\text{-}C_2B_9(C_7H_9)H_8Me_2\}]$ (R = C_6H_4Me-4), 61–72
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 52. Synthesis and reactions of the compounds $[RuW(\mu\text{-}CR)(CO)_3(\eta\text{-}C_5H_5)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[RuW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta\text{-}C_5H_5)(\eta^5\text{-}C_2B_9H_9Me_2)]$ and $[N(PPh_3)_2][RuW(\mu\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}\sigma,\eta^5\text{-}C_2B_9H_8Me_2)(CO)_3(\eta\text{-}C_5H_5)]$, 73–80
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 53. Alkylidyne(carbaborane)dimetal complexes with tungsten–molybdenum and –tungsten bonds; crystal structures of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ [M = Mo (two isomers) or W], 81–90
MIXED VALENCE
Magnetic and electrochemical studies on the series $[Cr_xFe_{3-x}(\mu_3\text{-}O)(gly)_6(H_2O)_3][NO_3]_7\cdot xH_2O$(gly = glycine; $n = 0$, 1, 2, or 3) and on the mixed-valence complex $[Fe^{II}Fe^{III}{}_2(\mu_3\text{-}O)(gly)_6(H_2O)_3]Cl_6$, 97–100
Mixed oxidation state trinuclear cobalt complexes with bridging sulphito and Schiff-base ligands. Part 1. Preparation of the complexes $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}L)_2Co^{III}{}_2(ROH)_2]$ (L = Schiff base anion, R = alkyl) and structure determination of $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}\alpha,\alpha'\text{-}Me_2\text{-}salpd)_2Co^{III}{}_2(Pr^nOH)_2]\cdot 2Pr^nOH$, 1305–12
Preparation and partial oxidation of tin(II) trifluoroacetate: spectroscopic properties and *X*-ray crystal structure of di-μ_3-oxo-octakis-μ-trifluoroacetato-tetratin(II)tin(IV), 1675–82
Reduction–oxidation properties of organotransition-metal

MIXED VALENCE (contd)
complexes. Part 27. Mixed-valence triazenido-bridged di-iridium compounds, 2737–40
MOLECULAR MECHANICS
Molecular mechanics studies of a series of dicopper complexes of a 20-membered macrocycle containing differing types of bridges between the metal centres, 2563–72
MOLECULAR ORBITAL
Structure and bonding in $(R_2SiX)_2$ molecules (X = O, NH, CH_2, or S; R = H), 271–4
Molecular orbital study of the bridging dioxygen ligand in $[Al_2(CH_3)_6(O_2)]^-$, 355–60
Nucleophilic attack upon ethylene co-ordinated to mercury(II): a molecular-orbital study concerning the origin of the acceleration by Hg^{II}, 361–8
A group theoretical paradigm for describing the skeletal molecular orbitals of cluster compounds. Part 1. Deltahedral clusters, 647–56
Extended Hückel molecular orbital study of the effects of edge-bridging hydrogen atoms on the lengths of boron–boron and metal–metal bonds in cluster compounds, and the crystal structure of benzyltrimethylammonium octahydrotriborate(1−), 1017–26
A group theoretical paradigm for describing the skeletal molecular orbitals of cluster compounds. Part 2. Bispherical clusters, 1445–56
Crystal and molecular structures of $[MCl(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf) and $[TiCl_{0.5}Br_{0.5}(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$: an unusual π-interaction with the metals and molecular orbital calculations on model compounds, 1641–6
Charge density in the $CoCl_4^{2-}$ ion: a comparison with spin density and theoretical calculations, 1737–46
MOLTEN SALT
Concentrated aqueous solutions and liquid hydrates of zinc chloride: a study of their solvent behaviour using intershell, charge-transfer, and *d*–*d* adsorption spectra, 1485–8
MOLYBDENUM
Synthesis of $[Mo(\eta\text{-}C_5H_4Pr^i)(PR_3)_2H_3]$ (R_3 = Me_3 or Me_2Ph), $[Mo(\eta\text{-}C_5H_4Pr^i)(Pr^i{}_2PCH_2CH_2PPr^i{}_2)H_3]$, and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of $[\{Mo(\eta\text{-}C_5H_4Pr^i)(\mu\text{-}Cl)_2\}_2]$, 9–20
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 53. Alkylidyne(carbaborane)dimetal complexes with tungsten–molybdenum and –tungsten bonds; crystal structures of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ [M = Mo (two isomers) or W], 81–90
Mechanism of the reaction between *trans*-$[W(NNH_2)(p\text{-}MeC_6H_4SO_3)(Ph_2PCH_2CH_2PPh_2)_2]^+$ and NEt_3: the influence of the metal on the rates of proton transfer of hydrazido(2−)-ligands, 197–200
Bimetallic systems. Part 16. Formation of Group 6 metal(0)–platinum(II) hydride complexes, by ring opening of *fac*- or *mer*-$[M(CO)_3(dppm\text{-}PP')(dppm\text{-}P)]$ (M = Cr, Mo, or W; dppm = $Ph_2PCH_2PPh_2$). Reactions of $[(CO)_3Mo(\mu\text{-}dppm)_2\text{-}PtH(Cl)]$, 221–6
Structure and properties of tris(quinoxaline-2,3-dithiolato)-molybdate(IV), 297–302
Preparations, crystal structures, and molybdenum-95 nuclear magnetic resonance spectroscopic studies of triphenylphosphine–gold–tetrathiomolybdate(VI) clusters, 303–6
The addition of isonitriles to $[\{M(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (M = Mo or W). Syntheses and properties of $[M_2(\mu\text{-}\eta^2\text{-}CNR)(CO)_4(\eta\text{-}C_5H_5)_2]$ (M = Mo or W, R = Me or Bu^t) and $[\{Mo(CNR)(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (R = Me or Bu^t). *X*-Ray crystal structures of $[Mo_2(\mu\text{-}\eta^2\text{-}CNBu^t)(CO)_4(\eta\text{-}(\eta\text{-}C_5H_5)_2]$ and $[\{Mo(CNBu^t)(CO)_2(\eta\text{-}C_5H_5)\}_2]$, 341–8
Syntheses and reactions of acyclic and cyclic carbene complexes *cis*-$[MI(=CMeNHMe)(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo or W) and *cis*-$[MI\{=\overline{C(CH_2)_n}NMe\}(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo, *n* = 3 or 4; M = W, *n* = 3) obtained through alkyl migration to methyl isocyanide co-ordinated to molybdenum or tungsten. *X*-Ray structure of *cis*-$[MI\{=\overline{C(CH_2)_3}NMe\}(CO)_2(\eta\text{-}C_5H_5)]$ (1986, 2127), 479–80
Kinetics and mechanism of ring closure for a series of photoproduced $[M(CO)_5L]$ intermediates (M = Cr, Mo, or W; L = pyridine 2-carbaldehyde imine ligand), 513–8
Some binuclear molybdenum(VI) complexes obtained from condensation reactions: their crystal and molecular structures, 523–8
The chemistry of the hydrazido(1−)-ligand. Preparations and crystal structures of $[Mo(NHNHCO_2Me)(NNCO_2Me)(S_2CNMe_2)_2]$, $[Mo(NHNMePh)(NNMePh)(S_2CNMe_2)_2]BPh_4$, and $[ReCl_2(NHNHCOPh)\text{-}(NNHCOPh)(PPh_3)_2]$. Mechanism of formation of substituted hydrazines from $[Mo(NNRPh)_2(S_2CNMe_2)_2]$ (R = Me or Ph), 529–40
Reactions of co-ordinated ligands. Part 39. The synthesis of carbyne complexes from alkyne molybdenum and tungsten cations; formation and crystal structures of $[Mo(\equiv CCH_2Bu^t)\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$ and $[\overline{Mo\{=C(SiMe_3)CH_2\}}\{P(OMe)_3\}_2(\eta^5\text{-}C_9H_7)]$, 591–604
A molybdenum-95 and nitrogen-14 nuclear magnetic resonance study of six-co-ordinate hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes containing a sixteen-electron $\{Mo(NO)\}^4$ core, 619–22
Heteronuclear 1,3-intramolecular metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of 2-thia-4-selenapentane. A dynamic nuclear magnetic resonance investigation, 757–60
Mono- and di-carbonyl complexes of molybdenum(II) with mixed bidentate ligands. *X*-Ray crystal structure of $[MoBr_2(CO)(bipy)(Ph_2PCH_2PPh_2)]$ (bipy = 2,2′-bipyridine): a novel type of $[M(bidentate)_2(unidentate)_3]$ structure, 819–24
New heterocyclic compounds containing niobium and molybdenum; crystal structure of a cyclomolybdaphosphazene, 847–50
Studies on transition-metal peroxo complexes. Part 7. Molybdenum(VI) and tungsten(VI) carboxylato peroxo complexes, and the *X*-ray crystal structure of $K_2[MoO(O_2)_2(glyc)]\cdot 2H_2O$, 991–6
Reactions of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}X_2]$ (M = Mo, X = I; M = W, X = Cl) with some amine-substituted pyridines. The molecular structures of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}(2\text{-}NHC_5NH_4)_2]$, 1063–70
The preparation, structure, and reactions of the mononuclear thiolate–hydride complexes $[MoH(SR)(dppe)_2]$ (R = bulkyl alkyl or aryl group, dppe = $Ph_2PCH_2CH_2PPh_2$); *X*-ray structure determination of *cis*-$[MoH(SC_6H_2Pr^i{}_3\text{-}2,4,6)(dppe)_2]$, 1115–22
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 56. Synthesis of iron–molybdenum compounds; crystal structures of $[FeMo(\mu\text{-}CR)(CO)_6(\eta\text{-}C_5H_5)]$ and $[FeMo_2(\mu_3\text{-}RC_2R)(CO)_6(\eta\text{-}C_5H_5)_2]$ (R = C_6H_4Me-4), 1209–14
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 57. Reactions of iron– molybdenum complexes with oxygen and sulphur; crystal structure of $[FeMo(\mu\text{-}\eta^2\text{-}SCC_6H_4Me\text{-}4)(CO)_5(\eta\text{-}C_5H_5)]$, 1215–20
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 58. Some reactions of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (M = Mo or W); crystal structures of the complexes $[W_2(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_2(L)(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$-[L = $N_2C(C_6H_4Me\text{-}4)_2$ or O], 1221–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 59. Reactions of the alkylidyne compounds $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Cr, Mo, or W, R = C_6H_4Me-4; M = W, R = Me, $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate] with d^{10} complexes; crystal structure of $[PtW_2(\mu\text{-}CMe)_2(CO)_4\text{-}\{HB(pz)_3\}_2]$, 1229–34
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 60. Di- and tri-metal compounds prepared from the metal alkylidyne complexes $[W(\equiv CR)(CO)_2(\eta\text{-}C_5Me_5)]$ (R = Me or C_6H_4Me-4) and $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Mo, R = C_6H_4Me-4; M = W, R = Me; $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate], 1235–42
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 61. Reactions of iron– molybdenum compounds with tertiary phosphines, 1243–8
Reactions of co-ordinated ligands. Part 41. Synthesis and reactions of cationic functionalised alkyne molybdenum complexes; formation of carbyne and vinylidene complexes and crystal structure of $[Mo(C{=}CHPh)Br\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$, 1319–28
Reaction paths and kinetics in the Cr^{2+} reduction of the $Mo^V{}_2$ and

MOLYBDENUM (contd)
Mo^{IV}_3 ions $[Mo_2O_4(H_2O)_6]^{2+}$ and $[Mo_3O_4(H_2O)_9]^{4+}$, 1397–402
Bimetallic systems. Part 19. Chromium(0)–, molybdenum(0)–, or tungsten(0)–iron(0) carbonyl complexes containing one or two bridging $Ph_2PCH_2PPh_2$ ligands: crystal structure of $[(OC)_3Mo(\mu\text{-}Ph_2PCH_2PPh_2)_2Fe(CO)_3]$, 1489–94
Bimetallic systems. Part 20. Synthesis of Group 6 metal(0)–iridium(I) or –rhodium(I) complexes bridged by $Ph_2PCH_2PPh_2$, 1495–500
Bimetallic systems. Part 22. Molybdenum(0)– and manganese(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the reversible uptake of carbon monoxide: crystal structure of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)\text{-}Mo(CO)_4]\cdot 0.5C_6H_5Cl$, 1509–14
Synthesis and substitution reactions of a heterodimetallic molybdenum–manganese complex, $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_6]$; *X*-ray crystal structure of $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_4(dppm\text{-}PP')]$, 1557–64
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato-molybdenum and -tungsten nitrosyls. Part 4. The effects of ligating atom type on the reduction potentials of monometallic complexes, 1733–6
Reactions of main group homopolynuclear cations with transition-metal carbonyls. Stabilisation of $\mu\text{-}Se_2^{2+}$ and $\eta^3\text{-}Te_3^{2+}$ dicationic ligands and crystal structures of $[Mo(Te_3)(CO)_4][SbF_6]_2$ and $[FeW(Se_2)(CO)_8][SbF_6]_2$, 2111–6
The hydrolysis of metal ions. Part 10. Kinetic and equilibrium measurements of molybdenum(VI), 2149–58
Some chemistry of the heterobimetallic complexes $[(OC)_3Mo(\mu\text{-}dppm)_2Rh(CO)Cl]$ and $[(OC)_3Mo(\mu\text{-}dppm)_2\text{-}M(CO)_2]PF_6$ (M = Rh or Ir, dppm = $Ph_2PCH_2PPh_2$). Structure of $[(OC)_3Mo(\mu\text{-}dppm)_2(\mu\text{-}C{\equiv}CMe)Rh(CO)]$, 2171–6
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 62. Synthesis of penta-, hexa-, and hepta-heteronuclear metal cluster compounds involving tungsten or molybdenum with platinum or nickel: crystal structures of the chain compounds $[Pt_3W_3(\mu\text{-}CMe)(\mu_3\text{-}CMe)_2(CO)_6(cod)(\eta\text{-}C_5Me_5)_3]\cdot 0.5CH_2Cl_2$ and $[Pt_3W_4(\mu\text{-}CC_6H_4Me\text{-}4)_2(\mu_3\text{-}CC_6H_4Me\text{-}4)_2(CO)_8(\eta\text{-}C_5H_5)_4]$, 2177–88
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 64. Addition of methylene groups to iron–molybdenum complexes; crystal structures of $[FeMo(\mu\text{-}CH_2)\{\mu\text{-}\sigma\text{:}\eta\text{-}C(C_6H_4Me\text{-}4){=}CH_2\}(CO)_5(\eta\text{-}C_5H_5)]$ and $[FeMo\{\mu\text{-}C(C_6H_4Me\text{-}4)C(OMe)C(H)\}(CO)_5(\eta\text{-}C_5H_5)]$, 2201–10
Crystal structure of tris(quinoxaline-2,3-dithiolato)molybdate(V); a distorted trigonal prismatic complex, 2267–72
Ab initio study of stable bis(carbon dioxide)molybdenum complexes, 2373–8
Synthesis, molecular structure, and nuclear magnetic resonance spectra of the anionic monocarbametallaborane $[NEt_4][(CO)_3(PPh_3)(OH)\text{-}closo\text{-}MoCB_{10}H_{10}]\cdot CH_2Cl_2$, 2385–90
Determination of the relative signs of $^2J(^{31}P\text{–}^{31}P)$ in complexes of tungsten(0) and molybdenum(0) using two-dimensional [^{31}P–^{31}P]-COSY-45 nuclear magnetic resonance chemical shift correlation, 2401–6
Subvalent group 14 metal compounds. Part 10. Syntheses and structures of some bis(metallo)plumbylenes $[Pb\{Mo(R)(CO)_3\}_2(OC_4H_8)]$ [R = $\eta\text{-}C_5H_5$, $\eta\text{-}C_5H_3(SiMe_3)_2$-1,3, or $\eta\text{-}C_5Me_5$]; *X*-ray crystal structures of $[Pb\{Mo(\eta\text{-}C_5Me_5)(CO)_3\}_2(OC_4H_8)]$ and $[\{Pb[Mo(\eta\text{-}C_5Me_5)(CO)_3][Mo(\eta\text{-}C_5Me_5)(CO)_2(\mu\text{-}CO)]\}_2]$, 2635–42
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato molybdenum and tungsten nitrosyls. Part 6. Synthesis and electrochemistry of complexes containing saturated heterocyclic amide ligands. *X*-Ray crystal and molecular structure of $[\{Mo(NO)[HB(Me_2pz)_3]Cl\}_2(NC_4H_8N)]$, 2653–60
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η^2-vinyl complexes of molybdenum and tungsten, $[M\{\eta^3\text{-}C(CF_3)C(CF_3)(L\text{–}L)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (L–L = $MeCO_2^-$, $Me_2PS_2^-$, $Et_2NCS_2^-$, or $Me_2NCS_2^-$), 2715–22
Heteronuclear complexes containing Group 6 transition metals; chemical, spectroscopic, and theoretical studies on some binuclear complexes and the *X*-ray crystal structure determination of $[WNi(CO)_3(PPh_3)_2(\eta^5\text{-}C_5H_5)]$, 2741–50
Carbene complexes. Part 21. Synthesis and characterisation of bis(carbene)molybdenum(II) complexes and dimetal(0) complexes of the Group 6 elements containing novel bridging bis(carbene) ligands; *X*-ray structures of $[Mo(CO)_2(L^{Et})_2(OSO_2CF_3)_2]$ [L^{Et} = $\equiv\overbrace{CN(Et)(CH_2)_2}^{}NEt]$ and $[\overbrace{W(CO)_5\{C(OEt)\text{-}CH_2C_6H_4CH_2C(OEt)\text{-}o\}}^{}W(CO)_5]$, 2843–52
Instant ligands. Part 2. Preparation of some bidentate ligands containing PF_2O Groups from aromatic hydroxy compounds and their reactions to form molybdenum and tungsten complexes, 3023–8
Chemistry of molybdenum. Part 5. Synthesis, spectroscopic and electrochemical properties of bis(2-aminocyclopent-1-ene-1-dithiocarboxylato)(arylimido)dihalogenomolybdenum(VI) complexes, 3043–8
The chemistry of cycloheptatrienyl complexes of molybdenum and tungsten: the synthesis and reactions of some vinylidene and alkynyl derivatives, 3049–54

MOLYBDENUM-95
Preparations, crystal structures, and molybdenum-95 nuclear magnetic resonance spectroscopic studies of triphenylphosphine–gold–tetrathiomolybdate(VI) clusters, 303–6
A molybdenum-95 and nitrogen-14 nuclear magnetic resonance study of six-co-ordinate hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes containing a sixteen-electron $\{Mo(NO)\}^4$ core, 619–22

MÖSSBAUER
Organotin biocides. Part 8. The crystal structure of triphenyltin formate and a comparative variable-temperature tin-119 Mössbauer spectroscopic study of organotin formates and acetates, 101–6
Mössbauer spectroscopic studies on molecular rotation and lattice dynamics of cyclodextrin clathrates of bis(η-cyclopentadienyl)-iron(II) and some of its ring-substituted derivatives, 627–32
Magnetic exchange interactions in perovskite solid solutions. Part 7. A study of the non-stoicheiometric phase $SrCoO_{3-y}$ by ^{57}Fe Mössbauer spectroscopy, 667–72
The configuration and lattice dynamics of complexes of dialkyltin(IV) with adenosine 5′-monophosphate and phenyl phosphates, 789–94
New spin-crossover iron(III) complexes with large hysteresis effects and time dependence of their magnetism, 1341–8
Magnetic exchange interactions in perovskite solid solutions. Part 8. A study of the solid solution $SrFe_{1-x}Co_xO_{3-y}$ by iron-57 Mössbauer spectroscopy, 1419–24
Complexes of 2,5,7,9-tetrathiabicyclo[4.3.0]non-1(6)-ene-8-thione (L^1). Part 1. Synthesis and antimony-121 Mössbauer spectroscopic characterisation of $SbCl_3(L^1)_{1.5}$, crystal and molecular structures of $SbCl_3(L^1)_{1.5}$ and uncomplexed L^1, 1605–10
Preparation and partial oxidation of tin(II) trifluoroacetate: spectroscopic properties and *X*-ray crystal structure of di-μ_3-oxo-octakis-μ-trifluoroacetato-tetratin(II)tin(IV), 1675–82
Iron antimonate, 1727–32
Transition metal Schiff-base complexes as ligands in tin chemistry. Part 1. A tin-119 Mössbauer spectroscopic investigation of the adducts $SnX_4\cdot ML$, $SnMe_2(NCS)_2\cdot ML$, and $SnRCl_3\cdot ML$ (X = halide; M = Cu^{II} or Ni^{II}; L = quadridentate Schiff-base ligand; R = phenyl or n-butyl), 2261–6
Gold-197 Mössbauer spectra of some gold–ruthenium cluster compounds, 2333–6
The reaction of di-2-pyridyl ketone with antimony(III) fluoride. Preparation, properties, Mössbauer spectrum and crystal structure of difluoro(methoxydi-2-pyridylmethoxo-*NN′O*)antimony(III), 2337–40
Preparation, crystal and molecular structure, and antimony-121 Mössbauer spectrum of trifluoro(1,10-phenanthroline)(thiourea)-antimony(III), 2451–4
Iron–ligand bonding properties of synthetic iron–porphyrin complexes with oxygen transporting ability in aqueous media, 2455–8

NEUTRON DIFFRACTION
Determination of the deuterium position in DNb_6I_{11}, 431–4

NICKEL
Co-ordination abilities of thioamide analogues of amino acids and peptides. Copper(II) and nickel(II) complexes, 1–4
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 5. Unsubstituted fifteen-membered-ring macrocycles containing 2,2′:6′,2″-terpyridine, 5–8
Reactions between bis(*O*-alkyl dithiocarbonato)nickel(II) complexes

NICKEL (contd)
and phosphines. Formation of a dithiocarbonate complex of nickel(II): $[Ni(S_2CO)(Ph_2PCH_2CH_2PPh_2)]$, 281–4
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution, 435–42
Studies of pendant-arm macrocyclic ligands. Part 5. Synthesis of two pyridine-containing penta-aza macrocycles with single pyrrolidinyl pendant arms and characterisation of their nickel(II) and copper(II) complexes. Crystal structure of perchlorato{7-[2′-(1″-pyrrolidinyl)ethyl]-3,7,11,17-tetraazabicyclo[11.3.1]heptadeca-1(17),13,15-triene}nickel(II) perchlorate, 545–50
The vibrational spectrum of tricarbonyl(trichlorostibine)nickel(0), $[Ni(CO)_3(SbCl_3)]$, 681–2
Vibrational studies of interstitial hydrogen in metal carbonyl clusters, 685–90
Metallacyclic complexes. Synthesis of metallathietane-3,3-dioxide (metallacyclosulphone) complexes of platinum(II), palladium(II), and nickel(II); crystal structure of *trans*-2,4-diphenyl-1,1-bis(triphenylarsine)platinathietane-3,3-dioxide, 733–40
Bimetallic systems. Part 18. Nickel(II)–mercury(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$ ligands, 777–80
Metal–ligand bonding parameters and magnetic properties of some previously reported tetragonal nickel(II) complexes, 825–30
Reactions of cyclobutenedione and cyclopentenetrione with zero-valent nickel triad complexes of isocyanide. *X*-Ray crystal structures of (3—4-η-3,4-diphenylcyclobut-3-ene-1,2-dione)- and (4—5-η-4,5-diphenylcyclopent-4-ene-1,2,3-trione)-tris(t-butyl isocyanide)nickel, 853–8
Synthesis of pyridine-containing tetra-aza macrocycles: 3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), its 3,11-dibenzyl (L^2) and 3,7,11-tribenzyl (L^3) derivatives, and their nickel(II), copper(II), and zinc(II) complexes: crystal structures of L^2·HCl and $[Ni(L^2)Cl]ClO_4·H_2O$, 889–94
Thermally induced phase transition and *X*-ray crystal structure of bis(*N*,*N*′-dimethyl-1,2-ethanediamine)di-isothiocyanatonickel(II), 997–1000
Studies of the pyridine-containing tetra-azamacrocycle 3,7,11-trimethyl-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1) and its complexes with Ni^{2+}, Cu^{2+}, and Zn^{2+}. Crystal structures of the unsymmetric isomer of $[Ni(L^1)][ClO_4]_2$, and the five-co-ordinate complexes $[Ni(L^1)(dmso)][ClO_4]_2$ (dmso = dimethyl sulphoxide) and $[Ni(L^1)Cl]ClO_4$, 1107–14
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 54. Compounds derived from $[W(\equiv CC_6H_4Me\text{-}4)(CO)_2(\eta\text{-}C_5Me_5)]$ with bonds between tungsten and iron, cobalt, or nickel; crystal structure of $[FeW\{\mu\text{-}C(C_6H_4Me\text{-}4)C(O)O\}(CO)_5(\eta\text{-}C_5Me_5)]$, 1191–200
Crystal and molecular structure of a novel mixed-spin binuclear nickel(II) complex with *N*,*N*′,*N*″,*N*‴-tetrakis(2-aminoethyl)-1,4,8,11-tetra-azacyclotetradecane, 1261–2
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1283–4
Design of transition metal chelates with biological activity. Potentiometric study of complex formation equilibria between 2-amino-*N*-hydroxy-3-(*p*-hydroxyphenyl)propanamide and nickel(II), copper(II), and hydrogen ions in aqueous solution, 1409–14
Bimetallic systems. Part 21. Synthesis of vinylidene- and phenylene-bridged complexes of dinickel(II), nickel(II)–palladium(II), and nickel(II)–platinum(II) containing $Ph_2PCH_2PPh_2$ (dppm): crystal structures of $[(SCN)Ni(\mu\text{-dppm})_2(\mu\text{-}C{=}CH_2)Ni(NCS)]$ and $[ClPd(\mu\text{-dppm})_2(\mu\text{-}C{=}CCl_2)NiCl]$, 1501–8
Raman and resonance-Raman studies of platinum group tetrathiomolybdates and tetrathiotungstates, 1535–44
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1583–4
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), and hydrogen ions in aqueous solution (1987, 435), 1583–4
Preparation, structure, and properties of $[Ni\{S_2C_2(CF_3)_2\}_2]^-$ and $[Ni\{Se_2C_2(CF_3)_2\}_2]^-$ salts with small cations, 1683–6
Studies on transition-metal–peptide complexes. Part 13. Copper(II) and nickel(II) complexes of amino acids and peptides containing a thioether group, 1717–20
On the induced circular dichroism of *d*–*d* transitions of a 14-membered macrocyclic tetra-amine nickel(II) complex in β-cyclodextrin, 1801–4
Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNN*′*N*′-tetra-acetate (cdta): structural and magnetic characterization of $[(H_2O)_4Cu(cdta)Ni]·3H_2O$ and $[(H_2O)_5Ni(cdta)Cu]·H_2O$, 1847–52
Thermal investigation and stereochemical studies of some cyclic diamine complexes of nickel(II), zinc(II), and cadmium(II) in the solid state, 2089–94
Photochemical reduction of carbon dioxide to carbon monoxide in water using a nickel(II) tetra-azamacrocycle complex as catalyst, 2105–10
Studies in Werner clathrates. Part 4. Structures of tetrakis(4-ethylpyridine)di-isothiocyanatonickel(II) and its clathrates with *p*-, *m*-, and *o*-xylene, carbon disulphide, and carbon tetrachloride, 2125–40
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 62. Synthesis of penta-, hexa-, and hepta-heteronuclear metal cluster compounds involving tungsten or molybdenum with platinum or nickel: crystal structures of the chain compounds $[Pt_3W_3(\mu\text{-}CMe)(\mu_3\text{-}CMe)_2(CO)_6(cod)(\eta\text{-}C_5Me_5)_3]·0.5CH_2Cl_2$ and $[Pt_3W_4(\mu\text{-}CC_6H_4Me\text{-}4)_2(\mu_3\text{-}CC_6H_4Me\text{-}4)_2(CO)_8(\eta\text{-}C_5H_5)_4]$, 2177–88
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 63. Synthesis of eight-membered-ring metallacycles: *X*-ray crystal structures of $[Pt_4W_4(\mu\text{-}CR)(\mu_3\text{-}CR)_3(\mu\text{-}CO)(CO)_7(\eta\text{-}C_5H_5)_4]·CH_2Cl_2$, $[Ni_2Pt_2W_4(\mu\text{-}CR)(\mu_3\text{-}CR)_3(\mu\text{-}CO)(CO)_7(\eta\text{-}C_5H_5)_4]·CH_2Cl_2$ (R = C_6H_4Me-4), and $[Ni_2Pt_2W_4(\mu_3\text{-}CPh)_4(CO)_8(\eta\text{-}C_5H_5)_4]$, 2189–200
Studies involving mixed donor macrocyclic ligands. Comparative *X*-ray diffraction, infrared, and fast atom bombardment mass spectrometric studies of the nickel thiocyanate complexes of three O_2N_3-donor macrocycles, 2537–42
Studies of metal-ion recognition. The interaction of Co^{II}, Ni^{II}, and Cu^{II} with new oxygen–nitrogen donor macrocycles; *X*-ray structures of complexes of Cu^{II} and Ni^{II} with a 15-membered O_2N_3 derivative, 2543–54
Complexes of cobalt(II) and nickel(II) with 3- and 4-acetylpyridine, 2595–8
Synthetic and equilibrium studies on copper(II), nickel(II), and palladium(II) complexes of *N*,*N*′,*N*″,*N*‴-tetrakis(2-hydroxyethyl)-1,4,8,11-tetra-azacyclotetradecane (L); kinetics of the acid dissociation of the copper(II) complex and *X*-ray structural characterisation of the nickel(II) complex $[NiLH_{-1}][ClO_4]$, 2605–14
Bicyclam [6,6′-bi(1,4,8,11-tetra-azacyclotetradecane)]: a ditopic receptor for homo- and hetero-bimetallic complexes, 2631–4
Nickel(II), copper(II), and zinc (II) complexes of penta-azamacrocyclic ligands. Crystal and molecular structure of 3,6,9,12-tetramethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-trienezinc(II) perchlorate, 2643–8
Synthesis and study of monomeric and dimeric amidine complexes of nickel(II) and platinum(II). Crystal structure of bis(*N*,*N*′-diphenylbenzamidino)platinum(II), $[Pt\{PhNC(Ph)NPh\}_2]$, 2687–94
Heteronuclear complexes containing Group 6 transition metals; chemical, spectroscopic, and theoretical studies on some binuclear complexes and the *X*-ray crystal structure determination of $[WNi(CO)_3(PPh_3)_2(\eta^5\text{-}C_5H_5)]$, 2741–50
Preparation and structure of nickel(II) complexes of N-substituted tetra-aza macrocycles: molecular structure of [(1*RS*,4*SR*,5*RS*,8*SR*,11*RS*,12*SR*)-1,5,8,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane]nickel(II) diperchlorate, 3093–8

NIOBIUM
The synthesis and structure of Group 5 metal alkyl and alkylidene complexes containing 2,6-dialkylphenoxide ligands: *X*-ray crystal structures of $[Ta(OC_6H_3Me_2\text{-}2,6)_2(CH_2Ph)_3]$, $[Ta(OC_6H_3Me_2\text{-}2,6)_4Me]$, and $[Ta(OC_6H_3Bu^t_2\text{-}2,6)_2(=CHSiMe_3)(CH_2SiMe_3)]$, 155–62
Electronic absorption spectra and photochemical reactivity of Group 5 metal alkyl compounds: photochemical α-hydrogen abstraction, 163–8
Trimethylphosphine complexes of niobium(IV) and tantalum(IV). Crystal, molecular, and electronic structures of $[M_2Cl_4(\mu\text{-}Cl)_4(PMe_3)_4]$ (M = Nb or Ta), 307–14

NIOBIUM(contd)
Determination of the deuterium position in DNb_6I_{11}, 431–4
New heterocyclic compounds containing niobium and molybdenum; crystal structure of a cyclomolybdaphosphazene, 847–50
Synthesis and characterization of monocyclopentadienyl niobium derivatives. *X*-Ray crystal structure of di-μ-chloro-μ-oxo-bis[dichloro(η-trimethylsilylcyclopentadienyl)niobium(v)], $[\{Nb(\eta\text{-}C_5H_4SiMe_3)Cl_2\}_2(\mu\text{-}O)(\mu\text{-}Cl)_2]$, 953–6
New dicyclopentadienyl phosphine, phosphite, and acetylene niobium(III) complexes. Crystal structure of $[Nb(\eta\text{-}C_5H_4SiMe_3)_2Cl(PhC{\equiv}CPh)]$, 975–80

NITRIC OXIDE
Equilibrium and kinetic study of nitric oxide binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 369–72
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 10. The reversible binding of nitric oxide by $[MnX_2(PR_3)]$ (X = Cl, Br, I; R_3 = Pr^n_3, Bu^n_3, $PhMe_2$, $PhEt_2$) in the solid state and in tetrahydrofuran solution. The effect of the nature of the halogen and the phosphine on stability and reversibility, 2497–502

NITRIDE
The radical cations of sulphur ($S_8^{\bullet+}$) and tetrasulphur tetranitride ($S_4N_4^{\bullet+}$): a radiation–electron spin resonance study, 729–32
An electron spin resonance study of the equilibrium between tetrahalogeno-and pentahalogeno-nitridotechnetate(VI) ions in solution, 1721–6

NITRILE
The chemistry of chromium nitrile complexes of 1,2-bis(dimethylphosphino)ethane. *X*-Ray crystal structures of *trans*-$[Cr^{IV}Cl(NEt)(dmpe)_2]CF_3SO_3$, *trans*-$[Cr^{IV}(N{=}CHMe)_2(dmpe)_2][BPh_4]_2$, and *trans*-$[Co^0(H{\cdot}NCMe)(dmpe)_2][BPh_4]_2$, 2947–54

NITROGEN
Studies of phosphazenes. Part 30. Reactions of hexachlorocyclotriphosphazene with aromatic primary amines: interplay of geminal and non-geminal modes of chlorine replacement, 579–84
Studies of phosphazenes. Part 31. Kinetic studies of the reactions of hexachlorocyclotriphosphazene with aromatic primary amines: evidence for the intermediacy of a reactive three-co-ordinated phosphorus(V) species, 585–90
The radical cations of sulphur ($S_8^{\bullet+}$) and tetrasulphur tetranitride ($S_4N_4^{\bullet+}$): a radiation–electron spin resonance study, 729–32
Electron diffraction study of the molecular structure of *N*-methylbis(trimethylsilyl)amine in the gas phase, 785–8
Some investigations of $S_5N_5^+$ compounds; *X*-ray crystal structure of $[S_5N_5]Cl$, 915–24
Structure and basicity. Part 10. Conductometric and potentiometric titrations of some cyclotriphosphazatrienes in non-aqueous media. The conductivities of alkali-metal acetates on titration with perchloric acid in acetic acid, 925–30
Reactions of $(NSCl)_3$ with Lewis acids and sulphur: a convenient preparation of $[S_2N][SbCl_6]$, 1565–8
Phosphorus–nitrogen compounds. Part 51. Carbon-13 and phosphorus-31 nuclear magnetic resonance properties of phenylphosphonothioic di(monoalkylamides). Determination of aromatic substituent constants by ^{13}C nuclear magnetic resonance spectroscopy, 2479–84

NITROGEN-14
A molybdenum-95 and nitrogen-14 nuclear magnetic resonance study of six-co-ordinate hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes containing a sixteen-electron $\{Mo(NO)\}^4$ core, 619–22
Nitrogen-14 nuclear quadrupole resonance study of bonding in transition-metal cyanides, 1637–40

NITROGEN-15
Nitrogen-15 and oxygen-17 nuclear magnetic resonance chemical shifts and ^{15}N-^{29}Si, ^{13}C-^{29}Si, and ^{15}N–^1H coupling constants in cyclosilazoxanes, 1593–6
Preparation and characterisation of mononuclear organoimido complexes of tungsten: *X*-ray crystal structures of $[WCl_4(NC_6H_4Me\text{-}p)(thf)]$, $[P(CH_2Ph)Ph_3][WCl_5(NC_6H_4Me\text{-}p)]$, and $[WCl_2(NPh)_2(bipy)]$, 2067–76

NITROPHENYL
Synthesis of *o*-nitrophenylplatinum(IV) complexes; crystal and molecular structure of dichlorobis(2-nitrophenyl-*C′O*)-platinum(IV), 881–4

NITROSOBENZENE
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\{ON(Ph)C(NPh)S\}(PPh_3)_2]$ and $[Pt\{ON(Ph)C(O)NPh\}(PPh_3)_2]$, 605–8
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of $[Pt\{ON(Ph)C(NPh)S\}(PPh_3)_2]$ and $[Pt\{ON(Ph)C(O)NPh\}(PPh_3)_2]$ (1987, 605), 1583–4

NITROSYL
Preparation and magnetism of the binuclear iron(II) complexes $[\{Fe(C_9H_{21}N_3)X_2\}_2]$ (X = NCS, NCO, or N_3) and their reaction with NO. Crystal structures of $[\{Fe(C_9H_{21}N_3)(NCS)_2\}_2]$ and $[Fe(C_9H_{21}N_3)(NO)(N_3)_2]$, 187–92
Reduction–oxidation properties of organotransition-metal complexes. Part 25. The one-electron reduction of cationic cyclopentadienyl–cobalt and –rhodium nitrosyl derivatives, 467–72
A molybdenum-95 and nitrogen-14 nuclear magnetic resonance study of six-co-ordinate hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes containing a sixteen-electron $\{Mo(NO)\}^4$ core, 619–22
Reactions of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}X_2]$ (M = Mo, X = I; M = W, X = Cl) with some amine-substituted pyridines. The molecular structures of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}(2\text{-}NHC_5NH_4)_2]$, 1063–70
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato-molybdenum and -tungsten nitrosyls. Part 4. The effects of ligating atom type on the reduction potentials of monometallic complexes, 1733–6
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato molybdenum and tungsten nitrosyls. Part 6. Synthesis and electrochemistry of complexes containing saturated heterocyclic amide ligands. *X*-Ray crystal and molecular structure of $[\{Mo(NO)[HB(Me_2pz)_3]Cl\}_2(NC_4H_8N)]$, 2653–60

N.M.R.
Synthesis and properties of heterotrimetallic complexes containing M–Pt–Hg (M = Mo, W, Fe, Mn, or Co) arrays, 119–22
Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of $[5,6,6\text{-}(PPh_3)_3\text{-}6\text{-}H\text{-}$*nido*$\text{-}6\text{-}RuB_9H_{12}]$ (1985, 2397), 275–6
Magnetostructural correlations in parallel square-planar chloride bridged copper(II) dimers: structure, dynamic nuclear magnetic resonance study, and magnetic properties of $[Cu_2(terpy)_2Cl_2][PF_6]_2$, 285–92
trans-Dichlorotetrakis(pyridine)platinum(IV) nitrate: a classical co-ordination compound, 293–6
Preparations, crystal structures, and molybdenum-95 nuclear magnetic resonance spectroscopic studies of triphenylphosphine–gold–tetrathiomolybdate(VI) clusters, 303–6
Tin-119 nuclear magnetic resonance studies of some pseudohalogeno-derivatives of $[SnX_6]^{2-}$ (X = Cl or Br), 315–8
Ruthenium(II) complexes of *closo* dodecaboranyl anions and the molecular structure of the *pileo* thirteen-vertex ruthenaborane $[(PPh_3)_2ClRuB_{12}H_{11}(NEt_3)]$, 451–6
Phosphorus-tellurium compounds. Part 3. Cyclopentadienyliron tellurophosphorane complexes: tellurium-bridged iron phosphine compounds?, 541–4
Studies of phosphazenes. Part 30. Reactions of hexachlorocyclotriphosphazene with aromatic primary amines: interplay of geminal and non-geminal modes of chlorine replacement, 579–84
A molybdenum-95 and nitrogen-14 nuclear magnetic resonance study of six-co-ordinate hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes containing a sixteen-electron $\{Mo(NO)\}^4$ core, 619–22
Bis[bis(diphenylphosphino)methane]platinum(II) salts: evidence for the formation of a five-co-ordinate bis[bis(diphenylphosphino)-methane]halogenoplatinum(II) cation, 705–8
Heteronuclear 1,3-intramolecular metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of 2-thia-4-selenapentane. A dynamic nuclear magnetic resonance investigation, 757–60
Menthyl-substituted organophosphorus compounds. Part 6. Preparation of P(L-men)RCl and P(L-men)R(S)Cl (R = Me, Et,

N.M.R. (contd)
Pr^i Bu^t, or Ph; men = *cyclo*-C_6H_9-2-Pr^i-5-Me) and nuclear magnetic resonance studies of configuration and halogen exchange, 795–806
Proton and phosphorus-31 nuclear magnetic resonance study of zinc(II) *O,O'*-dialkyl dithiophosphates in solution, 807–14
New heterocyclic compounds containing niobium and molybdenum; crystal structure of a cyclomolybdaphosphazene, 847–50
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η^2-vinyl complexes of tungsten, $[W\{\eta^3\text{-}C(CF_3)C(CF_3)(SR)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (R = Me, Et, Pr^n, Pr^i, or Bu^t), 895–906
Lewis-base adducts of Group 11 metal(I) compounds. Part 27. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance, far-infrared, and structural studies on the mononuclear 2:1 adducts of triphenylphosphine with copper(I) and gold(I) halides, 1089–98
Lewis-base adducts of Group 11 metal(I) compounds. Part 28. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance and structural studies on the mononuclear 3:1 adducts of triphenylphosphine with copper(I) halides, 1099–106
The preparation of $[Ru(NCMe)_6][7\text{-}(\eta^6\text{-}C_6Me_6)\text{-}nido\text{-}7\text{-}RuB_{10}H_{13}]_2$ and its characterization by single-crystal *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy: a triply hydrogen-bridged eleven-vertex *nido*-7-metallaundecaborane, 1169–74
Carbon-13 and phosphorus-31 nuclear magnetic resonance studies of the fluxional behaviour of the complexes $[Fe_3(CO)_{10}\{P(OMe)_3\}_2]$ and $[Fe_3(CO)_{10}\{P(OMe)_3\}\{P(OPh)_3\}]$, 1249–52
Fluorine-19 nuclear magnetic resonance studies of tetraethylammonium fluoride in glacial ethanoic acid, 1271–4
Triaquahenicosatungstodiphosphate(6−) heteropolyanion, $[P_2W_{21}O_{71}(OH_2)_3]^{6-}$: *X*-ray crystallographic and ^{183}W nuclear magnetic resonance structural studies (1986, 2237), 1283–4
Pentamethylcyclopentadienylrhodaborane chemistry. Part 2. The reaction of $[6\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}6\text{-}RhB_9H_{13}]$ with dimethylphenylphosphine, and the characterization of $[5\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}5\text{-}RhB_9H_{11}\text{-}7\text{-}(PMe_2Ph)]$, $[2\text{-}(\eta^5\text{-}C_5Me_5)\text{-}closo\text{-}2\text{-}RhB_9H_7\text{-}3,10\text{-}(PMe_2Ph)_2]$, and $[2\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}2\text{-}RhB_8H_{10}\text{-}5\text{-}(PMe_2Ph)]$ by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 1431–44
Phosphorus-31 nuclear magnetic resonance parameters for compounds with the α-tetraphosphorus trisulphide skeleton; substituent constants for an inorganic substrate, 1515–20
The triphenylphosphine cone angle and restricted rotation about the chromium–phosphorus bond in dicarbonyl(η^6-hexaalkylbenzene)(triphenylphosphine)chromium(0) complexes. Crystal and molecular structure of dicarbonyl(η^6-hexa-n-propylbenzene)(triphenylphosphine)chromium(0), 1545–50
Identification of the *endo,exo* isomer of 6,9-$(PMe_2Ph)_2$-*arachno*-$B_{10}H_{12}$ by nuclear magnetic resonance spectroscopy, 1573–6
Nitrogen-15 and oxygen-17 nuclear magnetic resonance chemical shifts and ^{15}N-^{29}Si, ^{13}C-^{29}Si, and ^{15}N–^1H coupling constants in cyclosilazoxanes, 1593–6
Syntheses, *X*-ray structures, and properties of complexes of macrocyclic hexaimines with lanthanide nitrates, 1665–74
Solid-state nuclear magnetic resonance spectroscopic study of γ-zirconium phosphate, 1877–82
Dithio-oxamides as ligands: crystal structures and vibrational analyses of bis(*N,N'*-dicyclohexyldithio-oxamidato)palladium(II), bis(*N,N'*-dibenzyldithio-oxamide)copper(II) diperchlorate, and dichloro(*N,N'*-dimethyldithio-oxamide)zinc(II) and the carbon-13 nuclear magnetic resonance spectra of related compounds, 1921–8
Reactions of rhodium(v) hydrido–silyl complexes with butyl-lithium and with lithium triethylhydridoborate, 1963–6
Phosphorus–nitrogen compounds. Part 50. Further studies on the formation of bicyclic cyclotetraphosphazatetraene derivatives, 2023–30
Studies on borate esters. Part 8. Interactions of cations with oxyacid anion-bridged esters of D-glucarate in alkaline media, 2051–8
Preparation and characterisation of mononuclear organoimido complexes of tungsten: *X*-ray crystal structures of $[WCl_4(NC_6H_4Me\text{-}p)(thf)]$, $[P(CH_2Ph)Ph_3][WCl_5(NC_6H_4Me\text{-}p)]$, and $[WCl_2(NPh)_2(bipy)]$, 2067–76
Oxygen-17 and aluminium-27 nuclear magnetic resonance spectroscopic investigations of aluminium(III) hydrolysis products, 2317–22
Aspects of germanium-73 nuclear magnetic resonance spectroscopy, 2365–72
Synthesis, molecular structure, and nuclear magnetic resonance spectra of the anionic monocarbametallaborane $[NEt_4][(CO)_3(PPh_3)(OH)\text{-}closo\text{-}MoCB_{10}H_{10}]\cdot CH_2Cl_2$, 2385–90
Determination of the relative signs of $^2J(^{31}P\text{–}^{31}P)$ in complexes of tungsten(0) and molybdenum(0) using two-dimensional $[^{31}P\text{–}^{31}P]$-COSY-45 nuclear magnetic resonance chemical shift correlation, 2401–6
Pentamethylcyclopentadienylrhodaborane chemistry. Part 3. The reaction of $[\{Rh(\eta^5\text{-}C_5Me_5)Cl_2\}_2]$ with *closo*-$[B_{10}H_{10}]^{2-}$ and *arachno*-$B_{10}H_{12}$-6,9-$(PMe_2Ph)_2$, and the characterisation of some novel rhodaboranes by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 2417–30
Phosphorus–nitrogen compounds. Part 51. Carbon-13 and phosphorus-31 nuclear magnetic resonance properties of phenylphosphonothioic di(monoalkylamides). Determination of aromatic substituent constants by ^{13}C nuclear magnetic resonance spectroscopy, 2479–84
Preparation, nuclear magnetic resonance spectra, and *X*-ray diffraction study of bis(*N*-benzylacetimidic acid ethyl ester)dichloropalladium(II), 2503–4
Preparation and spectroscopic study of functionally substituted cyclopentadienides of thallium(I), potassium, rhodium(I), and iridium(I), 2615–20
The mechanism of fluxionality of $[Ru(\eta^6\text{-}C_8H_8)(\eta^4\text{-}C_7H_8)]$, $[Ru(\eta^4\text{-}C_8H_8)(\eta^4\text{-}C_7H_8)L]$ [L = PEt_3 or $P(OMe)_3$], and $[Os(\eta^6\text{-}C_8H_8)(\eta^4\text{-}C_8H_{12})]$ (C_8H_8 = cyclo-octatetraene, C_7H_8 = norborna-2,5-diene, and C_8H_{12} = cyclo-octa-1,5-diene), 2649–52
A two-dimensional nuclear magnetic resonance exchange study of the fluxionality of $[PtIMe_3(MeSCH_2CH_2SEt)]$, 2695–702
The formation and characterisation of (η^2-ethene)hydrido(η^5-pentamethylcyclopentadienyl)(trisubstituted-silyl)rhodium complexes; intermediates in catalytic dehydrogenative silylation reactions, 2709–14
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η^2-vinyl complexes of molybdenum and tungsten, $[M\{\eta^3\text{-}C(CF_3)C(CF_3)(L\text{–}L)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (L–L = $MeCO_2^-$, $Me_2PS_2^-$, $Et_2NCS_2^-$, or $Me_2NCS_2^-$), 2715–22
Gold–boron chemistry. Part 1. Synthetic, structural, and spectroscopic studies on the compounds [5,6-μ-(AuPR$_3$)-*nido*-$B_{10}H_{13}$] (R = cyclo-C_6H_{11} or C_6H_4Me-2), 2761–8
Diplatinum-(II) and -(III) complexes with bridging μ-diphosphonito-*P,P'* or μ-methylenebis(phosphonito)-*P,P'* ligands. Synthesis, structure, and nuclear magnetic resonance analysis of the 'lantern' complexes. Crystal structures of $K_4[Pt_2\{\mu\text{-}CH_2[P(O)OH]_2\}_4]\cdot 6H_2O$ and $K_4[Pt_2\{\mu\text{-}CH_2[P(O)OH]_2\}_4Cl_2]\cdot 8H_2O$, 2769–80
Organoruthenaborane chemistry. Part 5. Products of the reaction between *closo*-$[B_{10}H_{10}]^{2-}$ and $[(\eta^6\text{-}C_6Me_6)ClRuB_3H_8]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[\{(\eta^6\text{-}C_6Me_6)_2Ru_2H_4\}RuB_{10}H_8(OEt)_2]$, 2781–8
Ligand intra- and inter-molecular exchange processes in dioxopentakis(trimethylurea)uranium(VI). A proton nuclear magnetic resonance study, 2885–8
Complexes formed between zinc(II) and diethylenetriamine: a carbon-13 nuclear magnetic resonance study, 2889–96
Carbon-13 nuclear magnetic resonance study of the zinc(II) complexes of 3,7-diazanonane-1,9-diamine and 1,2-diaminoethane, 2897–904

N.Q.R.
Nitrogen-14 nuclear quadrupole resonance study of bonding in transition-metal cyanides, 1637–40

NUCLEOTIDE
The configuration and lattice dynamics of complexes of dialkyltin(IV) with adenosine 5'-monophosphate and phenyl phosphates, 789–94
Metal–nucleotide interactions: crystal structures of $Ca[Cu(5'\text{-}GMP)_2(en)(H_2O)_2]\cdot 8H_2O$ (5'-GMP = guanosine 5'-monophosphate) and $Ca[Cu(5'\text{-}IMP)_2(H_2O)_4]\cdot 6.7H_2O$ (5'-IMP = inosine 5'-monophosphate), 2999–3002

OLEFIN
Hydride addition reactions of the olefin complexes $[Fe(\eta^5\text{-}C_5H_5)(CO)_2(olefin)][BF_4]$; formyl formation as the kinetically preferred

OLEFIN (contd)
process; regio- and stereo-selectivity during addition to the co-ordinated olefins, 2857–62
OLIGOMERIZATION
Co-oligomerization of diphenylketene and ethylene promoted by (η^5-cyclopentadienyl)bis(ethylene)cobalt(I): synthesis and *X*-ray structure of a heterocycle containing cobalt(III), 1789–92
OSMIUM
Oxoaryls of rhenium-(V) and -(VI) and osmium(VI). *X*-Ray crystal structures of dimesityldioxorhenium(VI), tetramesityloxorhenium(VI) and dimesityldioxo-osmium(VI), 169–76
Reaction of $[Os_5H_2(CO)_{15}]$ with pyridine: crystal and molecular structures of $[Os_5H_3(CO)_{14}(C_5H_4N)]$ and $[Os_5H_2(CO)_{14}(C_5H_5N)]$, 327–34
Triosmium clusters containing ligands derived from tropone (cycloheptatrienone): molecular structure of an oxidative addition product $[Os_3H(\mu_3\text{-}C_7H_5O)(CO)_9]$, 551–6
The synthesis and *X*-ray crystal structures of homoleptic tetrahedral aryls of osmium(IV) and of cyclohexyls of ruthenium(IV), osmium(IV), and chromium(IV), 557–62
Co-ordination chemistry of higher oxidation states. Part 22. *trans*-Dioxo-osmium(VI) complexes of diamines, diphosphines, and diarsines, 633–8
Trinuclear osmium clusters, containing bidentate $Ph_2PCH_2PPh_2$ and $Me_2PCH_2PMe_2$ ligands: *X*-ray crystal structures of the complexes $[Os_3H(CO)_9(Me_2PCHPMe_2)]$ and $[Os_3H(OH)(CO)_8(Ph_2PCH_2PPh_2)]$, 931–8
Phosphinoalkylsilyl complexes. Part 7. Synthesis and reactivity of octahedral, bis(chelate)dicarbonyl complexes of ruthenium(II) and osmium(II). Conformational properties of the bis(chelate) framework and *X*-ray crystal structures of $[M(PPh_2CH_2CH_2SiMe_2)_2(CO)_2]$ (M = Ru or Os), 969–74
High nuclearity osmium cluster radicals, 1051–4
The preparation, characterisation, and some reactions of $[Os_3H(X)(CO)_9(NMe_3)]$ (X = Cl, Br, or I), 1285–8
Systematic cluster build up by the reaction of $[Os_3H(X)(CO)_9(MeCN)]$ (X = SR or NHMe; R = Me, Et, or Ph) with $[OsH_2(CO)_4]$, 1289–92
Substitution reactions on triosmium clusters of the type $[Os_3(\mu\text{-}H)(\mu\text{-}X)(CO)_{10}]$ [X = NHMe, O_2CR, C_5H_4N, or OC(NHMe); R = H, Me, or Ph], 1293–8
Triosmium clusters containing the ligands CNH_2, NCHCN, and NCMeCN derived from cyanide ion, cyanogen, and acetonitrile, 1457–62
Intermediates in the conversion of $[Os(CO)_{11}(PRPh_2)]$ (R = Me or Ph) into $[Os_3(\mu\text{-}C_6H_4)(\mu_3\text{-}PR)(CO)_9]$: crystal and molecular structures of $[Os_3(\mu\text{-}H)(\mu_3\text{-}C_6H_4PMePh\text{-}o)(CO)_9]$ and $[Os_3(\mu_3\text{-}C_6H_4PMe\text{-}o)(CO)_{10}]$, 1529–34
Substituted osmium clusters containing labile ligands. The preparation, characterisation, and some reactions of $[Os_3H(OR)(CO)_9(MeCN)]$ R = H, Me, Et, or Ph) and the isomers of $[Os_3H(SR)(CO)_9(MeCN)]$ (R = Me, Et, or Ph); the crystal and molecular structure of $[Os_3H(SPh)(CO)_9(PEt_3)]$, 1623–30
Decarboxylation of vinylene carbonate at triosmium clusters; *X*-ray crystal structure of the μ_3-formylmethylidene compound $[Os_3(CHCHO)(CO)_{10}]$, 1861–4
Three isomeric hydride compounds formed by protonation of decacarbonylbis(dimethylphenylphosphine)triosmium and the *X*-ray structures of two of these: 1,1- and 1,2-$[Os_3H(CO)_{10}(PMe_2Ph)_2][PF_6]$, 1869–76
A reversible metal framework rearrangement in high-nuclearity osmium-platinum cluster compounds. *X*-Ray crystal structures of $[Os_6Pt(CO)_{17}(\mu_3\text{-}NCMe)(C_8H_{12})]$, $[Os_6Pt(CO)_{17}(\mu_4\text{-}NCMe)(C_8H_{12})]$ (C_8H_{12} = cyclo-octa-1,5-diene), and $[Os_6(CO)_{19}(\mu_3\text{-}NCMe)]$: compounds with novel metal–acetonitrile bonding modes, 2245–52
Synthesis, reactivity, and *X*-ray crystal structures of $[Os_6Pt_2H_2(CO)_{16}(\mu\text{-}C_8H_{10})(C_6H_{12})]\cdot CH_2Cl_2$ and $[Os_6H(CO)_{16}(\mu\text{-}C_8H_{11})]$. Model compounds for C–H activation, 2253–60
Attempted synthesis of osmium(VI) and iridium(VI) thiofluorides; the preparation of $OsF_5\cdot SF_4$ and $IrF_5\cdot SF_4$, 2285–8
The cyclometallation of benzoic acid to give rhodium, iridium, and osmium *C,O*-benzoates. *X*-Ray structure determination of the dibenzoate $[(C_5Me_5)Rh(OOCPh)_2(H_2O)]$, 2459–66
Substitution reactions of the osmium(I) dinuclear compound $[Os_2(MeCO_2)_2(CO)_6]$. *X*-Ray crystal structure of $[N(PPh_3)_2][Os_2(MeCO_2)_2Cl(CO)_5]$, 2473–8
The mechanism of fluxionality of $[Ru(\eta^6\text{-}C_8H_8)(\eta^4\text{-}C_7H_8)]$, $[Ru(\eta^4\text{-}C_8H_8)(\eta^4\text{-}C_7H_8)L]$ [L = PEt_3 or $P(OMe)_3$], and $[Os(\eta^6\text{-}C_8H_8)(\eta^4\text{-}C_8H_{12})]$ (C_8H_8 = cyclo-octatetraene, C_7H_8 = norborna-2,5-diene, and C_8H_{12} = cyclo-octa-1,5-diene), 2649–52
The syntheses and crystal and molecular structures of some ten-vertex (MCB_8; M = Ru or Os) metallacarbaboranes: 9,9,9-$(CO)(PPh_3)_2$-*nido*-9,6-MCB_8H_{10}-5-PPh_3)] (M = Ru or Os) and [9,9,9-$(\eta\text{-}C_5H_5)(H)(PPh_3)$-*arachno*-9,6-$RuCB_8H_{12}$], 2793–802
Synthesis using metal atoms of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[OsH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[MnH(\eta\text{-}C_6H_6)\text{-}(PMe_3)_2]$, and $[Mn(\eta\text{-}C_5H_5)(PMe_3)_3]$: arene ligand-exchange reactions of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, 2823–30
Reactions of allene with triosmium and triruthenium clusters including coupling and oxidative addition: *X*-ray structure of the coupled allene compound $[Os_3(C_6H_8)(CO)_{10}]$, 2935–40
Preparation of two mixed-metal carbonyl clusters based on the platinumtriosmium core. *X*-Ray crystal structure of $[Os_3PtH_2(cod)(CH_2)(CO)_9]$ (cod = cyclo-octa-1,5-diene), 3103–6
OXADIAZAGERMINE
Formation of germanium–nitrogen double bonds in reactions of the electron-rich germylene bis[bis(trimethylsilyl)amido]-germanium(II) with a range of diazo-compounds, 501–8
OXADIAZASTANNINE
Reaction of an enolisable diazo-compound with the electron-rich stannylene $Sn[N(SiMe_3)_2]_2$: preparation of a novel spiro[1,3,4,2-oxadiazastannine-6,4′-pyrazole] derivative, 509–12
OXALATE
Preparation and crystal structures of bis(tetraphenylarsonium) tris(oxalato)technetate(IV), and tetraphenylarsonium tris(benzene-1,2-dithiolato)technetate(V): octahedral *versus* trigonal-prismatic geometry for tris-bidentate complexes of technetium, 2305–10
OXIDATION
Photo-oxidations of acetoin and biacetyl catalysed by tetrabutylammonium decatungstate(4−) in acetonitrile under excess of oxygen, 961–2
OXIDE
Magnetic and electrochemical studies on the series $[Cr_nFe_{3-n}(\mu_3\text{-}O)(gly)_6(H_2O)_3][NO_3]_7\cdot xH_2O$(gly = glycine; *n* = 0, 1, 2, or 3) and on the mixed-valence complex $[Fe^{II}Fe^{III}_2(\mu_3\text{-}O)(gly)_6\text{-}(H_2O)_3]Cl_6$, 97–100
Oxoaryls of rhenium-(V) and -(VI) and osmium(VI). *X*-Ray crystal structures of dimesityldioxorhenium(VI), tetramesityloxorhenium(VI) and dimesityldioxo-osmium(VI), 169–76
Some binuclear molybdenum(VI) complexes obtained from condensation reactions: their crystal and molecular structures, 523–8
Intercalation compounds of (diphosphato)oxotungsten(VI), $[WO(P_2O_7)]$, 609–12
Co-ordination chemistry of higher oxidation states. Part 22. *trans*-Dioxo-osmium(VI) complexes of diamines, diphosphines, and diarsines, 633–8
Magnetic exchange interactions in perovskite solid solutions. Part 7. A study of the non-stoicheiometric phase $SrCoO_{3-y}$ by ^{57}Fe Mössbauer spectroscopy, 667–72
A comparative synthetic and structural study of μ-oxo and μ-sulphido derivatives of *N,N′*-ethylenebis(acetylacetoneiminato)-iron(III), including the synthesis and structural characterization of the parent compound, 709–14
Synthesis and characterization of monocyclopentadienyl niobium derivatives. *X*-Ray crystal structure of di-μ-chloro-μ-oxo-bis[dichloro(η-trimethylsilylcyclopentadienyl)niobium(V)], $[\{Nb(\eta\text{-}C_5H_4SiMe_3)Cl_2\}_2(\mu\text{-}O)(\mu\text{-}Cl)_2]$, 953–6
Studies on transition-metal peroxo complexes. Part 7. Molybdenum(VI) and tungsten(VI) carboxylato peroxo complexes, and the *X*-ray crystal structure of $K_2[MoO(O_2)_2(glyc)]\cdot 2H_2O$, 991–6
Synthesis, characterization, and reactions of chromyl(VI) alkoxides, 1253–6
Magnetic exchange interactions in perovskite solid solutions. Part 8. A study of the solid solution $SrFe_{1-x}Co_xO_{3-y}$ by iron-57 Mössbauer spectroscopy, 1419–24
Preparation and partial oxidation of tin(II) trifluoroacetate: spectroscopic properties and *X*-ray crystal structure of di-μ_3-oxo-octakis-μ-trifluoroacetato-tetratin(II)tin(IV), 1675–82

OXIDE (contd)
Technetium(v) and rhenium(v) complexes with *N*-(2-mercaptophenyl)salicylideneiminate. Crystal structure of chloro(oxo)[*N*-(2-sulphidophenyl)salicylideneiminato-*NOS*]technetium(v), 1693–700
Iron antimonate, 1727–32
Synthesis and characterisation of some new manganese(II) and manganese(III) complexes of tetradentate Schiff-base ligands and their reaction with molecular oxygen, 2009–12
Analysis and prediction of acid–base reactions between oxides and oxysalts using the optical basicity concept, 2323–8
A structural comparison of the two polymorphs of $Sr_2Co_2O_5$ by cobalt *K*-edge extended *X*-ray absorption fine structure spectroscopy, 2359–64
Reactions in mixed non-aqueous solution containing sulphur dioxide. Part 9. Mechanisms of dissolution of metals, oxides, and sulphites, 2995–8
Surface characterisation of α-, β-, γ-, and δ-UO_3 using *X*-ray photoelectron spectroscopy, 3009–16
OXIME
Synthesis of *SS'*-bis(4'-benzo[15-crown-5])dithioglyoxime and its complexes with copper(II), nickel(II), cobalt(II), cobalt(III), palladium(II), platinum(II), and platinum(IV), 1827–32
OXYGEN-17
Nitrogen-15 and oxygen-17 nuclear magnetic resonance chemical shifts and ^{15}N-^{29}Si, ^{13}C-^{29}Si, and ^{15}N–^{1}H coupling constants in cyclosilazoxanes, 1593–6
Oxygen-17 and aluminium-27 nuclear magnetic resonance spectroscopic investigations of aluminium(III) hydrolysis products, 2317–22
PALLADIUM
Bis(dialkylphosphino)methane complexes of palladium(II) and the molecular structure of $[Pd_2Cl_3(\mu\text{-}Pr^i_2PCH_2PPr^i_2)(\mu\text{-}Pr^i_2PCH_2PHPr^i)]\cdot 2CHCl_3$ containing an unusual phosphidophosphine bridging ligand, 201–6
Metallacyclic complexes. Synthesis of metallathietane-3,3-dioxide (metallacyclosulphone) complexes of platinum(II), palladium(II), and nickel(II); crystal structure of *trans*-2,4-diphenyl-1,1-bis(triphenylarsine)platinathietane-3,3-dioxide, 733–40
Reactions of cyclobutenedione and cyclopentenetrione with zero-valent nickel triad complexes of isocyanide. *X*-Ray crystal structures of (3—4-η-3,4-diphenylcyclobut-3-ene-1,2-dione)- and (4—5-η-4,5-diphenylcyclopent-4-ene-1,2,3-trione)-tris(t-butylisocyanide)nickel, 853–8
Rational synthesis of metalla–sulphur–nitrogen complexes: use of $[Me_2SnS_2N_2]_2$ in the preparation of complexes of the type $[M(S_2N_2H)(PR_3)_2]X$ (M = Pt or Pd). *X*-Ray structures of $[Pt(S_2N_2H)(PEt_3)_2][Me_2SnCl_3]$, $[Pt(S_2N_2H)(PBu^n_3)_2]_2[PF_6]Cl$, and $[\{Co(S_2N_2H)_2\}_2\{Ph_2P(O)CH_2CH_2P(O)Ph_2\}]$, 907–14
Reactions of planar pentadentate ligands with palladium and platinum chlorides. Crystal structures of *cis*-dichlorobis(*N'*-isopropylidenebenzohydrazide)platinum(II) and the cyclopalladated dianion of *N',N'''*-(pyridine-2,6-diyldiethylidyne)bis(benzohydrazide), 1175–80
Syntheses and characterization of dinuclear palladium and platinum complexes with a single unsupported bridging group. Crystal structure of $[\{C_6H_3(CH_2NMe_2)_2\text{-}o,o'\}Pd(\mu\text{-}Cl)Pd\{C_6H_3(CH_2NMe_2)_2\text{-}o,o'\}]BF_4$, 1359–66
Synthesis and characterization of palladium(II)–η^3-allyl–ylide complexes. *X*-Ray crystal structure of $[PdCl(\eta^3\text{-}2\text{-}MeC_3H_4)\{Ph_3PC(H)COMe\}]$, 1381–8
Cyclopalladation to form planar tridentate $[N\text{–}C\text{–}N]^-$ intramolecular co-ordination systems involving pyridine donor groups, including ligand synthesis and *X*-ray structural studies., 1477–84
Bimetallic systems. Part 21. Synthesis of vinylidene- and phenylene-bridged complexes of dinickel(II), nickel(II)–palladium(II), and nickel(II)–platinum(II) containing $Ph_2PCH_2PPh_2$ (dppm): crystal structures of $[(SCN)Ni(\mu\text{-}dppm)_2(\mu\text{-}C{=}CH_2)Ni(NCS)]$ and $[ClPd(\mu\text{-}dppm)_2(\mu\text{-}C{=}CCl_2)NiCl]$, 1501–8
Raman and resonance-Raman studies of platinum group tetrathiomolybdates and tetrathiotungstates, 1535–44
Neutral and anionic 2-methyl-6-nitrophenyl palladium(II) complexes. The *X*-ray crystal structure of bis(benzyltriphenylphosphonium) *trans*-di-μ-chloro-bis[chloro(2-methyl-6-nitrophenyl)palladate(II)]–acetone (1/2), 1655–8
Dithio-oxamides as ligands: crystal structures and vibrational analyses of bis(*N,N'*-dicyclohexyldithio-oxamidato)palladium(II), bis(*N,N'*-dibenzyldithio-oxamide)copper(II) diperchlorate, and dichloro(*N,N'*-dimethyldithio-oxamide)zinc(II) and the carbon-13 nuclear magnetic resonance spectra of related compounds, 1921–8
The synthesis and reactions of palladium–iron carbonyl complexes containing bridging $Ph_2PCH_2PPh_2$ ligands, 2005–8
Pentafluorophenyl complexes of palladium and platinum containing chelating, unidentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 2077–82
Preparation, nuclear magnetic resonance spectra, and *X*-ray diffraction study of bis(*N*-benzylacetimidic acid ethyl ester)dichloropalladium(II), 2503–4
Synthetic and equilibrium studies on copper(II), nickel(II), and palladium(II) complexes of *N,N',N'',N'''*-tetrakis(2-hydroxyethyl)-1,4,8,11-tetra-azacyclotetradecane (L); kinetics of the acid dissociation of the copper(II) complex and *X*-ray structural characterisation of the nickel(II) complex $[NiLH_{-1}][ClO_4]$, 2605–14
Mixed-metal phosphinito complexes of platinum(II) and palladium(II) with lanthanide and actinide elements. The single-crystal *X*-ray structure determination of $[UO_2(OH_2)\{(OPPh_2)_2Pd(S_2CNEt_2)\}_2]$, 2853–6
Platinum(II)– and palladium(II)–iron(0) complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand. Crystal structures of $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}CO)PtBr_2]$ and $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}I)PtI]\cdot CH_2Cl_2$, 3079–84
PENTA-AZABICYCLO-OCTADECATRIENE
Nickel(II), copper(II), and zinc (II) complexes of penta-azamacrocyclic ligands. Crystal and molecular structure of 3,6,9,12-tetramethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-trienezinc(II) perchlorate, 2643–8
PENTAFLUOROPHENYL
Pentafluorophenyl complexes of palladium and platinum containing chelating, unidentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 2077–82
Mono- and bi-nuclear gold(I) and gold(III) complexes with $S_2C\text{–}PR_3$ ligands. *X*-Ray crystal structures of $[Au(C_6F_5)_3(S_2C\text{–}PEt_3)]$ and $[Au_2(\mu\text{-}S_2CPEt_3)(C_6F_5)_6]\cdot CH_2Cl_2$, 3017–22
PENTAMETHYLCYCLOPENTADIENYL
Binuclear rhodium and iridium complexes containing pentamethylcyclopentadienyl and pyrazolate ligands, 639–46
Reaction of elemental sulphur with bis(pentamethylcyclopentadienyl)vanadium derivatives: disulphido and μ-sulphido complexes. Crystal structures of $[V(\eta\text{-}C_5Me_5)_2(S_2)]$ and $[V_2(\eta\text{-}C_5Me_5)_2Cl_2(\mu\text{-}S)_2]$, 2099–104
PENTAOXACYCLOPENTADECANE
Stereochemical role of lone pairs in main-group elements. Part 3. Structure and bonding in trichloro(1,4,7,10,13-pentaoxacyclopentadecane)antimony(III) studied by means of *X*-ray crystallography at 120 K, 427–30
PEPTIDE
Co-ordination abilities of thioamide analogues of amino acids and peptides. Copper(II) and nickel(II) complexes, 1–4
A thermodynamic and spectroscopic study of the proton and copper(II) complexes of L-prolyl-L-histidine, D-prolyl-L-histidine, L-histidyl-L-histidine, and D-histidyl-L-histidine, 661–6
Copper(II) complexes of tyrosine-containing tripeptides, 1263–6
Studies on transition-metal–peptide complexes. Part 13. Copper(II) and nickel(II) complexes of amino acids and peptides containing a thioether group, 1717–20
PERCHLORATE
A quantitative approach to host–guest interactions for matrix-isolated alkali-metal salts of hexafluoroanions and perchlorates, 1521–8
PEROVSKITE
Magnetic exchange interactions in perovskite solid solutions. Part 7. A study of the non-stoicheiometric phase $SrCoO_{3-y}$ by ^{57}Fe Mössbauer spectroscopy, 667–72
Magnetic exchange interactions in perovskite solid solutions. Part 8. A study of the solid solution $SrFe_{1-x}Co_xO_{3-y}$ by iron-57 Mössbauer spectroscopy, 1419–24
PEROXIDASE
A realistic model for heme-containing catalases and peroxidases: the *X*-ray structural characterisation of a non-porphyrin iron(III) macrocyclic complex, and the mechanism of its peroxidation of aromatic substrates, 2505–10
PEROXIDATION
A realistic model for heme-containing catalases and peroxidases: the

PEROXIDATION (contd)
X-ray structural characterisation of a non-porphyrin iron(III) macrocyclic complex, and the mechanism of its peroxidation of aromatic substrates, 2505–10
PEROXIDE
Studies on transition-metal peroxo complexes. Part 7. Molybdenum(VI) and tungsten(VI) carboxylato peroxo complexes, and the *X*-ray crystal structure of $K_2[MoO(O_2)_2(glyc)]\cdot 2H_2O$, 991–6
PHASE
Analysis and prediction of acid–base reactions between oxides and oxysalts using the optical basicity concept, 2323–8
PHASE TRANSITION
Thermally induced phase transition and *X*-ray crystal structure of bis(*N,N'*-dimethyl-1,2-ethanediamine)di-isothiocyanatonickel(II), 997–1000
PHENANTHROLINE
Metal ion-bridged intramolecular stacking interaction between the tryptophyl residue and the aromatic heterocyclic amine within the ternary complex (1,10-phenanthroline)(L-tryptophanato-*O,N*)copper(II) perchlorate–water (1/2.5): an *X*-ray study, 2017–22
Crystal structures and properties of $[Au(phen)\{(CN)_{0.92}Br_{0.08}\}_2]Br$ and $[Au(phen)(CN)\{(CN)_{0.82}Br_{0.18}\}]\cdot 0.5$*trans*-$[Au(CN)_2Br_2]\cdot 0.5Br\cdot phen$ (phen = 1,10-phenanthroline) obtained by disproportionation of five-co-ordinate bromodicyano(1,10-phenanthroline)gold(III). Two examples of secondary co-ordination and CN/Br disorder in square-planar gold(III) complexes, 2235–40
Preparation, crystal and molecular structure, and antimony-121 Mössbauer spectrum of trifluoro(1,10-phenanthroline)-(thiourea)antimony(III), 2451–4
PHENOLATE
Structure and magnetism of analogous *O*-bridged (phenolato) and *S*-bridged (thiophenolato) binuclear copper(II) complexes. Crystal structure of the binuclear *S*-bridged complex μ-[2,6-bis(4′-cyclohexyl-4′-hydroxy-2′,3′-diazabuta-1′,3′-dien-1′-yl)-4-methylthiophenolato(3−)-*S,N*^{2},*O*(Cu^{1,2})]-methanol-μ-(pyrazolyl-*N,N'*)-dicopper(II), 1585–92
An electron spin resonance study of the copper(II) chelate of 2-(2′-pyridylmethylenehydrazonomethyl)phenol and related compounds, 1755–62
An electron spin resonance study of the dioxygen adduct of bis[2-(2′-pyridylmethylenehydrazonomethyl)phenolato]cobalt(II) and related materials, 2341–6
PHENOXIDE
The synthesis and structure of Group 5 metal alkyl and alkylidene complexes containing 2,6-dialkylphenoxide ligands: *X*-ray crystal structures of $[Ta(OC_6H_3Me_2\text{-}2,6)_2(CH_2Ph)_3]$, $[Ta(OC_6H_3Me_2\text{-}2,6)_4Me]$, and $[Ta(OC_6H_3Bu^t_2\text{-}2,6)_2(=CHSiMe_3)(CH_2SiMe_3)]$, 155–62
PHENYL
Phenyl complexes of ruthenium(II): competition between carbonyl substitution and formation of benzoyl complexes, and an *X*-ray study of a benzoyl complex, 27–34
Methyl- and phenyl-mercury(II) derivatives of 2-mercaptopyridine. Crystal and molecular structure of methyl(pyridine-2-thiolato)-mercury(II) (1986, 1945), 683–4
Syntheses and characterization of dinuclear palladium and platinum complexes with a single unsupported bridging group. Crystal structure of $[\{C_6H_3(CH_2NMe_2)_2\text{-}o,o'\}Pd(\mu\text{-}Cl)Pd\{C_6H_3(CH_2NMe_2)_2\text{-}o,o'\}]BF_4$, 1359–66
Cyclopalladation to form planar tridentate [N–C–N]$^-$ intramolecular co-ordination systems involving pyridine donor groups, including ligand synthesis and *X*-ray structural studies., 1477–84
PHENYLENE
Bimetallic systems. Part 21. Synthesis of vinylidene- and phenylene-bridged complexes of dinickel(II), nickel(II)–palladium(II), and nickel(II)–platinum(II) containing $Ph_2PCH_2PPh_2$ (dppm): crystal structures of $[(SCN)Ni(\mu\text{-}dppm)_2(\mu\text{-}C{=}CH_2)Ni(NCS)]$ and $[ClPd(\mu\text{-}dppm)_2(\mu\text{-}C{=}CCl_2)NiCl]$, 1501–8
PHOSPHATE
The configuration and lattice dynamics of complexes of dialkyltin(IV) with adenosine 5′-monophosphate and phenyl phosphates, 789–94
Solid-state nuclear magnetic resonance spectroscopic study of γ-zirconium phosphate, 1877–82
Lamellar inorganic ion exchangers. Proton–copper(II) ion exchange in γ-titanium bis(hydrogenphosphate), 2045–50
PHOSPHAZENE
Phosphorus–nitrogen compounds. Part 50. Further studies on the formation of bicyclic cyclotetraphosphazatetraene derivatives, 2023–30
PHOSPHIDE
Bis(dialkylphosphino)methane complexes of palladium(II) and the molecular structure of $[Pd_2Cl_3(\mu\text{-}Pr^i_2PCH_2PPr^i_2)(\mu\text{-}Pr^i_2PCH_2PHPr^i)]\cdot 2CHCl_3$ containing an unusual phosphidophosphine bridging ligand, 201–6
Intermediates in the conversion of $[Os(CO)_{11}(PRPh_2)]$ (R = Me or Ph) into $[Os_3(\mu_3\text{-}C_6H_4)(\mu_3\text{-}PR)(CO)_9]$: crystal and molecular structures of $[Os_3(\mu\text{-}H)(\mu_3\text{-}C_6H_4PMePh\text{-}o)(CO)_9]$ and $[Os_3(\mu_3\text{-}C_6H_4PMe\text{-}o)(CO)_{10}]$, 1529–34
Synthesis and substitution reactions of a heterodimetallic molybdenum–manganese complex, $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_6]$; *X*-ray crystal structure of $[MoMn(\mu\text{-}H)(\mu\text{-}PPh_2)(\eta^5\text{-}C_5H_5)(CO)_4(dppm\text{-}PP')]$, 1557–64
The reaction of $[Mn_2\{\sigma{:}\eta^2\text{-}C(R){=}CHR\}(\mu\text{-}PPh_2)(CO)_7]$ (R = H or Ph) with nucleophiles: *X*-ray crystal structures of $[N(PPh_3)_2][Mn_2(\mu\text{-}H)\{\mu\text{-}C(O)C(Ph){=}CHPh\}(\mu\text{-}PPh_2)(CO)_6]$, $[Mn_2\{PPh_2C(Ph){=}CHPh\}(CO)_9]$, $[Mn_2(\mu\text{-}PPh_2\text{-}CH{=}CH_2)(CO)_8]$, $[Mn_2\{\mu\text{-}C(O)C(Ph){=}CHPh\}(\mu\text{-}PPh_2)(CO)_6(CNBu^t)_2]$, $[Mn_2\{\mu\text{-}\sigma{:}\eta^2\text{-}C(Ph){=}CHPh\}(\mu\text{-}PPh_2)(CO)_6(CNBu^t)]$, and $[Mn(\mu\text{-}PPh_2CH{=}CH_2)(CO)_7(PEt_3)]$, 2669–82
PHOSPHINE
Synthesis of $[Mo(\eta\text{-}C_5H_4Pr^i)(PR_3)_2H_3]$($R_3$ = Me_3 or Me_2Ph), $[Mo(\eta\text{-}C_5H_4Pr^i)(Pr^i_2PCH_2CH_2PPr^i_2)H_3]$, and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of $[\{Mo(\eta\text{-}C_5H_4Pr^i)(\mu\text{-}Cl)_2\}_2]$, 9–20
Synthesis and properties of heterotrimetallic complexes containing M–Pt–Hg (M = Mo, W, Fe, Mn, or Co) arrays, 119–22
Synthesis and *X*-ray crystal structure of a cuboctahedral platinum cluster, 259–62
Reactions between bis(*O*-alkyl dithiocarbonato)nickel(II) complexes and phosphines. Formation of a dithiocarbonate complex of nickel(II): $[Ni(S_2CO)(Ph_2PCH_2CH_2PPh_2)]$, 281–4
Trimethylphosphine complexes of niobium(IV) and tantalum(IV). Crystal, molecular, and electronic structures of $[M_2Cl_4(\mu\text{-}Cl)_4(PMe_3)_4]$ (M = Nb or Ta), 307–14
Preparation of isocyanide and mixed dinitrogen–isocyanide complexes of rhenium(I) from reactions of *trans*-$[ReCl(N_2)(PMe_2Ph)_4]$, *mer*-$[Re(S_2PPh_2)(N_2)(PMe_2Ph)_3]$, or *mer*-$[Re(S_2CNEt_2)(N_2)(PMe_2Ph)_3]$ with methyl isocyanide; crystal structure of *mer*-$[Re(S_2PPh_2)(N_2)(CNMe)(PMe_2Ph)_3]$, 319–26
Chemistry of platinum sulphido-complexes. Part 6. Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with the rhodium complexes $[(RhL_2Cl)_2]$ (L = CO or C_2H_4) and $[Rh(2,6\text{-}Me_2C_6H_3NC)_4]PF_6$. Crystal and molecular structures of $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(CO)_2][Rh(CO)_2Cl_2]\cdot C_3H_6O$ and $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(2,6\text{-}Me_2C_6H_3NC)_2]PF_6$, 335–40
Thermal and photochemical behaviour of organotetraruthenium clusters: solution structures and dynamics of phosphine-substituted derivatives, 349–54
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au), 379–90
η-Benzenebis(trimethylphosphine)iron as a precursor to $[Fe(\eta\text{-}C_5R_5)(PMe_3)_2]$ (R = H or Me) derivatives: the equilibrium $[Fe](PMe_3)Et \rightleftharpoons [Fe](\eta\text{-}C_2H_4)H + PMe_3$, where $[Fe] = Fe(\eta\text{-}C_5Me_5)(PMe_3)$, 411–6
Reactivity of the polyhydride complexes $[ReH_5(PPh_3)_3]$, $[ReH_3(dppe)_2]$ (dppe = $Ph_2PCH_2CH_2PPh_2$), $[ReH_3(PPh_3)_3L]$, and $[ReH_4(PPh_3)_3L]PF_6$ (L = MeCN or Bu^{t}NC) towards electrophiles and nucleophiles, 715–20
Menthyl-substituted organophosphorus compounds. Part 6. Preparation of P(L-men)RCl and P(L-men)R(S)Cl (R = Me, Et, Pr^i Bu^t, or Ph; men = *cyclo*-C_6H_9-2-Pr^i-5-Me) and nuclear magnetic resonance studies of configuration and halogen exchange, 795–806
Redox behaviour of the iron–sulphur cluster $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{2+}$. Synthesis and crystal structure of the new paramagnetic monopositive species $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^+$ as its $[PF_6]^-$ salt, 831–6
Reactions of aluminopolyhydride complexes of tungsten. *X*-Ray

PHOSPHINE (contd)
crystal structures of $[(Me_3P)_3H_3W(\mu\text{-}H)_2AlCl(\mu\text{-}N{=}CHEt)]_2$, $\{[(Me_3P)_3WH_4]Li\}_4$, $WH_2I(PMe_3)_3(SiMe_3)$, and $(Me_3P)_3H_2W(\mu\text{-}H)_3AlCl_2(NMe_3)$, 837–46
Rational synthesis of metalla–sulphur–nitrogen complexes: use of $[Me_2SnS_2N_2]_2$ in the preparation of complexes of the type $[M(S_2N_2H)(PR_3)_2]X$ (M = Pt or Pd). *X*-Ray structures of $[Pt(S_2N_2H)(PEt_3)_2][Me_2SnCl_3]$, $[Pt(S_2N_2H)(PBu^n{}_3)_2]_2[PF_6]Cl$, and $[\{Co(S_2N_2H)_2\}_2\{Ph_2P(O)CH_2CH_2P(O)Ph_2\}]$, 907–14
Complexes of the platinum metals. Part 32. Synthesis and chemistry of some iridium sulphonate derivatives, 947–52
New dicyclopentadienyl phosphine, phosphite, and acetylene niobium(III) complexes. Crystal structure of $[Nb(\eta\text{-}C_5H_4SiMe_3)_2Cl(PhC{\equiv}CPh)]$, 975–80
Lewis-base adducts of Group 11 metal(I) compounds. Part 27. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance, far-infrared, and structural studies on the mononuclear 2:1 adducts of triphenylphosphine with copper(I) and gold(I) halides, 1089–98
Lewis-base adducts of Group 11 metal(I) compounds. Part 28. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance and structural studies on the mononuclear 3:1 adducts of triphenylphosphine with copper(I) halides, 1099–106
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 61. Reactions of iron–molybdenum compounds with tertiary phosphines, 1243–8
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au) (1987, 379), 1283–4
The synthesis of silica-supported phosphine and phosphine oxide complexes of manganese(II) iodide and their reaction with sulphur dioxide, 1415–8
Intermediates in the conversion of $[Os(CO)_{11}(PRPh_2)]$ (R = Me or Ph) into $[Os_3(\mu_3\text{-}C_6H_4)(\mu_3\text{-}PR)(CO)_9]$: crystal and molecular structures of $[Os_3(\mu\text{-}H)(\mu_3\text{-}C_6H_4PMePh\text{-}o)(CO)_9]$ and $[Os_3(\mu_3\text{-}C_6H_4PMe\text{-}o)(CO)_{10}]$, 1529–34
The triphenylphosphine cone angle and restricted rotation about the chromium–phosphorus bond in dicarbonyl(η^6-hexaalkylbenzene)(triphenylphosphine)chromium(0) complexes. Crystal and molecular structure of dicarbonyl(η^6-hexa-n-propylbenzene)(triphenylphosphine)chromium(0), 1545–50
Comments on the attempts by Wickens and Abrams to synthesize manganese(II) phosphine complexes: a reply, 1581–2
Synthesis of (η-cyclopentadienyl)(trimethylphosphine)tungsten derivatives and related chemistry, 1611–8
The co-ordination of small molecules by manganese(II) phosphine complexes. The reaction of some $[MnX_2L]$ (X = Cl, Br, or I; L = $PPr^n{}_3$, PMe_2Ph, or PPh_3) complexes with tetracyanoethylene, 1785–8
Three isomeric hydride compounds formed by protonation of decacarbonylbis(dimethylphenylphosphine)triosmium and the *X*-ray structures of two of these: 1,1- and 1,2-$[Os_3H(CO)_{10}(PMe_2Ph)_2][PF_6]$, 1869–76
A general synthetic route for platinum cluster compounds containing carbonyl and tertiary phosphine ligands and a study of their reactions with unsaturated inorganic molecules, 1889–96
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 8. Irreversible oxidation of trimethylphosphinemanganese(II) dihalides, $[MnX_2(PMe_3)]$ (X = Cl, Br, or I), to bis(trimethylphosphine)manganese(III) trihalides, $[MnX_3(PMe_3)_2]$. *X*-Ray crystal structure of *trans*-triiodobis(trimethylphosphine)manganese(III), the first example of a trigonal-bipyramidal manganese(III) tertiary phosphine complex, 1999–2004
(η-Benzene)bis(trimethylphosphine)iron: a useful precursor to polyene iron derivatives. Crystal structure of 2—4-η:2′— 4′-η-1,1′-bi(cyclohex-3-en-2-yl)bis(trimethylphosphine)iron(II), 2031–8
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 9. The reaction of ethylene with $[MnX_2L]$ (X = Cl, Br, or I; L = $PPr^n{}_3$, $PBu^n{}_3$, $PPhMe_2$, $PPhEt_2$, or PPh_3) in tetrahydrofuran solution, 2241–4
Synthesis of chelate stabilised alcohol- and alkoxo–platinum(II) complexes. *X*-Ray crystal structure of $[Pt(PPh_2CH_2CMe_2O)_2]\cdot 3.5H_2O$, 2273–80
Gold-197 Mössbauer spectra of some gold–ruthenium cluster compounds, 2333–6
Ab initio study of stable bis(carbon dioxide)molybdenum complexes, 2373–8
Determination of the relative signs of $^2J(^{31}P\text{–}^{31}P)$ in complexes of tungsten(0) and molybdenum(0) using two-dimensional $[^{31}P\text{–}^{31}P]$-COSY-45 nuclear magnetic resonance chemical shift correlation, 2401–6
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 10. The reversible binding of nitric oxide by $[MnX_2(PR_3)]$ (X = Cl, Br, I; R_3 = $Pr^n{}_3$, $Bu^n{}_3$, $PhMe_2$, $PhEt_2$) in the solid state and in tetrahydrofuran solution. The effect of the nature of the halogen and the phosphine on stability and reversibility, 2497–502
Synthesis using metal atoms of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[OsH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, $[MnH(\eta\text{-}C_6H_6)\text{-}(PMe_3)_2]$, and $[Mn(\eta\text{-}C_5H_5)(PMe_3)_3]$: arene ligand-exchange reactions of $[RuH(\eta\text{-}C_6H_6)(PMe_3)(\sigma\text{-}C_6H_5)]$, 2823–30
Reactions of phosphine, arsine, and stibine with carbonylbis(triethylphosphine)iridium(I) halides. Part 1. Reactions in toluene; *X*-ray crystal structures of $[Ir(CO)ClH(PEt_3)_2(AsH_2)]$ and $[Ir(CO)XH(PEt_3)_2(\mu\text{-}ZH_2)RuCl_2(\eta^6\text{-}MeC_6H_4CHMe_2\text{-}p)]$ (X = Br, Z = P; X = Cl, Z = As), 2831–8
The mechanism of the photochemical production of hydrogen from aqueous solutions of hydridotris(triethylphosphine)platinum(II), 2941–6
A comparison of the efficiencies of polymer-supported phosphine and phosphine oxide complexes of manganese(II), $MnX_2(C_6H_4PPh_2)$ and $MnX_2\{C_6H_4P(O)Ph_2\}_n$ (*n* = 2 or 4; X = Cl, Br, or I), for the absorption of sulphur dioxide, 2959–62
Synthesis and chemistry of *trans*-$[RhX(CO)L_2]$ (X = anionic ligand, L = tertiary phosphine), 2969–74
Instant ligands. Part 2. Preparation of some bidentate ligands containing PF_2O Groups from aromatic hydroxy compounds and their reactions to form molybdenum and tungsten complexes, 3023–8

PHOSPHINE ALKENOATE
Co-ordination of highly substituted alkenes to transition-metal complexes: preparation and crystal and molecular structures of the complexes $[\{RhCl(Ph_2PO_2CCH{=}CMe_2)\}_2]$ and $[\{RhCl(Ph_2PO_2CCH{=}CHCH{=}CHMe)\}_2]$, 443–50

PHOSPHINE OXIDE
The synthesis of silica-supported phosphine and phosphine oxide complexes of manganese(II) iodide and their reaction with sulphur dioxide, 1415–8
A comparison of the efficiencies of polymer-supported phosphine and phosphine oxide complexes of manganese(II), $MnX_2(C_6H_4PPh_2)$ and $MnX_2\{C_6H_4P(O)Ph_2\}_n$ (*n* = 2 or 4; X = Cl, Br, or I), for the absorption of sulphur dioxide, 2959–62

PHOSPHINE SULPHIDE
Preparations, crystal structures, and molybdenum-95 nuclear magnetic resonance spectroscopic studies of triphenylphosphine–gold–tetrathiomolybdate(VI) clusters, 303–6

PHOSPHINITE
Mixed-metal phosphinito complexes of platinum(II) and palladium(II) with lanthanide and actinide elements. The single-crystal *X*-ray structure determination of $[UO_2(OH_2)\{(OPPh_2)_2Pd(S_2CNEt_2)\}_2]$, 2853–6

PHOSPHINOALKYLSILYL
Phosphinoalkylsilyl complexes. Part 7. Synthesis and reactivity of octahedral, bis(chelate)dicarbonyl complexes of ruthenium(II) and osmium(II). Conformational properties of the bis(chelate) framework and *X*-ray crystal structures of $[M(PPh_2CH_2CH_2SiMe_2)_2(CO)_2]$ (M = Ru or Os), 969–74

PHOSPHITE
trans to *cis* isomerisations of platinum(II) olefin phosphine complexes, $[PtX_2(ol)(PR_3)]$ (X = Cl or Br; ol = ethene, propene, or 1-heptene; PR_3 = PBu_3 or PMe_2Ph), 139–44
The chemistry of vitamin B_{12}. Part 27. The cage phosphite ester 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane and the anionic dimethyl phosphite as ligands for cobalt(III) corrinoids, 761–8
New dicyclopentadienyl phosphine, phosphite, and acetylene niobium(III) complexes. Crystal structure of $[Nb(\eta\text{-}C_5H_4SiMe_3)_2Cl(PhC{\equiv}CPh)]$, 975–80
Carbon-13 and phosphorus-31 nuclear magnetic resonance studies of the fluxional behaviour of the complexes

PHOSPHITE (contd)
$[Fe_3(CO)_{10}\{P(OMe)_3\}_2]$ and $[Fe_3(CO)_{10}\{P(OMe)_3\}\{P(OPh)_3\}]$, 1249–52
Cationic ruthenium(II) phosphite complexes. Preparation and properties of monocarbonyl and isocyanide derivatives, 1813–6
PHOSPHONATE
Solid-state nuclear magnetic resonance spectroscopic study of γ-zirconium phosphate, 1877–82
PHOSPHONIODITHIOCARBOXYLATE
Manganese(I) complexes with (tricyclohexylphosphonio)dithiocarboxylate as chelate and unidentate ligand. *X*-Ray crystal structure of *fac*-$[Mn(CO)_3\{S_2CP(C_6H_{11})_3\}_2]ClO_4 \cdot H_2O$, 2875–80
Mono- and bi-nuclear gold(I) and gold(III) complexes with $S_2C–PR_3$ ligands. *X*-Ray crystal structures of $[Au(C_6F_5)_3(S_2C–PEt_3)]$ and $[Au_2(\mu\text{-}S_2CPEt_3)(C_6F_5)_6] \cdot CH_2Cl_2$, 3017–22
PHOSPHORUS
Oxidation and reduction of phosphorus–phosphorus and arsenic–arsenic double bonds. An electrochemical study of two diphosphenes and a diarsene, 249–52
Direct synthesis of ammonium monofluorophosphate monohydrate, $[NH_4]_2[PO_3F] \cdot H_2O$, and potassium monofluorophosphate, $K_2[PO_3F]$, 477–8
Studies of phosphazenes. Part 30. Reactions of hexachlorocyclotriphosphazene with aromatic primary amines: interplay of geminal and non-geminal modes of chlorine replacement, 579–84
Studies of phosphazenes. Part 31. Kinetic studies of the reactions of hexachlorocyclotriphosphazene with aromatic primary amines: evidence for the intermediacy of a reactive three-co-ordinated phosphorus(V) species, 585–90
Menthyl-substituted organophosphorus compounds. Part 6. Preparation of P(L-men)RCl and P(L-men)R(S)Cl (R = Me, Et, Pr^i Bu^t, or Ph; men = *cyclo*-C_6H_9-2-Pr^i-5-Me) and nuclear magnetic resonance studies of configuration and halogen exchange, 795–806
Structure and basicity. Part 10. Conductometric and potentiometric titrations of some cyclotriphosphazatrienes in non-aqueous media. The conductivities of alkali-metal acetates on titration with perchloric acid in acetic acid, 925–30
Phosphorus-31 nuclear magnetic resonance parameters for compounds with the α-tetraphosphorus trisulphide skeleton; substituent constants for an inorganic substrate, 1515–20
Chemistry of 2,3-dihydro-2-metallaphenalenes. Part 2. Formation of $[\overline{CHMMe_2C(MMe_3)}C_{10}H_6]^x$ *via* base decomposition of $[1,8\text{-}(Me_3MCH_2)_2C_{10}H_6]^y$ (M = Ge, $x = 2-$, $y = 0$; M = P, $x = 0$, $y = 2+$); crystal structures of the novel bis-ylide $[\overline{CHPMe_2C(PMe_3)}C_{10}H_6$ and of a distorted *peri*-ylide, 1-Me-8-$(Me_3PCH)C_{10}H_6$, 1647–54
Kinetic study of the homogeneous methanolysis of tetraphosphorus decasulphide, 1897–900
Phosphorus–nitrogen compounds. Part 51. Carbon-13 and phosphorus-31 nuclear magnetic resonance properties of phenylphosphonothioic di(monoalkylamides). Determination of aromatic substituent constants by ^{13}C nuclear magnetic resonance spectroscopy, 2479–84
Mean phosphorus– and arsenic–sulphur bond enthalpies in diethyldithiocarbamate chelate complexes, 2955–8
PHOSPHORUS-31
Studies of phosphazenes. Part 30. Reactions of hexachlorocyclotriphosphazene with aromatic primary amines: interplay of geminal and non-geminal modes of chlorine replacement, 579–84
Proton and phosphorus-31 nuclear magnetic resonance study of zinc(II) *O,O'*-dialkyl dithiophosphates in solution, 807–14
New heterocyclic compounds containing niobium and molybdenum; crystal structure of a cyclomolybdaphosphazene, 847–50
Lewis-base adducts of Group 11 metal(I) compounds. Part 27. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance, far-infrared, and structural studies on the mononuclear 2:1 adducts of triphenylphosphine with copper(I) and gold(I) halides, 1089–98
Lewis-base adducts of Group 11 metal(I) compounds. Part 28. Solid-state phosphorus-31 cross-polarization magic-angle spinning nuclear magnetic resonance and structural studies on the mononuclear 3:1 adducts of triphenylphosphine with copper(I) halides, 1099–106
Phosphorus-31 nuclear magnetic resonance parameters for compounds with the α-tetraphosphorus trisulphide skeleton; substituent constants for an inorganic substrate, 1515–20
PHOTOCHEMISTRY
Synthesis of $[Mo(\eta\text{-}C_5H_4Pr^i)(PR_3)_2H_3]$($R_3$ = Me_3 or Me_2Ph), $[Mo(\eta\text{-}C_5H_4Pr^i)(Pr^i{}_2PCH_2CH_2PPr^i{}_2)H_3]$, and analogues and their ability to catalyse the photoinduced hydrogen-deuterium exchange of carbon-hydrogen bonds; crystal structure of $[\{Mo(\eta\text{-}C_5H_4Pr^i)(\mu\text{-}Cl)_2\}_2]$, 9–20
Electronic absorption spectra and photochemical reactivity of Group 5 metal alkyl compounds: photochemical α-hydrogen abstraction, 163–8
Thermal and photochemical behaviour of organotetraruthenium clusters: solution structures and dynamics of phosphine-substituted derivatives, 349–54
Kinetics and mechanism of ring closure for a series of photoproduced $[M(CO)_5L]$ intermediates (M = Cr, Mo, or W; L = pyridine 2-carbaldehyde imine ligand), 513–8
Photo-oxidations of acetoin and biacetyl catalysed by tetrabutylammonium decatungstate(4−) in acetonitrile under excess of oxygen, 961–2
Photoactivation of methane by η^5-cyclopentadienyl and substituted η^5-cyclopentadienyl Group 8 metal dicarbonyl complexes, $[M(\eta^5\text{-}C_5R_5)(CO)_2]$ (M = Rh or Ir, R = H or Me), and dicarbonyl(η^5-indenyl)iridium: a matrix isolation study, 1181–90
Water photolysis. Part 2. An investigation of the relative advantages of various components of the sensitiser–electron relay–metal colloid system for the photoproduction of hydrogen from water, and the use of these systems in the photohydrogenation of unsaturated organic substrates, 1299–304
Involvement of hydrogen-bonding protons in delocalization of the paramagnetic electron in a single crystal of photoreduced decatungstate, 1597–604
Photochemical preparation and reactivity of platinum(II) complexes with oxygen-donor ligands. Description and synthesis of di-μ-chloro-dichlorobis(substituted pyridine)diplatinum(II), 1687–92
Di-2-pyridyl ketone complexes of copper(I): efficient photocatalysts for norbornadiene–quadricyclane conversion, 1797–800
Photochemical reduction of carbon dioxide to carbon monoxide in water using a nickel(II) tetra-azamacrocycle complex as catalyst, 2105–10
The mechanism of the photochemical production of hydrogen from aqueous solutions of hydridotris(triethylphos-phine)platinum(II), 2941–6
Synthesis, characterisation, and photochemical properties of a series of ruthenium containing metallopolymers based on poly-*N*-vinylimidazole, 3073–8
PHOTOELECTRON
Application of ultraviolet photoelectron spectroscopy to molecular properties. Part 27. Stereochemistry of ligand exchange reactions of titanocene derivatives, 107–10
Bonding in and gas-phase pyrolysis of the arsoranes and stibanes EMe_3X_2 (E = As or Sb, X = F or Cl) and $SbMe_4F$: ultraviolet photoelectron and field-ionization mass spectrometric studies, 481–8
Iron antimonate, 1727–32
Surface characterisation of α-, β-, γ-, and δ-UO_3 using *X*-ray photoelectron spectroscopy, 3009–16
PHOTOSYNTHESIS
Formation of a (4 + 4) Schiff-base macrocyclic ligand by a template rearrangement. Crystal and molecular structures of two tetranuclear manganese(II) complexes, 2555–62
PHTHALATE
Correlation between thermal stability and molecular and crystal structure in a series of potassium–dibenzo-18-crown-6 solid compounds, 1779–84
PHTHALOCYANINATE
Equilibrium and kinetic study of nitric oxide binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 369–72
Disproportionation, electrochemistry, and electronic coupling involving mononuclear and binuclear cobalt phthalocyanine derivatives, 2511–8
PIPERAZINE
Thermal investigation and stereochemical studies of some cyclic diamine complexes of nickel(II), zinc(II), and cadmium(II) in the solid state, 2089–94

PLASTOCYANIN
Kinetic studies on 1:1 electron-transfer reactions involving blue copper proteins. Part 14. Reactions of poplar plastocyanin with inorganic complexes, 2573–8

PLATINUM
Linear-chain platinum(II) dithiocarboxylates: crystal structure of tetrakis(phenyldithioacetato)diplatinum(II), 35–40
trans to *cis* isomerisations of platinum(II) olefin phosphine complexes, $[PtX_2(ol)(PR_3)]$ (X = Cl or Br; ol = ethene, propene, or 1-heptene; PR_3 = PBu_3 or PMe_2Ph), 139–44
The synthesis of mono- and bi-metallic complexes containing the ligand bis(diphenylarsino)methane, 151–4
Bimetallic systems. Part 16. Formation of Group 6 metal(0)–platinum(II) hydride complexes, by ring opening of *fac*- or *mer*-$[M(CO)_3(dppm\text{-}P\,P')(dppm\text{-}P)]$ (M = Cr, Mo, or W; dppm = $Ph_2PCH_2PPh_2$). Reactions of $[(CO)_3Mo(\mu\text{-}dppm)_2PtH(Cl)]$, 221–6
Synthesis and *X*-ray crystal structure of a cuboctahedral platinum cluster, 259–62
trans-Dichlorotetrakis(pyridine)platinum(IV) nitrate: a classical co-ordination compound, 293–6
Chemistry of platinum sulphido-complexes. Part 6. Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with the rhodium complexes $[(RhL_2Cl)_2]$ (L = CO or C_2H_4) and $[Rh(2,6\text{-}Me_2C_6H_3NC)_4]PF_6$. Crystal and molecular structures of $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(CO)_2]\text{-}[Rh(CO)_2Cl_2]\cdot C_6H_6O$ and $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4\text{-}(2,6\text{-}Me_2C_6H_3NC)_2]PF_6$, 335–40
Sulphur–arsenic ligands. Part 1. Dimethyl(methylthio)arsine complexes of halogenotrimethylplatinum(IV) and tricarbonylhalogenorhenium(I), 417–20
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of
$[Pt\{ON(Ph)C(NPh)S\}(PPh_3)_2]$ and
$[Pt\{ON(Ph)C(O)NPh\}(PPh_3)_2]$, 605–8
Reactions of heteroboranes. Synthesis of $[2,2\text{-}(PPh_3)_2\text{-}1,2\text{-}SePtB_{10}H_{10}]\cdot CH_2Cl_2$, its crystal and molecular structure and that of $SeB_{11}H_{11}$, 699–704
Bis[bis(diphenylphosphino)methane]platinum(II) salts: evidence for the formation of a five-co-ordinate bis[bis(diphenylphosphino)-methane]halogenoplatinum(II) cation, 705–8
Metallacyclic complexes. Synthesis of metallathietane-3,3-dioxide (metallacyclosulphone) complexes of platinum(II), palladium(II), and nickel(II); crystal structure of *trans*-2,4-diphenyl-1,1-bis(triphenylarsine)platinathietane-3,3-dioxide, 733–40
Reactions of cyclobutenedione and cyclopentenetrione with zero-valent nickel triad complexes of isocyanide. *X*-Ray crystal structures of (3—4-η-3,4-diphenylcyclobut-3-ene-1,2-dione)- and (4—5-η-4,5-diphenylcyclopent-4-ene-1,2,3-trione)-tris(t-butyl isocyanide)nickel, 853–8
Synthesis of *o*-nitrophenylplatinum(IV) complexes; crystal and molecular structure of dichlorobis(2-nitrophenyl-*C'O*)-platinum(IV), 881–4
Rational synthesis of metalla–sulphur–nitrogen complexes: use of $[Me_2SnS_2N_2H_2]$ in the preparation of complexes of the type $[M(S_2N_2H)(PR_3)_2]X$ (M = Pt or Pd). *X*-Ray structures of $[Pt(S_2N_2H)(PEt_3)_2][Me_2SnCl_3]$, $[Pt(S_2N_2H)(PBu^n_3)_2]_2[PF_6]Cl$, and $[\{Co(S_2N_2H)_2\}_2\{Ph_2P(O)CH_2CH_2P(O)Ph_2\}]$, 907–14
Electron-transfer platinum complexes of esters of dithiocarbazic acid. The crystal and molecular structure of $[Pt\{N(CH_2Ph)NC(S)SMe\}_2]$, 1035–40
Reactions of planar pentadentate ligands with palladium and platinum chlorides. Crystal structures of *cis*-dichlorobis(*N'*-isopropylidenebenzohydrazide)platinum(II) and the cyclopalladated dianion of *N',N'''*-(pyridine-2,6-diyldiethylidyne)bis(benzohydrazide), 1175–80
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 59. Reactions of the alkylidyne compounds $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Cr, Mo, or W, R = C_6H_4Me-4; M = W, R = Me, $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate] with d^{10} complexes; crystal structure of $[PtW_2(\mu\text{-}CMe)_2(CO)_4\text{-}\{HB(pz)_3\}_2]$, 1229–34
Syntheses and characterization of dinuclear palladium and platinum complexes with a single unsupported bridging group. Crystal structure of $[\{C_6H_3(CH_2NMe_2)_2\text{-}o,o'\}Pd(\mu\text{-}Cl)Pd\text{-}\{C_6H_3(CH_2NMe_2)_2\text{-}o,o'\}]BF_4$, 1359–66
Kinetic studies on the oxidation of ethylenediaminebis(pyridine)- and bis(ethylenediamine)-platinum(II) complexes by hexachloroiridate(IV), 1473–6
Bimetallic systems. Part 21. Synthesis of vinylidene- and phenylene-bridged complexes of dinickel(II), nickel(II)–palladium(II), and nickel(II)–platinum(II) containing $Ph_2PCH_2PPh_2$ (dppm): crystal structures of $[(SCN)Ni(\mu\text{-}dppm)_2(\mu\text{-}C{=}CH_2)Ni(NCS)]$ and $[ClPd(\mu\text{-}dppm)_2(\mu\text{-}C{=}CCl_2)NiCl]$, 1501–8
Raman and resonance-Raman studies of platinum group tetrathiomolybdates and tetrathiotungstates, 1535–44
Insertion of CO_2-like molecules into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$: the *X*-ray structures of
$[Pt\{ON(Ph)C(NPh)S\}(PPh_3)_2]$ and
$[Pt\{ON(Ph)C(O)NPh\}(PPh_3)_2]$ (1987, 605), 1583–4
Unsymmetrical bridging of an acetylide ligand in the cation $[Pt_2(C{\equiv}CBu^t)_3(\mu\text{-}dppm)_2]^+$ (dppm = $Ph_2PCH_2PPh_2$): *X*-ray structure determination of the tetraphenylborate salt, 1659–64
Photochemical preparation and reactivity of platinum(II) complexes with oxygen-donor ligands. Description and synthesis of di-μ-chloro-dichlorobis(substituted pyridine)diplatinum(II), 1687–92
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H$(PPh_3)_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$], 1805–12
Mixed chloro-, *NN*-dialkyldithiocarbamato-, and *S*-alkyl *NN*-dialkyldithiocarbamate complexes of platinum(II). Crystal structure of $[PtCl(S_2CNEt_2)(MeS_2CNEt_2)]$, 1823–6
Electrochemical reactions of dichlorobis(aryl isocyanide)platinum(II) complexes, 1833–8
A general synthetic route for platinum cluster compounds containing carbonyl and tertiary phosphine ligands and a study of their reactions with unsaturated inorganic molecules, 1889–96
Pentafluorophenyl complexes of palladium and platinum containing chelating, unidentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 2077–82
Platinum(II) complexes with ligands $(RO)_2PCH_2P(OR)_2$ (R = Me, Et, Ph, or C_6H_4Me-4); crystal structure of *cis,cis*-$[Pt_2Me_4\{\mu\text{-}(EtO)_2PCH_2P(OEt)_2\}_2]$, 2117–24
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 62. Synthesis of penta-, hexa-, and hepta-heteronuclear metal cluster compounds involving tungsten or molybdenum with platinum or nickel: crystal structures of the chain compounds $[Pt_3W_3(\mu\text{-}CMe)(\mu_3\text{-}CMe)_2(CO)_6(cod)(\eta\text{-}C_5Me_5)_3]\cdot 0.5CH_2Cl_2$ and $[Pt_3W_4(\mu\text{-}CC_6H_4Me\text{-}4)_2(\mu_3\text{-}CC_6H_4Me\text{-}4)_2(CO)_8(\eta\text{-}C_5H_5)_4]$, 2177–88
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 63. Synthesis of eight-membered-ring metallacycles: *X*-ray crystal structures of $[Pt_4W_4(\mu\text{-}CR)(\mu_3\text{-}CR)_3(\mu\text{-}CO)(CO)_7(\eta\text{-}C_5H_5)_4]\cdot CH_2Cl_2$, $[Ni_2Pt_2W_4(\mu\text{-}CR)(\mu_3\text{-}CR)_3(\mu\text{-}CO)(CO)_7(\eta\text{-}C_5H_5)_4]\cdot CH_2Cl_2$ (R = C_6H_4Me-4), and $[Ni_2Pt_2W_4(\mu_3\text{-}CPh)_4(CO)_8(\eta\text{-}C_5H_5)_4]$, 2189–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 66. Carbaboranetungsten– platinum complexes. Polyhedral rearrangements of a 12-vertex cage system; crystal structures of $[PtW(CO)_2(PEt_3)_2\{\eta^6\text{-}C_2B_9H_8\text{-}(CH_2C_6H_4Me\text{-}4)Me_2\}]\cdot CH_2Cl_2$, $[PtW(\mu\text{-}H)\{\mu\text{-}\sigma{:}\eta^5\text{-}C_2B_9H_7(CH_2C_6H_4Me\text{-}4)Me_2\}(CO)_2(PMe_3)(PEt_3)_2]$, and related compounds, 2219–34
A reversible metal framework rearrangement in high-nuclearity osmium-platinum cluster compounds. *X*-Ray crystal structures of $[Os_6Pt(CO)_{17}(\mu_3\text{-}NCMe)(C_8H_{12})]$, $[Os_6Pt(CO)_{17}(\mu_4\text{-}NCMe)\text{-}(C_8H_{12})]$ (C_8H_{12} = cyclo-octa-1,5-diene), and $[Os_6(CO)_{19}(\mu_3\text{-}NCMe)]$: compounds with novel metal–acetonitrile bonding modes, 2245–52
Synthesis, reactivity, and *X*-ray crystal structures of $[Os_6Pt_2H_2(CO)_{16}(\mu\text{-}C_8H_{10})(C_6H_{12})]\cdot CH_2Cl_2$ and $[Os_6H(CO)_{16}(\mu\text{-}C_8H_{11})]$. Model compounds for C–H activation, 2253–60
Synthesis of chelate stabilised alcohol- and alkoxo–platinum(II) complexes. *X*-Ray crystal structure of
$[Pt(PPh_2CH_2CMe_2O)_2]\cdot 3.5H_2O$, 2273–80
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 67. Reactions of the salts $[X][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ [X = $P(CH_2Ph)Ph_3$, R = C_6H_4Me-4; X = NEt_4, R = Ph] with bis(cyclo-octa-1,5-diene)platinum, 2661–8
Synthesis and study of monomeric and dimeric amidine complexes of nickel(II) and platinum(II). Crystal structure of bis(*N,N'*-diphenylbenzamidino)platinum(II), $[Pt\{PhNC(Ph)NPh\}_2]$, 2687–94

PLATINUM (contd)
A two-dimensional nuclear magnetic resonance exchange study of the fluxionality of $[PtIMe_3(MeSCH_2CH_2SEt)]$, 2695–702
Diplatinum-(II) and -(III) complexes with bridging μ-diphosphonito-*P*,*P*′ or μ-methylenebis(phosphonito)-*P*,*P*′ ligands. Synthesis, structure, and nuclear magnetic resonance analysis of the 'lantern' complexes. Crystal structures of $K_4[Pt_2\{\mu\text{-}CH_2[P(O)OH]_2\}_4]\cdot 6H_2O$ and $K_4[Pt_2\{\mu\text{-}CH_2[P(O)OH]_2\}_4Cl_2]\cdot 8H_2O$, 2769–80
Mixed-metal phosphinito complexes of platinum(II) and palladium(II) with lanthanide and actinide elements. The single-crystal *X*-ray structure determination of $[UO_2(OH_2)\{(OPPh_2)_2Pd(S_2CNEt_2)\}_2]$, 2853–6
The mechanism of the photochemical production of hydrogen from aqueous solutions of hydridotris(triethylphosphine)platinum(II), 2941–6
Platinum(II)– and palladium(II)–iron(0) complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand. Crystal structures of $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}CO)PtBr_2]$ and $[(OC)_3Fe(\mu\text{-}Ph_2PCH_2PPh_2)(\mu\text{-}I)PtI]\cdot CH_2Cl_2$, 3079–84
Preparation of two mixed-metal carbonyl clusters based on the platinumtriosmium core. *X*-Ray crystal structure of $[Os_3PtH_2(cod)(CH_2)(CO)_9]$ (cod = cyclo-octa-1,5-diene), 3103–6
Metallacarboranes derived from the $[C_2B_{10}H_{12}]^{2-}$ anion. The preparation, molecular structure, and properties of [4-(dppe)-4-Pd-1,6-$C_2B_{10}H_{12}$], [4,4,4-H(PPh$_3$)$_2$-2-OMe-4-Ir-1,6-$C_2B_{10}H_{11}$], and [3-(dppe)-3-Pt-1,2-$C_2B_9H_{11}$] (1987, 1805), 3107–8
PLATINUM-195
Bis[bis(diphenylphosphino)methane]platinum(II) salts: evidence for the formation of a five-co-ordinate bis[bis(diphenylphosphino)-methane]halogenoplatinum(II) cation, 705–8
POLYMER
Infrared spectroscopic studies of uranyl(VI) species adsorbed from aqueous $[UO_2(CO_3)_3]^{4-}$ solutions on to a polymer bearing amidoxime groups, 2489–92
Synthesis, characterisation, and photochemical properties of a series of ruthenium containing metallopolymers based on poly-*N*-vinylimidazole, 3073–8
PORPHYRIN
Reverse Bohr effect on the oxygen-binding affinity of heme embedded in a bilayer of liposome as a hemoglobin model: pH-induced oxygen uptake and evolution by aqueous synthetic lipid-heme solution, 1917–20
Iron–ligand bonding properties of synthetic iron–porphyrin complexes with oxygen transporting ability in aqueous media, 2455–8
Dynamics of oxygen binding to modified iron(II)–porphyrin complexes embedded in the bilayer of phospholipid liposome, 2493–6
Oxygen- and carbon monoxide-binding to a lipophilic diporphyrinatocopperiron complex solubilized in an aqueous medium with a micelle, 2963–8
An electron spin resonance study of frozen aqueous solutions containing 5,10,15,20-tetrakis(*N*-methyl-4′-pyridinio) porphyrinatocobalt(II), 3099–102
POTASSIUM
Correlation between thermal stability and molecular and crystal structure in a series of potassium–dibenzo-18-crown-6 solid compounds, 1779–84
Preparation and spectroscopic study of functionally substituted cyclopentadienides of thallium(I), potassium, rhodium(I), and iridium(I), 2615–20
PROPANAMIDE
Design of transition metal chelates with biological activity. Potentiometric study of complex formation equilibria between 2-amino-*N*-hydroxy-3-(*p*-hydroxyphenyl)propanamide and nickel(II), copper(II), and hydrogen ions in aqueous solution, 1409–14
PROTEIN
Kinetic studies on 1:1 electron-transfer reactions involving blue copper proteins. Part 14. Reactions of poplar plastocyanin with inorganic complexes, 2573–8
PROTON TRANSFER
The base catalysed hydrolysis of *trans*-dichloro-bis(1,2-diaminoethane)cobalt(III) and -(1,3-diaminopropane)cobalt(III) cations: rate-limiting proton transfer, 1775–8
PULSE RADIOLYSIS
Reactions of chromium(III) complexes of 1,10-phenanthroline, 2,2′-bipyridyl, and oxalate with the pulse radiolytically generated aquated electron, zinc(I), and cadmium(I), 1425–30
PYRAZOLATE
Binuclear rhodium and iridium complexes containing pentamethylcyclopentadienyl and pyrazolate ligands, 639–46
PYRAZOLE
Reactivity of ruthenium(II) complexes with 1-hydroxymethyl- or 1-benzyl-3,5-dimethylpyrazole and similar functionalized bipyrazoles. *X*-Ray crystal structure of carbonylchlorohydrido(3,5-dimethylpyrazole-N^2)-bis(triphenylphosphine)ruthenium(II), 183–6
Metal–ligand bonding parameters and magnetic properties of some previously reported tetragonal nickel(II) complexes, 825–30
PYRAZOLYL
A molybdenum-95 and nitrogen-14 nuclear magnetic resonance study of six-co-ordinate hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes containing a sixteen-electron $\{Mo(NO)\}^4$ core, 619–22
Reactions of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}X_2]$ (M = Mo, X = I; M = W, X = Cl) with some amine-substituted pyridines. The molecular structures of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}(2\text{-}NHC_5NH_4)_2]$, 1063–70
Structure and magnetism of analogous *O*-bridged (phenolato) and *S*-bridged (thiophenolato) binuclear copper(II) complexes. Crystal structure of the binuclear *S*-bridged complex μ-[2,6-bis(4′-cyclohexyl-4′-hydroxy-2′,3′-diazabuta-1′,3′-dien-1′-yl)-4-methylthiophenolato(3−)-*S*,N^2,*O*(Cu^{1,2})]-methanol-μ-(pyrazolyl-*N*,*N*′)-dicopper(II), 1585–92
PYRAZOLYLBORATE
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 59. Reactions of the alkylidyne compounds $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Cr, Mo, or W, R = C_6H_4Me-4; M = W, R = Me, $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate] with d^{10} complexes; crystal structure of $[PtW_2(\mu\text{-}CMe)_2(CO)_4\{HB(pz)_3\}_2]$, 1229–34
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 60. Di- and tri-metal compounds prepared from the metal alkylidyne complexes $[W(\equiv CR)(CO)_2(\eta\text{-}C_5Me_5)]$ (R = Me or C_6H_4Me-4) and $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Mo, R = C_6H_4Me-4; M = W, R = Me; $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate], 1235–42
PYRIDINE
Investigation into aroylhydrazones as chelating agents. Part 8. Synthesis and spectroscopic characterization of complexes of Co, Ni, Cu, Zn, and Cd with 2,6-diacetylpyridine bis(salicyloylhydrazone); *X*-ray crystal structure of dichloro[2,6-diacetylpyridine bis(salicyloylhydrazone)]cadmium(II)–chloroform–methanol (1/1/1), 177–82
trans-Dichlorotetrakis(pyridine)platinum(IV) nitrate: a classical co-ordination compound, 293–6
Reaction of $[Os_5H_2(CO)_{15}]$ with pyridine: crystal and molecular structures of $[Os_5H_3(CO)_{14}(C_5H_4N)]$ and $[Os_5H_2(CO)_{14}(C_5H_5N)]$, 327–34
Synthetic, spectroscopic, and *X*-ray crystallographic studies on binuclear copper(II) complexes with a tridentate NNS-bonding 2-formylpyridine thiosemicarbazone ligand. The characterization of both neutral and deprotonated co-ordinated ligand structures, 493–500
Metal–ligand bonding parameters and magnetic properties of some previously reported tetragonal nickel(II) complexes, 825–30
Reactions of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}X_2]$ (M = Mo, X = I; M = W, X = Cl) with some amine-substituted pyridines. The molecular structures of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}(2\text{-}NHC_5NH_4)_2]$, 1063–70
Kinetic studies on the oxidation of ethylenediaminebis(pyridine)- and bis(ethylenediamine)-platinum(II) complexes by hexachloroiridate(IV), 1473–6
Cyclopalladation to form planar tridentate $[N\text{–}C\text{–}N]^-$ intramolecular co-ordination systems involving pyridine donor groups, including ligand synthesis and *X*-ray structural studies., 1477–84
Synthetic, spectroscopic, and *X*-ray crystallographic studies on binuclear copper(II) complexes with a tridentate NNS-bonding 2-formylpyridine thiosemicarbazone ligand. The characterization of both neutral and deprotonated co-ordinated ligand structures (1987, 493), 1583–4
New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states, 2621–30

PYRIDINE (contd)
Electronic structure and redox reactions of vanadium(II) polypyridine, isocyanide, and phosphine complexes, 2881–4
Structure and electrochemical properties of bis(pyridine)cobalt(III) complexes of *N,N'*-ethylenebis(salicylideneimine) homologues with an aryl substituent on the ethylene backbone. Crystal structures of $[Co(salphen)(py)_2]PF_6 \cdot 0.5Et_2O$ and $[Co(salcppn)(py)_2]PF_6 \cdot 2py$ (py = pyridine), 3055–60
Syntheses and crystal structures of complexes $[M_2R_2]$, [M = Cu, Ag, or Au; R = 2-C(SiMe$_3$)$_2$C$_5$H$_4$N] and $[Cu_4R'_4]$[R' = 2-CH(SiMe$_3$)C$_5$H$_4$N]; electrochemical generation of $[Cu_2R_2]^{2+}$, 3085–92

PYRIDINECARBALDIMINE
Kinetics and mechanism of ring closure for a series of photoproduced $[M(CO)_5L]$ intermediates (M = Cr, Mo, or W; L = pyridine 2-carbaldehyde imine ligand), 513–8

PYRIDINECARBOXYLATE
Ternary metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of copper(II)–2,2'-bipyridyl– pyridoxic acid and copper(II)–2,2'-bipyridyl–pyridoxic acid γ-lactone complexes, 1009–16

PYRIDINETHIOLATE
Methyl- and phenyl-mercury(II) derivatives of 2-mercaptopyridine. Crystal and molecular structure of methyl(pyridine-2-thiolato)mercury(II) (1986, 1945), 683–4

PYRIDINETHIONE
Crystal structure and mass spectrometry of dichlorodimethylbis[2(1*H*)-pyridinethione-*S*]tin(IV), 815–8
The synthesis, crystal and molecular structure of a copper(I) complex of 2(1*H*)-pyridinethione: an intermediate in Cu_4S_4 heterocubane build up, 2281–4

PYRIDINONE
Ternary metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of copper(II)–2,2'-bipyridyl– pyridoxic acid and copper(II)–2,2'-bipyridyl–pyridoxic acid γ-lactone complexes, 1009–16

PYRIDONATE
2-Pyridonate (Opy) rhodium(I) carbonyl complexes. Crystal structure of the unusual tetranuclear complex $[Rh_4(\mu\text{-Opy})_4(\mu\text{-CO})_2(CO)_4]$, 981–4

PYRIDOXAL
Thiosemicarbazones as co-ordinating agents. Part 2. Synthesis, spectroscopic characterization, and *X*-ray structure of aquachloro(pyridoxal thiosemicarbazone)manganese(II) chloride and aqua(pyridoxal thiosemicarbazonato)copper(II) chloride monohydrate, 227–34

PYRIDYLTRIAZOLE
Synthesis, spectroscopic, and electrochemical properties of bis(2,2'-bipyridyl)ruthenium compounds of some pyridyl-1,2,4-triazoles, 1389–96

PYROLYSIS
Bonding in and gas-phase pyrolysis of the arsoranes and stibanes EMe_3X_2 (E = As or Sb, X = F or Cl) and $SbMe_4F$: ultraviolet photoelectron and field-ionization mass spectrometric studies, 481–8

PYRROLIDINE
Synthesis and stereochemical studies of the chiral ruthenium complexes $[Ru(\eta\text{-}C_5H_5)\{(S)\text{dpompyr-}PP'\}X]$ [dpompyr = *N*-diphenylphosphino-2-(diphenylphosphinoxymethyl)pyrrolidine, X = H or Cl]. Crystal structure of $[(S)Ru(\eta\text{-}C_5H_5)\{(S)\text{dpompyr-}PP'\}Cl]$, 1149–56

QUINOLINE
Characterizations of the redox orbitals of mixed-ligand 2,2'-bipyridine–2-(2'-pyridyl)quinoline ruthenium(II) complexes, 2467–72

QUINOLINOLATE
The single-crystal electron spin resonance spectrum of oxobis(2-methylquinolin-8-olato)vanadium(IV) as a pure compound and diluted in chlorobis(2-methylquinolin-8-olato)gallium(III), 111–8

QUINOXALINEDITHIOLATE
Structure and properties of tris(quinoxaline-2,3-dithiolato)-molybdate(IV), 297–302
A structural comparison of bis(quinoxaline-2,3-dithiolato)cuprate complexes, 1083–8
Crystal structure of tris(quinoxaline-2,3-dithiolato)molybdate(V); a distorted trigonal prismatic complex, 2267–72

RADICAL
Hydrogen halide radical anions, 421–6
High nuclearity osmium cluster radicals, 1051–4

RADICAL CATION
The radical cations of sulphur ($S_8^{\cdot+}$) and tetrasulphur tetranitride ($S_4N_4^{\cdot+}$): a radiation–electron spin resonance study, 729–32

RADIOLYSIS
Hydrogen halide radical anions, 421–6

RAMAN
Spectrochemistry of solutions. Part 19. Complexation of gold(I) by cyanide in liquid ammonia: an infrared and Raman spectroscopic study, 673–6
The vibrational spectrum of tricarbonyl(trichlorostibine)nickel(0), $[Ni(CO)_3(SbCl_3)]$, 681–2
Raman and resonance-Raman studies of platinum group tetrathiomolybdates and tetrathiotungstates, 1535–44
The preparation and crystal structure of $5SbF_3 \cdot 3SbF_5$, 1989–98
Studies of metal carbonate equilibria. Part 15. The beryllium(II)–water–carbon dioxide(g) system in acidic 3.0 mol dm^{-3} perchlorate media, 2439–44
Preparations, properties, and vibrational spectra of some (dimethylamino)halogenosilanes, 3029–34

REDOX POTENTIAL
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato-molybdenum and -tungsten nitrosyls. Part 4. The effects of ligating atom type on the reduction potentials of monometallic complexes, 1733–6
The reversible one-electron oxidation of *trans*-$[Mn^IL(CO)(Ph_2PCH_2PPh_2)_2]^{z+}$: synthesis and characterisation of stable manganese(II) derivatives, 1769–74
Co-ordination chemistry of higher oxidation states. Part 23. Synthesis and properties of tetrahalogenoiridium(IV) complexes, $[IrL_2X_4]$ [X = Cl or Br; L = pyridine, PR_3, AsR_3, SbR_3, SR_2, or SeR_2]. Crystal and molecular structure of *trans*-$[Ir(AsEt_3)_2Br_4]$, 1901–10

REDUCTION
Electrocatalytic reduction of carbon dioxide to methanol. Part 5. Relationship between the ability of metal complexes to engage in homogeneous catalysis and their co-ordination chemistry, 1377–80

RESOLUTION
Chiral recognition in ion pairing of an optically active tris(ethylenediamine)cobalt(III) cation and application to chromatographic resolution of metal complex anions, 1977–82

RHENIUM
Isomers in *fac*-$[ReBr(CO)_3(L_c)_2]$ (L_c = *cis*-3,4-dihydro-2,3,4,5-tetraphenyl-2*H*-1,2,3-diazaphosphole) arising from chirality of phosphorus ligands: separation and *X*-ray structural studies, 21–6
Oxoaryls of rhenium-(V) and -(VI) and osmium(VI). *X*-Ray crystal structures of dimesityldioxorhenium(VI), tetramesityloxorhenium(VI) and dimesityldioxo-osmium(VI), 169–76
Preparation of isocyanide and mixed dinitrogen–isocyanide complexes of rhenium(I) from reactions of *trans*-$[ReCl(N_2)(PMe_2Ph)_4]$, *mer*-$[Re(S_2PPh_2)(N_2)(PMe_2Ph)_3]$, or *mer*-$[Re(S_2CNEt_2)(N_2)(PMe_2Ph)_3]$ with methyl isocyanide; crystal structure of *mer*-$[Re(S_2PPh_2)(N_2)(CNMe)(PMe_2Ph)_3]$, 319–26
Activation of alkanes involving rhenium atoms: synthesis and electronic structure of binuclear μ-alkylidene derivatives; molecular structure of $[\{Re(\eta\text{-}C_6H_6)\}_2(\mu\text{-}CHBu^t)(\mu\text{-}H)_2]$, 391–402
Synthesis of η-arene rhenium compounds using rhenium atoms. Crystal structures of $[Re_2(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)_2(\mu\text{-}H)_2]$ and $[Re(\eta\text{-}C_6H_6)(\eta\text{-}C_6H_8)H]$, 403–10
Sulphur–arsenic ligands. Part 1. Dimethyl(methylthio)arsine complexes of halogenotrimethylplatinum(IV) and tricarbonylhalogenorhenium(I), 417–20
The chemistry of the hydrazido(1−)-ligand. Preparations and crystal structures of $[Mo(NHNHCO_2Me)(NNCO_2Me)(S_2CNMe_2)_2]$, $[Mo(NHNMePh)(NNMePh)(S_2CNMe_2)_2]BPh_4$, and $[ReCl_2(NHNHCOPh)\text{-}(NNHCOPh)(PPh_3)_2]$. Mechanism of formation of substituted hydrazines from $[Mo(NNRPh)_2(S_2CNMe_2)_2]$ (R = Me or Ph), 529–40
Preparation of the nitrenes $ReF_5(NF)$, $ReF_5(NCl)$, and $ReF_5(NBr)$. *X*-Ray crystal structures of $ReF_5(NF)$ and $ReF_5(NCl)$, 567–72

RHENIUM contd)
Crystal structure of *trans*-dicarbonylbis(dimethylphenylphosphine)-(*N*-phenylsalicylideneaminato)rhenium(I) and *mer*-trichloro-(dimethylphenylphosphine)(*N*-phenylsalicylideneaminato)-rhenium(IV). Comparison of Re^{I}, Re^{III}, Re^{IV}, and Re^{V} complexes with the salicylideneaminato ligand and reaction mechanisms, 613–8
Reactivity of the polyhydride complexes $[ReH_5(PPh_3)_3]$, $[ReH_3(dppe)_2]$ (dppe = $Ph_2PCH_2CH_2PPh_2$), $[ReH_3(PPh_3)_3L]$, and $[ReH_4(PPh_3)_3L]PF_6$ (L = MeCN or Bu^tNC) towards electrophiles and nucleophiles, 715–20
Rhenium carbonyl fluorides: preparation of $[Re(CO)_6][ReF_6]$ and some reactions of $[Re(CO)_5F{\cdot}ReF_5]$, 741–6
Reactions of co-ordinated ligands. Part 40. Synthesis, structure, and reactivity of dirhenium μ-allylidene complexes; crystal structures of $[Re_2(\mu\text{-}\sigma{:}\eta^3\text{-}CHCHCMe_2)(CO)_8]$ and $[Re_2(\mu\text{-}\sigma{:}\eta^3\text{-}CHCH\text{-}CMe_2)(CO)_9]$, 1313–8
Preparation and reactivity of halide-bridged binuclear carbonyl manganese and rhenium anions, $[M_2(\mu\text{-}X)(CO)_{8-2n}\{(EtO)_2POP(OEt)_2\}_n]^-$ (X = halide; M = Mn, n = 1 or 2; M = Re, n = 1). *X*-Ray crystal structure of $[Mn_2Au(\mu\text{-}Br)(CO)_6(\mu\text{-}tedip)(PPh_3)]$, 1551–6
Technetium(V) and rhenium(V) complexes with *N*-(2-mercaptophenyl)salicylideneiminate. Crystal structure of chloro(oxo)[*N*-(2-sulphidophenyl)salicylideneiminato-*NOS*]-technetium(V), 1693–700
Synthesis and reactions of $[(OC)_2Re(\mu\text{-}dppm)_2(\mu\text{-}Cl)_2Rh(\eta^2\text{-}C_2H_4)]$ (dppm = $Ph_2PCH_2PPh_2$): crystal structure of the 1:2 chlorobenzene solvate, 1763–8
Substitution reactions and luminescence spectra of the mixed complexes $[ReCl_xBr_{6-x}]^{2-}$, 1929–32
Carbon-13 nuclear magnetic resonance study of the properties of the unsaturated anion $[Re_3(\mu\text{-}H)_4(CO)_9(NCMe)]^-$ at low temperature, 2083–6
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S{=}C(NHPh)\text{-}S\}]{\cdot}[Re(CO)_2(PPh_3)_2\{O\text{–}C(NHPh){=}S\}]$, 2299–304
Chelate ring-opening of rhenium–diphosphine complexes by diazonium salts, 3067–72
Reactivity of rhenium(I) pseudo-allyl complexes with heterocumulenes such as RNCS (R = Me or Ph, *p*-MeC_6H_4) and PhNCO. Part 1. Crystal structure determination of two isostructural monothio- and dithio-carbamate rhenium(I) complexes co-crystallized as $[Re(CO)_2(PPh_3)_2\{S{=}C(NHPh)\text{–}S\}]{\cdot}[Re(CO)_2(PPh_3)_2\{O\text{–}C(NHPh){=}S\}]$ (1987, 2299), 3107–8

RHODABORANE
Pentamethylcyclopentadienylrhodaborane chemistry. Part 2. The reaction of $[6\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}6\text{-}RhB_9H_{13}]$ with dimethylphenylphosphine, and the characterization of $[5\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}5\text{-}RhB_9H_{11}\text{-}7\text{-}(PMe_2Ph)]$, $[2\text{-}(\eta^5\text{-}C_5Me_5)\text{-}closo\text{-}2\text{-}RhB_9H_7\text{-}3,10\text{-}(PMe_2Ph)_2]$, and $[2\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}2\text{-}RhB_8H_{10}\text{-}5\text{-}(PMe_2Ph)]$ by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 1431–44

RHODIUM
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 51. Alkylidyne(carbaborane)tungsten-gold and -rhodium complexes; crystal structures of $[AuW(\mu\text{-}CR)(CO)_2(PPh_3)(\eta^5\text{-}C_2B_9H_9Me_2)]$, $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2\text{-}(\eta^5\text{-}C_2B_9H_9Me_2)]$, and $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2\text{-}\{\eta^5\text{-}C_2B_9(C_7H_9)H_8Me_2\}]$ (R = C_6H_4Me-4), 61–72
Chemistry of platinum sulphido-complexes. Part 6. Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with the rhodium complexes $[(RhL_2Cl)_2]$ (L = CO or C_2H_4) and $[Rh(2,6\text{-}Me_2C_6H_3NC)_4]PF_6$. Crystal and molecular structures of $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(CO)_2]\text{-}[Rh(CO)_2Cl_2]{\cdot}C_3H_6O$ and $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4\text{-}(2,6\text{-}Me_2C_6H_3NC)_2]PF_6$, 335–40
Co-ordination of highly substituted alkenes to transition-metal complexes: preparation and crystal and molecular structures of the complexes $[\{RhCl(Ph_2PO_2CCH{=}CMe_2)\}_2]$ and $[\{RhCl(Ph_2PO_2CCH{=}CHCH{=}CHMe)\}_2]$, 443–50
Reduction–oxidation properties of organotransition-metal complexes. Part 25. The one-electron reduction of cationic cyclopentadienyl–cobalt and –rhodium nitrosyl derivatives, 467–72
Binuclear rhodium and iridium complexes containing pentamethylcyclopentadienyl and pyrazolate ligands, 639–46
Synthesis and characterisation of $[N(PPh_3)_2][RuRh_5(CO)_{16}]$ by *X*-ray crystallography and by natural-abundance carbon-13 nuclear magnetic resonance spectroscopy, 781–4
Preparation and reactions of bis(isocyanide)(η^5-indenyl)rhodium(I) complexes, 851–2
2-Pyridonate (Opy) rhodium(I) carbonyl complexes. Crystal structure of the unusual tetranuclear complex $[Rh_4(\mu\text{-}Opy)_4(\mu\text{-}CO)_2(CO)_4]$, 981–4
Reduction–oxidation properties of organotransition-metal complexes. Part 26. The synthesis of paramagnetic triazenido-and aryl(1-aryliminoethyl)amido-bridged dirhodium carbonyl derivatives containing the $[Rh_2]^{3+}$ core, 1403–8
Pentamethylcyclopentadienylrhodaborane chemistry. Part 2. The reaction of $[6\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}6\text{-}RhB_9H_{13}]$ with dimethylphenylphosphine, and the characterization of $[5\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}5\text{-}RhB_9H_{11}\text{-}7\text{-}(PMe_2Ph)]$, $[2\text{-}(\eta^5\text{-}C_5Me_5)\text{-}closo\text{-}2\text{-}RhB_9H_7\text{-}3,10\text{-}(PMe_2Ph)_2]$, and $[2\text{-}(\eta^5\text{-}C_5Me_5)\text{-}nido\text{-}2\text{-}RhB_8H_{10}\text{-}5\text{-}(PMe_2Ph)]$ by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 1431–44
Bimetallic systems. Part 20. Synthesis of Group 6 metal(0)–iridium(I) or –rhodium(I) complexes bridged by $Ph_2PCH_2PPh_2$, 1495–500
Synthesis and reactions of $[(OC)_2Re(\mu\text{-}dppm)_2(\mu\text{-}Cl)_2Rh(\eta^2\text{-}C_2H_4)]$ (dppm = $Ph_2PCH_2PPh_2$): crystal structure of the 1:2 chlorobenzene solvate, 1763–8
Activation energies for molecular tumbling and cyclopentadienyl rotation in $[M(\eta^5\text{-}C_5H_5)(\eta^4\text{-}cod)]$ (M = Rh or Ir; cod = cyclo-octa-1,5-diene) and the *X*-ray crystal structure of and bonding in $[Rh(\eta^5\text{-}C_5H_5)(\eta^4\text{-}cod)]$, 1947–52
Reactions of rhodium(V) hydrido–silyl complexes with butyl-lithium and with lithium triethylhydridoborate, 1963–6
Transformation of $[PPh_4][Fe_3RhC(CO)_{16}]$ under hydroformylation conditions: synthesis and crystal structure of $[PPh_4][Fe_3Rh_3C(CO)_{15}]$, 1983–8
Some chemistry of the heterobimetallic complexes $[(OC)_3Mo(\mu\text{-}dppm)_2Rh(CO)Cl]$ and $[(OC)_3Mo(\mu\text{-}dppm)_2M(CO)_2]PF_6$ (M = Rh or Ir, dppm = $Ph_2PCH_2PPh_2$). Structure of $[(OC)_3Mo(\mu\text{-}dppm)_2(\mu\text{-}C{\equiv}CMe)Rh(CO)]$, 2171–6
Pentamethylcyclopentadienylrhodaborane chemistry. Part 3. The reaction of $[\{Rh(\eta^5\text{-}C_5Me_5)Cl_2\}_2]$ with *closo*-$[B_{10}H_{10}]^{2-}$ and *arachno*-$B_{10}H_{12}$-6,9-$(PMe_2Ph)_2$, and the characterisation of some novel rhodaboranes by *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy, 2417–30
The cyclometallation of benzoic acid to give rhodium, iridium, and osmium *C,O*-benzoates. *X*-Ray structure determination of the dibenzoate $[(C_5Me_5)Rh(OOCPh)_2(H_2O)]$, 2459–66
Preparation and spectroscopic study of functionally substituted cyclopentadienides of thallium(I), potassium, rhodium(I), and iridium(I), 2615–20
The formation and characterisation of (η^2-ethene)hydrido(η^5-pentamethylcyclopentadienyl)(trisubstituted-silyl)rhodium complexes; intermediates in catalytic dehydrogenative silylation reactions, 2709–14
Rhodium(I)–iron(0) carbonyl complexes containing one bridging $Ph_2PCH_2PPh_2$ ligand, including the formation of a tetranuclear cluster from a heterobimetallic precursor: crystal structures of $[(OC)_4Fe(\mu\text{-}Ph_2PCH_2PPh_2)RhCl(CO)]$ and $[Fe_2Rh_2(\mu\text{-}Ph_2PCH_2PPh_2)_2(CO)_8]$, 2751–6
The chemistry of tetraphenylcyclopentadienone complexes of ruthenium and rhodium: the *X*-ray crystal structure of $[Ru\{\eta^5\text{-}C_5Ph_4OC(O)CH(OMe)Ph\}(CO)_2Cl]$, 2815–22
Synthesis and chemistry of *trans*-$[RhX(CO)L_2]$ (X = anionic ligand, L = tertiary phosphine), 2969–74
Dinuclear rhodium and iridium complexes of dicarboxyimidazolates; crystal structure of $[NBu_4][(cod)Rh(dci)Rh(cod)]{\cdot}2Pr^iOH$, 3003–8

RHODIUM-103
Reactions of rhodium(V) hydrido–silyl complexes with butyl-lithium and with lithium triethylhydridoborate, 1963–6

ROTATION
The triphenylphosphine cone angle and restricted rotation about the chromium–phosphorus bond in dicarbonyl(η^6-hexa-alkylbenzene)(triphenylphosphine)chromium(0) complexes. Crystal and molecular structure of dicarbonyl(η^6-hexa-n-propylbenzene)(triphenylphosphine)chromium(0), 1545–50
Activation energies for molecular tumbling and cyclopentadienyl rotation in $[M(\eta^5\text{-}C_5H_5)(\eta^4\text{-}cod)]$ (M = Rh or Ir; cod = cyclo-octa-1,5-diene) and the *X*-ray crystal structure of and bonding in $[Rh(\eta^5\text{-}C_5H_5)(\eta^4\text{-}cod)]$, 1947–52

RUTHENABORANE

Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [5,6,6-$(PPh_3)_3$-6-H-*nido*-6-RuB_9H_{12}] (1985, 2397), 275–6

RUTHENAUNDECABORATE

The preparation of [$Ru(NCMe)_6$][7-(η^6-C_6Me_6)-*nido*-7-$RuB_{10}H_{13}$]$_2$ and its characterization by single-crystal *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy: a triply hydrogen-bridged eleven-vertex *nido*-7-metallaundecaborane, 1169–74

RUTHENIUM

Phenyl complexes of ruthenium(II): competition between carbonyl substitution and formation of benzoyl complexes, and an *X*-ray study of a benzoyl complex, 27–34

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 50. Synthesis and crystal structures of the complexes [$RuW_3(\mu$-CO)(μ_3-η-CO)(μ-CPh)$_2$(μ_3-CPh)(CO)$_2$(η-C_5H_5)$_3$] and [$RuW_3(\mu$-CO)(μ_3-η-CO){μ-C(Ph)C(O)}(μ-CPh)(μ_3-CPh)(CO)($PMePh_2$)(η-C_5H_5)$_3$], 51–60

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 52. Synthesis and reactions of the compounds [$RuW(\mu$-CR)(CO)$_3$(η-C_5H_5)(η^5-$C_2B_9H_9Me_2$)] (R = C_6H_4Me-4 or Me); crystal structures of [$RuW(\mu$-CC_6H_4Me-4)(CO)$_3$(η-C_5H_5)(η^5-$C_2B_9H_9Me_2$)] and [$N(PPh_3)_2$][$RuW(\mu$-CC_6H_4Me-4)(μ-σ,η^5-$C_2B_9H_8Me_2$)(CO)$_3$(η-C_5H_5)], 73–80

High-yield syntheses of sulphido triruthenium carbonyl cluster complexes containing bis(diphenylphosphino)methane, 91–6

Reactivity of ruthenium(II) complexes with 1-hydroxymethyl- or 1-benzyl-3,5-dimethylpyrazole and similar functionalized bipyrazoles. *X*-Ray crystal structure of carbonylchlorohydrido(3,5-dimethylpyrazole-N^2)-bis(triphenylphosphine)ruthenium(II), 183–6

Synthesis of the alkoxo(hydrido)-clusters [$Ru_3(\mu$-H)(CO)$_{10}(\mu$-OR)] [R = Me, Et, Pr^n, or Bu^n] catalysed by dinuclear carbonyl iron complexes, 253–4

Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [5,6,6-$(PPh_3)_3$-6-H-*nido*-6-RuB_9H_{12}] (1985, 2397), 275–6

Reactions of metal carbonyl cluster complexes with multidentate phosphine ligands; the preparation and thermolysis of bis(dimethylphosphino)methane derivatives of [$Ru_3(CO)_{12}$], and the *X*-ray crystal structure of [$Ru_3(CO)_9$(H)($Me_2PCHPMe_2$)], 277–80

Thermal and photochemical behaviour of organotetraruthenium clusters: solution structures and dynamics of phosphine-substituted derivatives, 349–54

The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds [$M_2Ru_4H_2(CO)_{12}(PPh_3)_2$] (M = Cu, Ag, or Au), 379–90

Ruthenium(II) complexes of *closo* dodecaboranyl anions and the molecular structure of the *pileo* thirteen-vertex ruthenaborane [$(PPh_3)_2ClRuB_{12}H_{11}(NEt_3)$], 451–6

The synthesis and *X*-ray crystal structures of homoleptic tetrahedral aryls of osmium(IV) and of cyclohexyls of ruthenium(IV), osmium(IV), and chromium(IV), 557–62

Synthesis and characterisation of [$N(PPh_3)_2$][$RuRh_5(CO)_{16}$] by *X*-ray crystallography and by natural-abundance carbon-13 nuclear magnetic resonance spectroscopy, 781–4

Substitution reactions between penta-ammineaquaruthenium(II) and thiocyanato, protonated acetato, cyano, and oxalato ligands in aqueous acidic solutions, 957–60

Phosphinoalkylsilyl complexes. Part 7. Synthesis and reactivity of octahedral, bis(chelate)dicarbonyl complexes of ruthenium(II) and osmium(II). Conformational properties of the bis(chelate) framework and *X*-ray crystal structures of [$M(PPh_2CH_2CH_2SiMe_2)_2(CO)_2$] (M = Ru or Os), 969–74

The heteronuclear cluster chemistry of the Group 1B metals. Part 3. Synthesis, structures, and chemical properties of the trimetallic hexanuclear Group 1B metal cluster compounds [$MM'Ru_4(\mu_3$-H)$_2$(CO)$_{12}$(PPh_3)$_2$] (M = Cu, M′ = Ag or Au; M = Ag, M′ = Au). *X*-Ray crystal structure of the complex [$AgCuRu_4(\mu_3$-H)$_2$(CO)$_{12}$(PPh_3)$_2$]·CH_2Cl_2, 1001–8

Synthesis and stereochemical studies of the chiral ruthenium complexes [Ru(η-C_5H_5){(*S*)dpompyr-*PP*′}X] [dpompyr = *N*-diphenylphosphino-2-(diphenylphosphinoxymethyl)pyrrolidine, X = H or Cl]. Crystal structure of [(*S*)Ru(η-C_5H_5){(*S*)dpompyr-*PP*′}Cl], 1149–56

The preparation of [$Ru(NCMe)_6$][7-(η^6-C_6Me_6)-*nido*-7-$RuB_{10}H_{13}$]$_2$ and its characterization by single-crystal *X*-ray diffraction analysis and nuclear magnetic resonance spectroscopy: a triply hydrogen-bridged eleven-vertex *nido*-7-metallaundecaborane, 1169–74

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 55. Coupling of methylene and alkylidyne groups at a trimetal centre; crystal structures of [Ru_2W\{μ-σ:η-*trans*-CH=C(H)R}(μ-CO)(CO)$_3$(η-C_5H_5)$_2$(η-C_5Me_5)]·CH_2Cl_2 and [$Ru_2W(\mu_3$-$CCH_2R)(\mu$-CO)$_3$(CO)(η-C_5H_5)$_3$] (R = C_6H_4Me-4), 1201–8

Electronic *d*–*d* spectra and ligand-dissociation rates of pentacyanoruthenate(II) ions, 1267–70

The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds [$M_2Ru_4H_2(CO)_{12}(PPh_3)_2$] (M = Cu, Ag, or Au) (1987, 379), 1283–4

Synthesis, spectroscopic, and electrochemical properties of bis(2,2′-bipyridyl)ruthenium compounds of some pyridyl-1,2,4-triazoles, 1389–96

Conversion of uncharged methyl complexes of ruthenium(II) into a cationic acyl complex: a mechanistic study, 1713–6

Cationic ruthenium(II) phosphite complexes. Preparation and properties of monocarbonyl and isocyanide derivatives, 1813–6

The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of [$M_2Ru_3(\mu_3$-S)(μ-$Ph_2PCH_2PPh_2$)-(CO)$_9$] (M = Cu or Au), 1967–76

A trigonal bipyramidal pentaruthenium cluster, [$Ru_5(CO)_3(\mu_3$-CO)$_3$(η-C_5H_5)$_4$], 2087–8

Gold-197 Mössbauer spectra of some gold–ruthenium cluster compounds, 2333–6

Characterizations of the redox orbitals of mixed-ligand 2,2′-bipyridine–2-(2′-pyridyl)quinoline ruthenium(II) complexes, 2467–72

The mechanism of fluxionality of [Ru(η^6-C_8H_8)(η^4-C_7H_8)], [Ru(η^4-C_8H_8)(η^4-C_7H_8)L] [L = PEt_3 or $P(OMe)_3$], and [Os(η^6-C_8H_8)(η^4-C_8H_{12})] (C_8H_8 = cyclo-octatetraene, C_7H_8 = norborna-2,5-diene, and C_8H_{12} = cyclo-octa-1,5-diene), 2649–52

New decomposition pathways for formyl complexes of ruthenium(II), 2683–6

Organic chemistry of dinuclear metal centres. Part 9. Diruthenium complexes of buta-1,3-diene. *X*-Ray crystal structure of [$Ru_2(CO)_2(\mu$-*s*-*trans*-η^2:η^2-$CH_2CHCHCH_2$)(η-C_5H_5)$_2$], 2703–8

Reactions of tetra-μ-carboxylato-diruthenium(II,II) compounds. *X*-Ray crystal structures of $Ru_2(\mu$-$O_2CCF_3)_4$(thf)$_2$, $Ru_2(\mu$-$O_2CR)_4$-$(NO)_2$ (R = Et or CF_3), and {Na_2[$Ru_2(\mu$-$O_2CO)_4$]·$6H_2O$}$_n$, 2723–36

Organoruthenaborane chemistry. Part 5. Products of the reaction between *closo*-[$B_{10}H_{10}]^{2-}$ and [(η^6-C_6Me_6)$ClRuB_3H_8$]. Nuclear magnetic resonance studies and the crystal and molecular structure of [{(η^6-C_6Me_6)$_2Ru_2H_4$}$RuB_{10}H_8(OEt)_2$], 2781–8

The syntheses and crystal and molecular structures of some ten-vertex (MCB_8; M = Ru or Os) metallacarbaboranes: 9,9,9-(CO)(PPh_3)$_2$-*nido*-9,6-MCB_8H_{10}-5-PPh_3)] (M = Ru or Os) and [9,9,9-(η-C_5H_5)(H)(PPh_3)-*arachno*-9,6-$RuCB_8H_{12}$], 2793–802

Reactions of co-ordinated ligands. Part 42. The synthesis and reactions of cationic η^4-tetraphenylcyclobutadiene-ruthenium complexes; *X*-ray crystal structures of [Ru(NCMe)(η^4-C_4Ph_4)(η-C_5H_5)][BF_4] and [Ru{η^5-CH(Ph)C-(Ph)C(Ph)C(Ph)O}(η-C_5H_5)], 2803–14

The chemistry of tetraphenylcyclopentadienone complexes of ruthenium and rhodium: the *X*-ray crystal structure of [Ru{η^5-$C_5Ph_4OC(O)CH(OMe)Ph$}(CO)$_2$Cl], 2815–22

Synthesis using metal atoms of [RuH(η-C_6H_6)(PMe_3)(σ-C_6H_5)], [OsH(η-C_6H_6)(PMe_3)(σ-C_6H_5)], [MnH(η-C_6H_6)-(PMe_3)$_2$], and [Mn(η-C_5H_5)(PMe_3)$_3$]: arene ligand-exchange reactions of [RuH(η-C_6H_6)(PMe_3)(σ-C_6H_5)], 2823–30

Reactions of phosphine, arsine, and stibine with carbonylbis(triethylphosphine)iridium(I) halides. Part 1. Reactions in toluene; *X*-ray crystal structures of [Ir(CO)Cl$H(PEt_3)_2(AsH_2)$] and [Ir(CO)XH(PEt_3)$_2$(μ-

RUTHENIUM (contd)
$ZH_2)RuCl_2(\eta^6\text{-}MeC_6H_4CHMe_2\text{-}p)]$ (X = Br, Z = P; X = Cl, Z = As), 2831–8
Reactivity of a cluster-co-ordinated acyl. Reduction of a μ-acetyl ligand to acetaldehyde, 2839–42
Reactions of allene with triosmium and triruthenium clusters including coupling and oxidative addition: *X*-ray structure of the coupled allene compound $[Os_3(C_6H_8)(CO)_{10}]$, 2935–40
Preparation, isomerization, and reactions of hydride complexes of ruthenium(II), 2989–94
Synthesis, characterisation, and photochemical properties of a series of ruthenium containing metallopolymers based on poly-*N*-vinylimidazole, 3073–8
SALICYLATE
Complex formation of beryllium(II) with salicylate and hydroxide ions in 1 mol dm^{-3} sodium perchlorate aqueous solution at 25 °C, 1853–6
SCHIFF BASE
Complexes of ligands providing endogenous bridges. Part 1. The syntheses and crystal structures of barium and lead(II) complexes of macrocyclic Schiff bases derived from heterocyclic dicarbonyls and 1,*n*-diamino-*n'*-hydroxyalkanes (*n*,*n'* = 3,2; 4,2; or 5,3), 207–18
Complexes of ligands providing endogenous bridges. Part 2. The non-template synthesis of tetraimine Schiff base macrocycles derived from 1,3-diamino-2-hydroxypropane, 219–20
Crystal structure of *trans*-dicarbonylbis(dimethylphenylphosphine)-(*N*-phenylsalicylideneaminato)rhenium(I) and *mer*-trichloro-(dimethylphenylphosphine)(*N*-phenylsalicylideneaminato) rhenium(IV). Comparison of Re^{I}, Re^{III}, Re^{IV}, and Re^{V} complexes with the salicylideneaminato ligand and reaction mechanisms, 613–8
A comparative synthetic and structural study of μ-oxo and μ-sulphido derivatives of *N*,*N'*-ethylenebis(acetylacetoneiminato)-iron(III), including the synthesis and structural characterization of the parent compound, 709–14
Synthesis and characterization of technetium-(IV) and -(III) complexes with bidentate Schiff bases. *X*-Ray molecular structure of chloro(dimethylphenylphosphine)bis(*N*-phenylsalicylideneiminato)technetium(III), 867–72
Crystal structures of low- and high-spin iron(III) complexes with quadridentate Schiff bases, 1157–62
Water-soluble Schiff-base complexes of vanadyl(IV) and vanadium(III), 1279–82
Mixed oxidation state trinuclear cobalt complexes with bridging sulphito and Schiff-base ligands. Part 1. Preparation of the complexes $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}L)_2Co^{III}_2(ROH)_2]$ (L = Schiff base anion, R = alkyl) and structure determination of $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}\alpha,\alpha'\text{-}Me_2\text{-}salpd)_2Co^{III}_2(Pr^nOH)_2]\cdot 2Pr^nOH$, 1305–12
New spin-crossover iron(III) complexes with large hysteresis effects and time dependence of their magnetism, 1341–8
Reaction of elemental sulphur with cobalt(II)–Schiff-base complexes: synthesis of μ-disulphido and μ-tetrasulphido binuclear cobalt(III) complexes. Crystal structures of two binuclear cobalt(III)–Schiff-base complexes, 1367–76
Technetium(v) and rhenium(v) complexes with *N*-(2-mercaptophenyl)salicylideneiminate. Crystal structure of chloro(oxo)[*N*-(2-sulphidophenyl)salicylideneiminato-*NOS*]technetium(v), 1693–700
X-Ray structural study of [Fe(saltrien)]X (X = Br·$2H_2O$, BPh_4, or PF_6). Origin of unusual magnetic behaviour of the spin-crossover complex [Fe(saltrien)]PF_6, 1957–62
Synthesis and characterisation of some new manganese(II) and manganese(III) complexes of tetradentate Schiff-base ligands and their reaction with molecular oxygen, 2009–12
Transition metal Schiff-base complexes as ligands in tin chemistry. Part 1. A tin-119 Mössbauer spectroscopic investigation of the adducts $SnX_4\cdot ML$, $SnMe_2(NCS)_2\cdot ML$, and $SnRCl_3\cdot ML$ (X = halide; M = Cu^{II} or Ni^{II}; L = quadridentate Schiff-base ligand; R = phenyl or n-butyl), 2261–6
Complexes of ligands providing endogenous bridges. Part 4. Copper(II) complexes of macrocyclic Schiff bases derived from 2,6-diacetylpyridine and 1-*n*-diamino-*n'*-hydroxyalkanes (*n*,*n'* = 3,2; 4,2; and 5,3): synthesis, properties, and structures, 2519–30
Formation of a (4 + 4) Schiff-base macrocyclic ligand by a template rearrangement. Crystal and molecular structures of two tetranuclear manganese(II) complexes, 2555–62
Metallo-ligands containing terminal oxygen donors bonded to a manganese–carbonyl moiety, 2599–604
Structure and electrochemical properties of bis(pyridine)cobalt(III) complexes of *N*,*N'*-ethylenebis(salicylideneimine) homologues with an aryl substituent on the ethylene backbone. Crystal structures of $[Co(salphen)(py)_2]PF_6\cdot 0.5Et_2O$ and $[Co(salcppn)(py)_2]PF_6\cdot 2py$ (py = pyridine), 3055–60
SELENADODECABORANE
Reactions of heteroboranes. Synthesis of $[2,2\text{-}(PPh_3)_2\text{-}1,2\text{-}SePtB_{10}H_{10}]\cdot CH_2Cl_2$, its crystal and molecular structure and that of $SeB_{11}H_{11}$, 699–704
SELENIUM
Reactions of main group homopolynuclear cations with transition-metal carbonyls. Stabilisation of $\mu\text{-}Se_2^{2+}$ and $\eta^3\text{-}Te_3^{2+}$ dicationic ligands and crystal structures of $[Mo(Te_3)(CO)_4][SbF_6]_2$ and $[FeW(Se_2)(CO)_8][SbF_6]_2$, 2111–6
SILANE
Reaction of silicon difluoride with halogens: a reinvestigation, 1123–6
SILICATE
Synthesis and properties of gallium tetramethylammonium sodalite, $NMe_4GaSi_5O_{12}$, 623–6
SILICON
Structure and bonding in $(R_2SiX)_2$ molecules (X = O, NH, CH_2, or S; R = H), 271–4
Synthesis and properties of gallium tetramethylammonium sodalite, $NMe_4GaSi_5O_{12}$, 623–6
Electron diffraction study of the molecular structure of *N*-methylbis(trimethylsilyl)amine in the gas phase, 785–8
Reaction of silicon difluoride with halogens: a reinvestigation, 1123–6
Preparations, properties, and vibrational spectra of some (dimethylamino)halogenosilanes, 3029–34
Molecular structures of some (dimethylamino)halogenosilanes in the gas phase by electron diffraction and the crystal and molecular structures of mono-and di-chloro(dimethylamino)silane by *X*-ray diffraction at low temperatures, 3035–42
The structure of bis(dimethylamino)silane in the gas phase, determined by electron diffraction, 3061–6
SILVER
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au), 379–90
Thermodynamics of complex formation of silver with amines in dimethyl sulphoxide, 657–60
Bimetallic systems. Part 17. Synthesis of Group 6 metal(0)–copper(I), –silver(I), or –gold(I) complexes with two $Ph_2PCH_2PPh_2$ (dppm) bridging ligands: crystal structure of $[(OC)_2W(\mu\text{-}CO)(\mu\text{-}dppm)_2(\mu\text{-}Cl)Cu]\cdot CH_2Cl_2$, 769–76
The heteronuclear cluster chemistry of the Group 1B metals. Part 3. Synthesis, structures, and chemical properties of the trimetallic hexanuclear Group 1B metal cluster compounds $[MM'Ru_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]$ (M = Cu, M' = Ag or Au; M = Ag, M' = Au). *X*-Ray crystal structure of the complex $[AgCuRu_4(\mu_3\text{-}H)_2(CO)_{12}(PPh_3)_2]\cdot CH_2Cl_2$, 1001–8
The heteronuclear cluster chemistry of the Group 1B metals. Part 2. Synthesis, *X*-ray crystal structures, and dynamic behaviour of the bimetallic hexanuclear Group 1B metal cluster compounds $[M_2Ru_4H_2(CO)_{12}(PPh_3)_2]$ (M = Cu, Ag, or Au) (1987, 379), 1283–4
Crystal and molecular structure of $[\{Ag[S_2P(OEt)_2]\cdot PPh_3\}_2]$, 1569–72
Metal(II) hexafluoroantimonates: preparation and characterization of $MF_2\cdot 2SbF_5$ (M = Mg, Ni, Zn, Fe, Co, Cu, Cr, Ag, Cd, or Pb) and the *X*-ray structure determination of $AgF_2\cdot 2SbF_5$, 2379–84
σ-Alkynyl complexes of manganese(I) as η^2-bonding ligands for Group 1B metal–ligand fragments. *X*-Ray crystal structure of $[Mn_2Cu(\mu\text{-}CCBu^t)_2(CO)_6(dppe)_2]PF_6\cdot 0.5CH_2Cl_2$, 2867–74
Syntheses and crystal structures of complexes $[M_2R_2]$, [M = Cu, Ag, or Au; R = 2-$C(SiMe_3)_2C_5H_4N$] and $[Cu_4R'_4]$[R' = 2-$CH(SiMe_3)C_5H_4N$]; electrochemical generation of $[Cu_2R_2]^{2+}$, 3085–92
SILYL
Reactions of rhodium(v) hydrido–silyl complexes with butyl-lithium and with lithium triethylhydridoborate, 1963–6

SODALITE
Synthesis and properties of gallium tetramethylammonium sodalite, $NMe_4GaSi_5O_{12}$, 623–6
SOLID SOLUTION
Studies on alkaline-earth metal sulphites. Part 9. Solid solutions of $Ca_3(SO_4)_2SO_3{\cdot}12H_2O$ in $Ca_3(SO_3)_2SO_4{\cdot}12H_2O$: structure, stability, and dehydration products with amines in dimethyl sulphoxide, 657–60
Structure and basicity. Part 10. Conductometric and potentiometric titrations of some cyclotriphosphazatrienes in non-aqueous media. The conductivities of alkali-metal acetates on titration with perchloric acid in acetic acid, 925–30
SPIN DENSITY
Charge density in the $CoCl_4^{2-}$ ion: a comparison with spin density and theoretical calculations, 1737–46
SPIRO-PYRAZOLE
Reaction of an enolisable diazo-compound with the electron-rich stannylene $Sn[N(SiMe_3)_2]_2$: preparation of a novel spiro[1,3,4,2-oxadiazastannine-6,4′-pyrazole] derivative, 509–12
STABILITY
Co-ordination abilities of thioamide analogues of amino acids and peptides. Copper(II) and nickel(II) complexes, 1–4
Equilibria in aqueous solutions of some iron(II) complexes, 255–8
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution, 435–42
Interactions of the trimethyltin(IV) cation with carboxylic acids, amino acids, and related ligands, 563–6
Thermodynamics of complex formation of silver with amines in dimethyl sulphoxide, 657–60
A thermodynamic and spectroscopic study of the proton and copper(II) complexes of L-prolyl-L-histidine, D-prolyl-L-histidine, L-histidyl-L-histidine, and D-histidyl-L-histidine, 661–6
Uranyl(VI) complexes of 3-aminopropanoic acid and 4-aminobutanoic acid in aqueous solution: a potentiometric and calorimetric study, 695–8
The chemistry of vitamin B_{12}. Part 27. The cage phosphite ester 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane and the anionic dimethyl phosphite as ligands for cobalt(III) corrinoids, 761–8
Stability constants for hypochloritoborate and hypobromitoborate complex ions in aqueous solution, 943–6
Aluminium speciation studies in biological fluids. A new investigation of aluminium hydroxide equilibria under physiological conditions, 1145–8
Copper(II) complexes of tyrosine-containing tripeptides, 1263–6
Complex formation equilibria between 2-amino-*N*-hydroxyacetamide and 2-amino-*N*-hydroxypentanamide and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solutions (1986, 2587), 1283–4
Design of transition metal chelates with biological activity. Potentiometric study of complex formation equilibria between 2-amino-*N*-hydroxy-3-(*p*-hydroxyphenyl)propanamide and nickel(II), copper(II), and hydrogen ions in aqueous solution, 1409–14
Pattern of transition-metal chelates with biological activity. Complex formation equilibria between α-amino-*N*-hydroxy-1*H*-imidazole-4-propanamide and cobalt(II), nickel(II), and hydrogen ions in aqueous solution (1987, 435), 1583–4
Tungsten(VI) equilibria: a potentiometric and calorimetric investigation, 1701–6
Studies on transition-metal–peptide complexes. Part 13. Copper(II) and nickel(II) complexes of amino acids and peptides containing a thioether group, 1717–20
Correlation between thermal stability and molecular and crystal structure in a series of potassium–dibenzo-18-crown-6 solid compounds, 1779–84
Complex formation of beryllium(II) with salicylate and hydroxide ions in 1 mol dm^{-3} sodium perchlorate aqueous solution at 25 °C, 1853–6
The hydrolysis of metal ions. Part 10. Kinetic and equilibrium measurements of molybdenum(VI), 2149–58
Beryllium(II) hydrolysis in 3.0 mol dm^{-3} perchlorate, 2431–8
Studies of metal carbonate equilibria. Part 15. The beryllium(II)–water–carbon dioxide(g) system in acidic 3.0 mol dm^{-3} perchlorate media, 2439–44
Studies of metal carbonate equilibria. Part 16. The beryllium(II)–water–carbon dioxide(g) system in neutral-to-alkaline 3.0 mol dm^{-3} perchlorate media at 25 °C, 2445–50
Complexes formed between zinc(II) and diethylenetriamine: a carbon-13 nuclear magnetic resonance study, 2889–96
Carbon-13 nuclear magnetic resonance study of the zinc(II) complexes of 3,7-diazanonane-1,9-diamine and 1,2-diaminoethane, 2897–904
The effect of increase in chelate ring size beyond six-membered on the metal ion size selectivity patterns of tetra-aza macrocycles, 2911–6
STACKING
The ring-stacking principle in organolithium chemistry: its development through the isolation and crystal structures of hexameric iminolithium clusters $(RR'C{=}NLi)_6$ (R′ = Ph, R = Bu^t or Me_2N; R = R′ = Me_2N or Bu^t), 1071–82
Metal ion-bridged intramolecular stacking interaction between the tryptophyl residue and the aromatic heterocyclic amine within the ternary complex (1,10-phenanthroline)(L-tryptophanato-*O*,*N*)copper(II) perchlorate–water (1/2.5): an *X*-ray study, 2017–22
STEREOCHEMISTRY
Thermal investigation and stereochemical studies of some cyclic diamine complexes of nickel(II), zinc(II), and cadmium(II) in the solid state, 2089–94
STEREOSELECTIVITY
Application of ultraviolet photoelectron spectroscopy to molecular properties. Part 27. Stereochemistry of ligand exchange reactions of titanocene derivatives, 107–10
A thermodynamic and spectroscopic study of the proton and copper(II) complexes of L-prolyl-L-histidine, D-prolyl-L-histidine, L-histidyl-L-histidine, and D-histidyl-L-histidine, 661–6
Synthesis and stereochemical studies of the chiral ruthenium complexes $[Ru(\eta\text{-}C_5H_5)\{(S)dpompyr\text{-}PP'\}X]$ [dpompyr = *N*-diphenylphosphino-2-(diphenylphosphinoxymethyl)pyrrolidine, X = H or Cl]. Crystal structure of $[(S)Ru(\eta\text{-}C_5H_5)\{(S)dpompyr\text{-}PP'\}Cl]$, 1149–56
STIBINE
The vibrational spectrum of tricarbonyl(trichlorostibine)nickel(0), $[Ni(CO)_3(SbCl_3)]$, 681–2
Reactions of phosphine, arsine, and stibine with carbonylbis(triethylphosphine)iridium(I) halides. Part 1. Reactions in toluene; *X*-ray crystal structures of $[Ir(CO)ClH(PEt_3)_2(AsH_2)]$ and $[Ir(CO)XH(PEt_3)_2(\mu\text{-}ZH_2)RuCl_2(\eta^6\text{-}MeC_6H_4CHMe_2\text{-}p)]$ (X = Br, Z = P; X = Cl, Z = As), 2831–8
STRONTIUM
Magnetic exchange interactions in perovskite solid solutions. Part 7. A study of the non-stoicheiometric phase $SrCoO_{3-y}$ by ^{57}Fe Mössbauer spectroscopy, 667–72
A structural comparison of the two polymorphs of $Sr_2Co_2O_5$ by cobalt *K*-edge extended *X*-ray absorption fine structure spectroscopy, 2359–64
SUBSTITUTION
Substitution reactions between penta-ammineaquaruthenium(II) and thiocyanato, protonated acetato, cyano, and oxalato ligands in aqueous acidic solutions, 957–60
The relative nucleophilic efficiency of water and the azide and thiocyanate ions in the aqueous substitution of chloropentacyanocobaltate(III), 1059–62
Photochemical preparation and reactivity of platinum(II) complexes with oxygen-donor ligands. Description and synthesis of di-μ-chloro-dichlorobis(substituted pyridine)diplatinum(II), 1687–92
Preparation and properties of the tungsten(V) aqua dimer $[W_2O_4(H_2O)_6]^{2+}$: kinetic studies on the 1:1 reaction with NCS^-, 2293–8
The relative nucleophilic efficiency of water and dimethyl sulphoxide in the aqueous substitution of aquapentacyanocobaltate(III) and chloropentacyanocobaltate(III), 2311–6
SULPHATE
Studies on alkaline-earth metal sulphites. Part 9. Solid solutions of $Ca_3(SO_4)_2SO_3{\cdot}12H_2O$ in $Ca_3(SO_3)_2SO_4{\cdot}12H_2O$: structure, stability, and dehydration products, 235–40
Crystal and molecular structure of chromium(II) sulphate pentahydrate and single-crystal electronic spectra and bonding of $CrSO_4{\cdot}5H_2O$, $CuSO_4{\cdot}5H_2O$, and $CuSO_4{\cdot}5D_2O$, 1817–22
SULPHIDE
High-yield syntheses of sulphido triruthenium carbonyl cluster complexes containing bis(diphenylphosphino)methane, 91–6
Matrix isolation studies of monomeric boron trisulphide, 267–70
Chemistry of platinum sulphido-complexes. Part 6. Reactions of

SULPHIDE (contd)
$[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with the rhodium complexes $[(RhL_2Cl)_2]$ (L = CO or C_2H_4) and $[Rh(2,6\text{-}Me_2C_6H_3NC)_4]PF_6$. Crystal and molecular structures of $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4(CO)_2]\text{-}[Rh(CO)_2Cl_2]\text{-}C_3H_6O$ and $[Pt_2Rh(\mu\text{-}S)_2(PPh_3)_4\text{-}(2,6\text{-}Me_2C_6H_3NC)_2]PF_6$, 335–40
A comparative synthetic and structural study of μ-oxo and μ-sulphido derivatives of *N,N'*-ethylenebis(acetylacetoneiminato)-iron(III), including the synthesis and structural characterization of the parent compound, 709–14
Redox behaviour of the iron–sulphur cluster $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{2+}$. Synthesis and crystal structure of the new paramagnetic monopositive species $[Fe_6(\mu_3\text{-}S)_8(PEt_3)_6]^{+}$ as its $[PF_6]^{-}$ salt, 831–6
Dimeric tungsten(V) compounds containing $[(S)W(\mu\text{-}S)_2W(S)]^{2+}$. Syntheses and structures of di-μ-sulphido-bis[(diethyl dithiophosphato-*SS'*)sulphidotungsten(V)] and di-μ-sulphido-bis[(*N,N*-diethyldithiocarbamato-*SS'*)sulphidotungsten(V)], 1163–8
Reaction of elemental sulphur with cobalt(II)–Schiff-base complexes: synthesis of μ-disulphido and μ-tetrasulphido binuclear cobalt(III) complexes. Crystal structures of two binuclear cobalt(III)–Schiff-base complexes, 1367–76
Kinetic study of the homogeneous methanolysis of tetraphosphorus decasulphide, 1897–900
The heteronuclear cluster chemistry of the Group 1B metals. Part 4. Synthesis, structures, and dynamic behaviour of Group 1B metal–ruthenium cluster compounds containing sulphur ligands. *X*-Ray crystal structures of $[M_2Ru_3(\mu_3\text{-}S)(\mu\text{-}Ph_2PCH_2PPh_2)(CO)_9]$ (M = Cu or Au), 1967–76
Reaction of elemental sulphur with bis(pentamethylcyclopentadienyl)vanadium derivatives: disulphido and μ-sulphido complexes. Crystal structures of $[V(\eta\text{-}C_5Me_5)_2(S_2)]$ and $[V_2(\eta\text{-}C_5Me_5)_2Cl_2(\mu\text{-}S)_2]$, 2099–104
A two-dimensional nuclear magnetic resonance exchange study of the fluxionality of $[PtIMe_3(MeSCH_2CH_2SEt)]$, 2695–702

SULPHITE
Studies on alkaline-earth metal sulphites. Part 9. Solid solutions of $Ca_3(SO_4)_2SO_3\cdot12H_2O$ in $Ca_3(SO_3)_2SO_4\cdot12H_2O$: structure, stability, and dehydration products, 235–40
A kinetic and mechanistic study of the formation of a diaryl ditelluride in solution by the sulphite reduction of the aryltellurium trichloride, 985–90
Mixed oxidation state trinuclear cobalt complexes with bridging sulphito and Schiff-base ligands. Part 1. Preparation of the complexes $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}L)_2Co^{III}_2(ROH)_2]$ (L = Schiff base anion, R = alkyl) and structure determination of $[Co^{II}(\mu\text{-}SO_3)_2(\mu\text{-}\alpha,\alpha'\text{-}Me_2\text{-}salpd)_2Co^{III}_2(Pr^nOH)_2]\cdot2Pr^nOH$, 1305–12
Reactions in mixed non-aqueous solution containing sulphur dioxide. Part 9. Mechanisms of dissolution of metals, oxides, and sulphites, 2995–8

SULPHONATE
Complexes of the platinum metals. Part 32. Synthesis and chemistry of some iridium sulphonate derivatives, 947–52

SULPHUR
The radical cations of sulphur ($S_8^{\cdot+}$) and tetrasulphur tetranitride ($S_4N_4^{\cdot+}$): a radiation–electron spin resonance study, 729–32
Some investigations of $S_5N_5^{+}$ compounds; *X*-ray crystal structure of $[S_5N_5]Cl$, 915–24
Reaction of elemental sulphur with cobalt(II)–Schiff-base complexes: synthesis of μ-disulphido and μ-tetrasulphido binuclear cobalt(III) complexes. Crystal structures of two binuclear cobalt(III)–Schiff-base complexes, 1367–76
Phosphorus-31 nuclear magnetic resonance parameters for compounds with the α-tetraphosphorus trisulphide skeleton; substituent constants for an inorganic substrate, 1515–20
Reactions of $(NSCl)_3$ with Lewis acids and sulphur: a convenient preparation of $[S_2N][SbCl_6]$, 1565–8
Reaction of elemental sulphur with bis(pentamethylcyclopentadienyl)vanadium derivatives: disulphido and μ-sulphido complexes. Crystal structures of $[V(\eta\text{-}C_5Me_5)_2(S_2)]$ and $[V_2(\eta\text{-}C_5Me_5)_2Cl_2(\mu\text{-}S)_2]$, 2099–104
Attempted synthesis of osmium(VI) and iridium(VI) thiofluorides; the preparation of $OsF_5\cdot SF_4$ and $IrF_5\cdot SF_4$, 2285–8

SULPHUR DIOXIDE
The synthesis of silica-supported phosphine and phosphine oxide complexes of manganese(II) iodide and their reaction with sulphur dioxide, 1415–8
A comparison of the efficiencies of polymer-supported phosphine and phosphine oxide complexes of manganese(II), $MnX_2(C_6H_4PPh_2)$ and $MnX_2\{C_6H_4P(O)Ph_2\}_n$ (n = 2 or 4; X = Cl, Br, or I), for the absorption of sulphur dioxide, 2959–62
Reactions in mixed non-aqueous solution containing sulphur dioxide. Part 9. Mechanisms of dissolution of metals, oxides, and sulphites, 2995–8

SUPPORTED MATERIAL
The synthesis of silica-supported phosphine and phosphine oxide complexes of manganese(II) iodide and their reaction with sulphur dioxide, 1415–8
A comparison of the efficiencies of polymer-supported phosphine and phosphine oxide complexes of manganese(II), $MnX_2(C_6H_4PPh_2)$ and $MnX_2\{C_6H_4P(O)Ph_2\}_n$ (n = 2 or 4; X = Cl, Br, or I), for the absorption of sulphur dioxide, 2959–62

TANTALUM
The synthesis and structure of Group 5 metal alkyl and alkylidene complexes containing 2,6-dialkylphenoxide ligands: *X*-ray crystal structures of $[Ta(OC_6H_3Me_2\text{-}2,6)_2(CH_2Ph)_3]$, $[Ta(OC_6H_3Me_2\text{-}2,6)_4Me]$, and $[Ta(OC_6H_3Bu^t_2\text{-}2,6)_2(=CHSiMe_3)(CH_2SiMe_3)]$, 155–62
Electronic absorption spectra and photochemical reactivity of Group 5 metal alkyl compounds: photochemical α-hydrogen abstraction, 163–8
Trimethylphosphine complexes of niobium(IV) and tantalum(IV). Crystal, molecular, and electronic structures of $[M_2Cl_4(\mu\text{-}Cl)_4(PMe_3)_4]$ (M = Nb or Ta), 307–14

TECHNETIUM
Synthesis and characterization of technetium-(IV) and -(III) complexes with bidentate Schiff bases. *X*-Ray molecular structure of chloro(dimethylphenylphosphine)bis(*N*-phenylsalicylideneiminato)technetium(III), 867–72
Technetium(V) and rhenium(V) complexes with *N*-(2-mercaptophenyl)salicylideneiminate. Crystal structure of chloro(oxo)[*N*-(2-sulphidophenyl)salicylideneiminato-*NOS*]technetium(V), 1693–700
An electron spin resonance study of the equilibrium between tetrahalogeno-and pentahalogeno-nitridotechnetate(VI) ions in solution, 1721–6
Preparation and crystal structures of bis(tetraphenylarsonium) tris(oxalato)technetate(IV), and tetraphenylarsonium tris(benzene-1,2-dithiolato)technetate(V): octahedral *versus* trigonal-prismatic geometry for tris-bidentate complexes of technetium, 2305–10

TELLURIUM
A kinetic and mechanistic study of the formation of a diaryl ditelluride in solution by the sulphite reduction of the aryltellurium trichloride, 985–90
Reactions of main group homopolynuclear cations with transition-metal carbonyls. Stabilisation of $\mu\text{-}Se_2^{2+}$ and $\eta^3\text{-}Te_3^{2+}$ dicationic ligands and crystal structures of $[Mo(Te_3)(CO)_4][SbF_6]_2$ and $[FeW(Se_2)(CO)_8][SbF_6]_2$, 2111–6

TELLUROPHOSPHORANE
Phosphorus-tellurium compounds. Part 3. Cyclopentadienyliron tellurophosphorane complexes: tellurium-bridged iron phosphine compounds?, 541–4

TEMPLATE
Formation of a (4 + 4) Schiff-base macrocyclic ligand by a template rearrangement. Crystal and molecular structures of two tetranuclear manganese(II) complexes, 2555–62

TERPYRIDINE
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 5. Unsubstituted fifteen-membered-ring macrocycles containing 2,2′:6′,2″-terpyridine, 5–8
Magnetostructural correlations in parallel square-planar chloride bridged copper(II) dimers: structure, dynamic nuclear magnetic resonance study, and magnetic properties of $[Cu_2(terpy)_2Cl_2][PF_6]_2$, 285–92

TETRA-AZABICYCLOHEPTADECATRIENE
Studies of pendant-arm macrocyclic ligands. Part 5. Synthesis of two pyridine-containing penta-aza macrocycles with single pyrrolidinyl pendant arms and characterisation of their nickel(II) and copper(II) complexes. Crystal structure of perchlorato{7-[2′-(1″-pyrrolidinyl)ethyl]-3,7,11,17-tetra-azabicyclo[11.3.1]-heptadeca-1(17),13,15-triene}nickel(II) perchlorate, 545–50
Synthesis of pyridine-containing tetra-aza macrocycles: 3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), its 3,11-dibenzyl (L^2) and 3,7,11-tribenzyl (L^3) derivatives, and their

TETRA-AZABICYCLOHEPTADECATRIENE (contd)
nickel(II), copper(II), and zinc(II) complexes: crystal structures of L^2·HCl and $[Ni(L^2)Cl]ClO_4·H_2O$, 889–94
Studies of the pyridine-containing tetra-azamacrocycle 3,7,11-trimethyl-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1) and its complexes with Ni^{2+}, Cu^{2+}, and Zn^{2+}. Crystal structures of the unsymmetric isomer of $[Ni(L^1)][ClO_4]_2$, and the five-co-ordinate complexes $[Ni(L^1)(dmso)][ClO_4]_2$ (dmso = dimethyl sulphoxide) and $[Ni(L^1)Cl]ClO_4$, 1107–14

TETRA-AZACYCLODODECANE
Amino-acid complexation to a cobalt(III) complex of optically-active (2*R*,5*R*,8*R*,11*R*)-2,5,8,11-tetraethyl-1,4,7,10-tetra-azacyclododecane (L). Preparation, characterization, and structure determinations of $[Co^{III}\{(R)$- or (S)-alaO$\}$L]Br-$[ClO_4]·H_2O$, 721–8

TETRA-AZACYCLO-OCTACOSAOCTAENE
Binucleating N_6 24- and 26-membered macrocyclic ligands. Part 1. Dilead complexes: *X*-ray crystal structure determination of a macrocyclic dilead complex containing nitrogen-only bridging thiocyanate, 123–8

TETRA-AZACYCLOTETRADECANE
Preparation and structure of nickel(II) complexes of N-substituted tetra-aza macrocycles: molecular structure of [(1*RS*,4*SR*,5*RS*,8*SR*,11*RS*,12*SR*)-1,5,8,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane]nickel(II) diperchlorate, 3093–8

TETRACYANOETHYLENE
The co-ordination of small molecules by manganese(II) phosphine complexes. The reaction of some $[MnX_2L]$ (X = Cl, Br, or I; L = PPr^n_3, PMe_2Ph, or PPh_3) complexes with tetracyanoethylene, 1785–8
Dienophile cycloaddition to cycloheptatriene and related complexes of tricarbonyliron. *X*-Ray crystal structures of $[Fe(CO)_3(\eta^4$-$C_7H_7CN)]·C_2(CN)_4$ and $[Fe(CO)_3(\eta^4$-C_7H_7-$C_7H_7)]$, 2927–34

TETRAHYDROTHIOPHENE
Structure and solvation of mercury(II) iodide and bromide in tetrahydrothiophene solution, 2411–6

TETRATHIABICYCLONONENETHIONE
Complexes of 2,5,7,9-tetrathiabicyclo[4.3.0]non-1(6)-ene-8-thione (L^1). Part 1. Synthesis and antimony-121 Mössbauer spectroscopic characterisation of $SbCl_3(L^1)_{1.5}$, crystal and molecular structures of $SbCl_3(L^1)_{1.5}$ and uncomplexed L^1, 1605–10

THALLIUM
The chemistry of trichloro[tris(trimethylsilyl)methyl] and trichloro[tris(dimethylphenylsilyl)methyl] complexes of gallium, indium, and thallium. Crystal and molecular structures of $[Li(thf)_2(\mu$-$Cl)_2GaCl\{C(SiMe_2Ph)_3\}]$·thf, $[Li(thf)_3(\mu$-$Cl)InCl_2\{C(SiMe_3)_3\}]$, and $[\{(Me_3Si)_3C\}In(\mu$-$Cl)_2\{\mu$-$Fe(CO)_4\}In\{C(SiMe_3)_3\}]$ (thf = tetrahydrofuran), 747–56
Transition-metal mediated asymmetric synthesis. Part 5. Alternative methods for the formation of tricarbonyl(η^5-cyclohexadienyl)-iron(1+) salts, 885–8
Preparation and spectroscopic study of functionally substituted cyclopentadienides of thallium(I), potassium, rhodium(I), and iridium(I), 2615–20

THERAPY
Regioselective synthesis of $[1\text{-}B_{10}H_9(SH)]^{2-}$ and $[2\text{-}B_{10}H_9(SH)]^{2-}$: potential agents for boron–neutron capture therapy of brain tumours, 1953–6

THERMOCHEMISTRY
Thermal investigation and stereochemical studies of some cyclic diamine complexes of nickel(II), zinc(II), and cadmium(II) in the solid state, 2089–94

THERMODYNAMICS
Thermodynamics of complex formation of silver with amines in dimethyl sulphoxide, 657–60
Uranyl(VI) complexes of 3-aminopropanoic acid and 4-aminobutanoic acid in aqueous solution: a potentiometric and calorimetric study, 695–8
Tungsten(VI) equilibria: a potentiometric and calorimetric investigation, 1701–6
Mean phosphorus– and arsenic–sulphur bond enthalpies in diethyldithiocarbamate chelate complexes, 2955–8

THIASELENAPENTANE
Heteronuclear 1,3-intramolecular metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of 2-thia-4-selenapentane. A dynamic nuclear magnetic resonance investigation, 757–60

THIOAMIDE
Co-ordination abilities of thioamide analogues of amino acids and peptides. Copper(II) and nickel(II) complexes, 1–4

THIOCYANATE
Tin-119 nuclear magnetic resonance studies of some pseudohalogeno-derivatives of $[SnX_6]^{2-}$ (X = Cl or Br), 315–8
The relative nucleophilic efficiency of water and the azide and thiocyanate ions in the aqueous substitution of chloropentacyanocobaltate(III), 1059–62
Preparation and properties of the tungsten(V) aqua dimer $[W_2O_4(H_2O)_6]^{2+}$: kinetic studies on the 1:1 reaction with NCS^-, 2293–8
Studies involving mixed donor macrocyclic ligands. Comparative *X*-ray diffraction, infrared, and fast atom bombardment mass spectrometric studies of the nickel thiocyanate complexes of three O_2N_3-donor macrocycles, 2537–42
Formation of a (4 + 4) Schiff-base macrocyclic ligand by a template rearrangement. Crystal and molecular structures of two tetranuclear manganese(II) complexes, 2555–62
Binuclear copper(I) complexes of a 24-membered macrocycle (L^1) derived from 2,5-diformylthiophene and 3-thiapentane-1,5-diamine. Crystal and molecular structure of $[Cu_2L^1(NCS)][ClO_4]$; a unique example of a single end-to-end bridging thiocyanate between copper(I) atoms, 2975–80

THIOLATE
Syntheses, spectroscopic, and electrochemical properties of dinuclear copper(II) complexes bridged by a single thiolate sulphur atom; *X*-ray crystal structures of $[Cu_2(L^1)_2(SC_6H_4Cl$-$4)][PF_6]$ and $[Cu_2(L^2)_2(SC_6H_4CH_3$-$4)][ClO_4]$, 241–8
The preparation, structure, and reactions of the mononuclear thiolate–hydride complexes $[MoH(SR)(dppe)_2]$ (R = bulkyl alkyl or aryl group, dppe = $Ph_2PCH_2CH_2PPh_2$); *X*-ray structure determination of *cis*-$[MoH(SC_6H_2Pr^i_3$-2,4,6$)(dppe)_2]$, 1115–22
Systematic cluster build up by the reaction of $[Os_3H(X)(CO)_9(MeCN)]$ (X = SR or NHMe; R = Me, Et, or Ph) with $[OsH_2(CO)_4]$, 1289–92
Preparation and crystal structures of $[PPh_4]_2[M_2(SPh)_6]$ (M = Zn or Cd), 1577–80
Comments on the attempts by Wickens and Abrams to synthesize manganese(II) phosphine complexes: a comment, 1581–2
Substituted osmium clusters containing labile ligands. The preparation, characterisation, and some reactions of $[Os_3H(OR)(CO)_9(MeCN)]$ R = H, Me, Et, or Ph) and the isomers of $[Os_3H(SR)(CO)_9(MeCN)]$ (R = Me, Et, or Ph); the crystal and molecular structure of $[Os_3H(SPh)(CO)_9(PEt_3)]$, 1623–30
Polymeric thiolate complexes of Group 12 metals. Crystal and molecular structures of *catena*-[μ-(3-dimethylammonio-1-propanethiolate)]-dichlorocadmium(II) and bis[3-(dimethylammonio)propyl] disulphide tetrabromocadmate(II), 2391–6

THIOMOLYBDATE
Raman and resonance-Raman studies of platinum group tetrathiomolybdates and tetrathiotungstates, 1535–44

THIOPHENOL
High-yield syntheses of sulphido triruthenium carbonyl cluster complexes containing bis(diphenylphosphino)methane, 91–6

THIOPHOSPHINATE
Preparation of isocyanide and mixed dinitrogen–isocyanide complexes of rhenium(I) from reactions of *trans*-$[ReCl(N_2)(PMe_2Ph)_4]$, *mer*-$[Re(S_2PPh_2)(N_2)(PMe_2Ph)_3]$, or *mer*-$[Re(S_2CNEt_2)(N_2)(PMe_2Ph)_3]$ with methyl isocyanide; crystal structure of *mer*-$[Re(S_2PPh_2)(N_2)(CNMe)(PMe_2Ph)_3]$, 319–26

THIOSEMICARBAZONE
Thiosemicarbazones as co-ordinating agents. Part 2. Synthesis, spectroscopic characterization, and *X*-ray structure of aquachloro(pyridoxal thiosemicarbazone)manganese(II) chloride and aqua(pyridoxal thiosemicarbazonato)copper(II) chloride monohydrate, 227–34
Synthetic, spectroscopic, and *X*-ray crystallographic studies on binuclear copper(II) complexes with a tridentate NNS-bonding 2-formylpyridine thiosemicarbazone ligand. The characterization of both neutral and deprotonated co-ordinated ligand structures, 493–500
Synthetic, spectroscopic, and *X*-ray crystallographic studies on binuclear copper(II) complexes with a tridentate NNS-bonding 2-formylpyridine thiosemicarbazone ligand. The characterization of

THIOSEMICARBAZONE (contd)
both neutral and deprotonated co-ordinated ligand structures (1987, 493), 1583–4
THIOTOLUOYL
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 57. Reactions of iron– molybdenum complexes with oxygen and sulphur; crystal structure of $[FeMo(\mu\text{-}\eta^2\text{-}SCC_6H_4Me\text{-}4)(CO)_5(\eta\text{-}C_5H_5)]$, 1215–20
THIOTUNGSTATE
Raman and resonance-Raman studies of platinum group tetrathiomolybdates and tetrathiotungstates, 1535–44
TIN
Organotin biocides. Part 8. The crystal structure of triphenyltin formate and a comparative variable-temperature tin-119 Mössbauer spectroscopic study of organotin formates and acetates, 101–6
The first authenticated example of geometrical isomers in organotin(IV) co-ordination chemistry. The crystal and molecular structures of octahedral *cis*- and *trans*-SnR_2 isomeric adducts of dichlorobis(4-chlorophenyl)tin(IV) with 4,4′-dimethyl-2,2′-bipyridyl, 129–38
Tin-119 nuclear magnetic resonance studies of some pseudohalogeno-derivatives of $[SnX_6]^{2-}$ (X = Cl or Br), 315–8
Reaction of an enolisable diazo-compound with the electron-rich stannylene $Sn[N(SiMe_3)_2]_2$: preparation of a novel spiro[1,3,4,2-oxadiazastannine-6,4′-pyrazole] derivative, 509–12
Interactions of the trimethyltin(IV) cation with carboxylic acids, amino acids, and related ligands, 563–6
The configuration and lattice dynamics of complexes of dialkyltin(IV) with adenosine 5′-monophosphate and phenyl phosphates, 789–94
Crystal structure and mass spectrometry of dichlorodimethylbis[2(1*H*)-pyridinethione-*S*]tin(IV), 815–8
Preparation and partial oxidation of tin(II) trifluoroacetate: spectroscopic properties and *X*-ray crystal structure of di-μ_3-oxo-octakis-μ-trifluoroacetato-tetratin(II)tin(IV), 1675–82
X-Ray crystal structure and properties of dibenzotetrathiafulvalenium trichlorodiethylstannate(IV), 1793–6
Transition metal Schiff-base complexes as ligands in tin chemistry. Part 1. A tin-119 Mössbauer spectroscopic investigation of the adducts $SnX_4{\cdot}ML$, $SnMe_2(NCS)_2{\cdot}ML$, and $SnRCl_3{\cdot}ML$ (X = halide; M = Cu^{II} or Ni^{II}; L = quadridentate Schiff-base ligand; R = phenyl or n-butyl), 2261–6
TIN-119
Tin-119 nuclear magnetic resonance studies of some pseudohalogeno-derivatives of $[SnX_6]^{2-}$ (X = Cl or Br), 315–8
TITANIUM
Application of ultraviolet photoelectron spectroscopy to molecular properties. Part 27. Stereochemistry of ligand exchange reactions of titanocene derivatives, 107–10
Crystal and molecular structures of $[MCl(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf) and $[TiCl_{0.5}Br_{0.5}(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$: an unusual π-interaction with the metals and molecular orbital calculations on model compounds, 1641–6
Lamellar inorganic ion exchangers. Proton–copper(II) ion exchange in γ-titanium bis(hydrogenphosphate), 2045–50
Axially asymmetric metal alkyls. Part 5. Synthesis and reduction to Zr^{III} species of Group 4 metallepines $[ML(\eta\text{-}C_5H_5)_2]$[L = $(2\text{-}CHRC_6H_4)_2^{2-}$, R = H, M = Ti, Zr, or Hf), isomers of $[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), and *meso*-$[M\{1,8\text{-}(CHSiMe_3)_2C_{10}H_6\}(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf); *X*-ray crystal structures of $[TiL(\eta\text{-}C_5H_5)_2]$ (R = H) and λ-and δ-*RS*-$[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), 2347–58
TOSYLVALINATE
Spectroscopic, magnetic, and structural properties of diaquadimethanolbis(*N*-tosylvalinato)copper(II) and its 2,2′-bipyridine adduct, 1333–40
TRANSITION METAL
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 49. Synthesis of mixed-metal complexes with bonds between Cr, Mo or W and Co, Rh, Ir or Re; crystal structure of $[CrReRh_2(\mu_3\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}CO)(CO)_9(\mu\text{-}C_9H_7)_2]$, 41–50
Synthesis and properties of heterotrimetallic complexes containing M–Pt–Hg (M = Mo, W, Fe, Mn, or Co) arrays, 119–22
Investigation into aroylhydrazones as chelating agents. Part 8. Synthesis and spectroscopic characterization of complexes of Co, Ni, Cu, Zn, and Cd with 2,6-diacetylpyridine bis(salicyloylhydrazone); *X*-ray crystal structure of dichloro[2,6-diacetylpyridine bis(salicyloylhydrazone)]cadmium(II)–chloroform–methanol (1/1/1), 177–82
Extended Hückel molecular orbital study of the effects of edge-bridging hydrogen atoms on the lengths of boron–boron and metal–metal bonds in cluster compounds, and the crystal structure of benzyltrimethylammonium octahydrotriborate(1−), 1017–26
Water photolysis. Part 2. An investigation of the relative advantages of various components of the sensitiser–electron relay–metal colloid system for the photoproduction of hydrogen from water, and the use of these systems in the photohydrogenation of unsaturated organic substrates, 1299–304
Concentrated aqueous solutions and liquid hydrates of zinc chloride: a study of their solvent behaviour using intershell, charge-transfer, and *d*–*d* adsorption spectra, 1485–8
Nitrogen-14 nuclear quadrupole resonance study of bonding in transition-metal cyanides, 1637–40
Synthesis of *SS*′-bis(4′-benzo[15-crown-5])dithioglyoxime and its complexes with copper(II), nickel(II), cobalt(II), cobalt(III), palladium(II), platinum(II), and platinum(IV), 1827–32
Molten lithium carbonate–sodium carbonate–potassium carbonate eutectic: electronic spectroscopy of first-row transition metal compounds, 2329–32
Metal(II) hexafluoroantimonates: preparation and characterization of $MF_2{\cdot}2SbF_5$ (M = Mg, Ni, Zn, Fe, Co, Cu, Cr, Ag, Cd, or Pb) and the *X*-ray structure determination of $AgF_2{\cdot}2SbF_5$, 2379–84
TRIAZACYCLONONANE
Preparation and magnetism of the binuclear iron(II) complexes $[\{Fe(C_9H_{21}N_3)X_2\}_2]$ (X = NCS, NCO, or N_3) and their reaction with NO. Crystal structures of $[\{Fe(C_9H_{21}N_3)(NCS)_2\}_2]$ and $[Fe(C_9H_{21}N_3)(NO)(N_3)_2]$, 187–92
Studies on transition-metal macrocyclic complexes. Single-crystal *X*-ray structure and electrochemistry of the bis-macrocyclic complex $[Cu(L)_2][ClO_4]_2{\cdot}2H_2O$ (L = 1,4,7-triazacyclononane), 373–8
TRIAZENIDE
Reduction–oxidation properties of organotransition-metal complexes. Part 26. The synthesis of paramagnetic triazenido-and aryl(1-aryliminoethyl)amido-bridged dirhodium carbonyl derivatives containing the $[Rh_2]^{3+}$ core, 1403–8
Reactions of tetra-μ-carboxylato-diruthenium(II,II) compounds. *X*-Ray crystal structures of $Ru_2(\mu\text{-}O_2CCF_3)_4(thf)_2$, $Ru_2(\mu\text{-}O_2CR)_4(NO)_2$ (R = Et or CF_3), and $\{Na_3[Ru_2(\mu\text{-}O_2CO)_4]{\cdot}6H_2O\}_n$, 2723–36
Reduction–oxidation properties of organotransition-metal complexes. Part 27. Mixed-valence triazenido-bridged di-iridium compounds, 2737–40
TRIBORATE
Extended Hückel molecular orbital study of the effects of edge-bridging hydrogen atoms on the lengths of boron–boron and metal–metal bonds in cluster compounds, and the crystal structure of benzyltrimethylammonium octahydrotriborate(1−), 1017–26
TRIETHOXYETHANE
Some binuclear molybdenum(VI) complexes obtained from condensation reactions: their crystal and molecular structures, 523–8
TRIFLUOROACETATE
Preparation and partial oxidation of tin(II) trifluoroacetate: spectroscopic properties and *X*-ray crystal structure of di-μ_3-oxo-octakis-μ-trifluoroacetato-tetratin(II)tin(IV), 1675–82
TRIFLUOROMETHYL
The methyl/trifluoromethyl substitution effect in boranes. A gas-phase electron diffraction and vibrational study of bis(dimethylamino)(trifluoromethyl)borane and dimethylaminobis(trifluoromethyl)borane, 1839–46
TRIMETHYLSILYL
Electron diffraction study of the molecular structure of *N*-methylbis(trimethylsilyl)amine in the gas phase, 785–8
Alkylation of tricarbonyliron complexes by trimethylsilyl cyanide: synthetic and kinetic studies, 2013–6
TRIMETHYLSILYLCYCLOPENTADIENYL
New dicyclopentadienyl phosphine, phosphite, and acetylene niobium(III) complexes. Crystal structure of $[Nb(\eta\text{-}C_5H_4SiMe_3)_2Cl(PhC{\equiv}CPh)]$, 975–80
Metallocene derivatives of early transition elements. Part 4. Synthesis and crystal structures of a series of zirconocene(IV) halides $[Zr(\eta\text{-}C_5H_4SiMe_3)_2X_2]$ (X = Cl or Br) and $[Zr\{\eta\text{-}C_5H_3(SiMe_3)_2\text{-}1,3\}_2X_2]$ (X = F, Br, or I), 1463–72

TRIOXADIAZABICYCLOICOSANE
A structural study of the complexation of the lithium ion by the cryptand 4,7,13-trioxa-1,10-diazabicyclo[8.5.5]icosane, 489–92
TRIS(DIMETHYLPHENYLSILYL)METHYL
The chemistry of trichloro[tris(trimethylsilyl)methyl] and trichloro[tris(dimethylphenylsilyl)methyl] complexes of gallium, indium, and thallium. Crystal and molecular structures of $[Li(thf)_2(\mu\text{-}Cl)_2GaCl\{C(SiMe_2Ph)_3\}]\cdot thf$, $[Li(thf)_3(\mu\text{-}Cl)InCl_2\{C(SiMe_3)_3\}]$, and $[\{(Me_3Si)_3C\}In(\mu\text{-}Cl)_2\{\mu\text{-}Fe(CO)_4\}In\{C(SiMe_3)_3\}]$ (thf = tetrahydrofuran), 747–56
TRIS(TRIMETHYLSILYL)METHYL
The chemistry of trichloro[tris(trimethylsilyl)methyl] and trichloro[tris(dimethylphenylsilyl)methyl] complexes of gallium, indium, and thallium. Crystal and molecular structures of $[Li(thf)_2(\mu\text{-}Cl)_2GaCl\{C(SiMe_2Ph)_3\}]\cdot thf$, $[Li(thf)_3(\mu\text{-}Cl)InCl_2\{C(SiMe_3)_3\}]$, and $[\{(Me_3Si)_3C\}In(\mu\text{-}Cl)_2\{\mu\text{-}Fe(CO)_4\}In\{C(SiMe_3)_3\}]$ (thf = tetrahydrofuran), 747–56
TRYPTOPHANATE
Metal ion-bridged intramolecular stacking interaction between the tryptophyl residue and the aromatic heterocyclic amine within the ternary complex (1,10-phenanthroline)(L-tryptophanato-*O*,*N*)copper(II) perchlorate–water (1/2.5): an *X*-ray study, 2017–22
TUMOUR
Regioselective synthesis of $[1\text{-}B_{10}H_9(SH)]^{2-}$ and $[2\text{-}B_{10}H_9(SH)]^{2-}$: potential agents for boron–neutron capture therapy of brain tumours, 1953–6
TUNGSTEN
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 50. Synthesis and crystal structures of the complexes $[RuW_3(\mu\text{-}CO)(\mu_3\text{-}\eta\text{-}CO)(\mu\text{-}CPh)_2(\mu_3\text{-}CPh)(CO)_2(\eta\text{-}C_5H_5)_3]$ and $[RuW_3(\mu\text{-}CO)(\mu_3\text{-}\eta\text{-}CO)\{\mu\text{-}C(Ph)C(O)\}(\mu\text{-}CPh)(\mu_3\text{-}CPh)(CO)(PMePh_2)(\eta\text{-}C_5H_5)_3]$, 51–60
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 51. Alkylidyne(carbaborane)tungsten-gold and -rhodium complexes; crystal structures of $[AuW(\mu\text{-}CR)(CO)_2(PPh_3)(\eta^5\text{-}C_2B_9H_9Me_2)]$, $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$, and $[RhW(\mu\text{-}CR)(CO)_2(PPh_3)_2\{\eta^5\text{-}C_2B_9(C_7H_9)H_8Me_2\}]$ (R = C_6H_4Me-4), 61–72
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 52. Synthesis and reactions of the compounds $[RuW(\mu\text{-}CR)(CO)_3(\eta\text{-}C_5H_5)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[RuW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta\text{-}C_5H_5)(\eta^5\text{-}C_2B_9H_9Me_2)]$ and $[N(PPh_3)_2][RuW(\mu\text{-}CC_6H_4Me\text{-}4)(\mu\text{-}\sigma,\eta^5\text{-}C_2B_9H_8Me_2)(CO)_3(\eta\text{-}C_5H_5)]$, 73–80
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 53. Alkylidyne(carbaborane)dimetal complexes with tungsten–molybdenum and –tungsten bonds; crystal structures of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ [M = Mo (two isomers) or W], 81–90
Bimetallic systems. Part 16. Formation of Group 6 metal(0)–platinum(II) hydride complexes, by ring opening of *fac*- or *mer*-$[M(CO)_3(dppm\text{-}PP')(dppm\text{-}i')]$ (M = Cr, Mo, or W; dppm = $Ph_2PCH_2PPh_2$). Reactions of $[(CO)_3Mo(\mu\text{-}dppm)_2PtH(Cl)]$, 221–6
The addition of isonitriles to $[\{M(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (M = Mo or W). Syntheses and properties of $[M_2(\mu\text{-}\eta^2\text{-}CNR)(CO)_4(\eta\text{-}C_5H_5)_2]$ (M = Mo or W, R = Me or Bu^t) and $[\{Mo(CNR)(CO)_2(\eta\text{-}C_5H_5)\}_2]$ (R = Me or Bu^t). *X*-Ray crystal structures of $[Mo_2(\mu\text{-}\eta^2\text{-}CNBu^t)(CO)_4(\eta\text{-}(\eta\text{-}C_5H_5)_2]$ and $[\{Mo(CNBu^t)(CO)_2(\eta\text{-}C_5H_5)\}_2]$, 341–8
Syntheses and reactions of acyclic and cyclic carbene complexes *cis*-$[MI(\equiv CMeNHMe)(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo or W) and *cis*-$[MI\{=\overline{C(CH_2)_nNMe}\}(CO)_2(\eta\text{-}C_5H_5)]$ (M = Mo, *n* = 3 or 4; M = W, *n* = 3) obtained through alkyl migration to methyl isocyanide co-ordinated to molybdenum or tungsten. *X*-Ray structure of *cis*-$[MI\{=\overline{C(CH_2)_3NMe}\}(CO)_2(\eta\text{-}C_5H_5)]$ (1986, 2127), 479–80
Kinetics and mechanism of ring closure for a series of photoproduced $[M(CO)_5L]$ intermediates (M = Cr, Mo, or W; L = pyridine 2-carbaldehyde imine ligand), 513–8
Reactions of co-ordinated ligands. Part 39. The synthesis of carbyne complexes from alkyne molybdenum and tungsten cations; formation and crystal structures of $[Mo(\equiv CCH_2Bu^t)\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$ and $[\overline{Mo\{=C(SiMe_3)CH_2\}}\{P(OMe)_3\}_2(\eta^5\text{-}C_9H_7)]$, 591–604
Intercalation compounds of (diphosphato)oxotungsten(VI), $[WO(P_2O_7)]$, 609–12
Heteronuclear 1,3-intramolecular metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of 2-thia-4-selenapentane. A dynamic nuclear magnetic resonance investigation, 757–60
Reactions of aluminopolyhydride complexes of tungsten. *X*-Ray crystal structures of $[(Me_3P)_3H_3W(\mu\text{-}H)_2AlCl(\mu\text{-}N{=}CHEt)]_2$, $\{[(Me_3P)_3WH_5]Li\}_4$, $WH_2I(PMe_3)_3(SiMe_3)$, and $(Me_3P)_3H_2W(\mu\text{-}H)_3AlCl_2(NMe_3)$, 837–46
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η^2-vinyl complexes of tungsten, $[W\{\eta^3\text{-}C(CF_3)C(CF_3)(SR)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (R = Me, Et, Pr^n, Pr^i, or Bu^t), 895–906
Photo-oxidations of acetoin and biacetyl catalysed by tetrabutylammonium decatungstate(4−) in acetonitrile under excess of oxygen, 961–2
Studies on transition-metal peroxo complexes. Part 7. Molybdenum(VI) and tungsten(VI) carboxylato peroxo complexes, and the *X*-ray crystal structure of $K_2[MoO(O_2)_2(glyc)]\cdot 2H_2O$, 991–6
Carbene complexes in organic synthesis: a new tungstacarbene-promoted alkyne insertion–olefin cyclopropanation reaction, 1041–50
Reactions of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}X_2]$ (M = Mo, X = I; M = W, X = Cl) with some amine-substituted pyridines. The molecular structures of $[M(NO)\{HB(3,5\text{-}Me_2C_3N_2H)_3\}(2\text{-}NHC_5NH_4)_2]$, 1063–70
Chemistry of metallacyclic complexes containing the $\overline{FeCSCS}$ ring obtained by metal-promoted CS_2–CS coupling. *X*-Ray crystal structures of $[(CO)(cp)\overline{FeC\{SW(CO)_5\}SC(FeL_n)S}]$ and $[(CO)(cp)\overline{FeC(SEt)SC(FeL_n)S}]SO_3CF_3$ [FeL_n = Fe(cp)$(CO)_2$, cp = η-C_5H_5], 1133–44
Dimeric tungsten(V) compounds containing $[(S)W(\mu\text{-}S)_2W(S)]^{2+}$. Syntheses and structures of di-μ-sulphido-bis[(diethyl dithiophosphato-*SS'*)sulphidotungsten(V)] and di-μ-sulphido-bis[(*N*,*N*-diethyldithiocarbamato-*SS'*)sulphidotungsten(V)], 1163–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 54. Compounds derived from $[W(\equiv CC_6H_4Me\text{-}4)(CO)_2(\eta\text{-}C_5Me_5)]$ with bonds between tungsten and iron, cobalt, or nickel; crystal structure of $[FeW\{\mu\text{-}C(C_6H_4Me\text{-}4)C(O)O\}(CO)_5(\eta\text{-}C_5Me_5)]$, 1191–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 55. Coupling of methylene and alkylidyne groups at a trimetal centre; crystal structures of $[Ru_2W\{\mu\text{-}\sigma\text{:}\eta\text{-}trans\text{-}CH{=}C(H)R\}(\mu\text{-}CO)(CO)_3(\eta\text{-}C_5H_5)_2(\eta\text{-}C_5Me_5)]\cdot CH_2Cl_2$ and $[Ru_2W(\mu_3\text{-}CCH_2R)(\mu\text{-}CO)_3(CO)(\eta\text{-}C_5H_5)_3]$ (R = C_6H_4Me-4), 1201–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 58. Some reactions of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (M = Mo or W); crystal structures of the complexes $[W_2(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_2(L)(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$-[L = $N_2C(C_6H_4Me\text{-}4)_2$ or O], 1221–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 59. Reactions of the alkylidyne compounds $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Cr, Mo, or W, R = C_6H_4Me-4; M = W, R = Me, $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate] with d^{10} complexes; crystal structure of $[PtW_2(\mu\text{-}CMe)_2(CO)_4\{HB(pz)_3\}_2]$, 1229–34
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 60. Di- and tri-metal compounds prepared from the metal alkylidyne complexes $[W(\equiv CR)(CO)_2(\eta\text{-}C_5Me_5)]$ (R = Me or C_6H_4Me-4) and $[M(\equiv CR)(CO)_2\{HB(pz)_3\}]$ [M = Mo, R = C_6H_4Me-4; M = W, R = Me; $HB(pz)_3$ = hydrotris(pyrazol-1-yl)borate], 1235–42
Triaquahenicosatungstodiphosphate(6−) heteropolyanion, $[P_2W_{21}O_{71}(OH_2)_3]^{6-}$: *X*-ray crystallographic and ^{183}W nuclear magnetic resonance structural studies (1986, 2237), 1283–4
Bimetallic systems. Part 19. Chromium(0)–, molybdenum(0)–, or tungsten(0)–iron(0) carbonyl complexes containing one or two bridging $Ph_2PCH_2PPh_2$ ligands: crystal structure of $[(OC)_3Mo(\mu\text{-}Ph_2PCH_2PPh_2)_2Fe(CO)_3]$, 1489–94

TUNGSTEN (contd)
Bimetallic systems. Part 20. Synthesis of Group 6 metal(0)–iridium(I) or –rhodium(I) complexes bridged by $Ph_2PCH_2PPh_2$, 1495–500
Involvement of hydrogen-bonding protons in delocalization of the paramagnetic electron in a single crystal of photoreduced decatungstate, 1597–604
Synthesis of (η-cyclopentadienyl)(trimethylphosphine)tungsten derivatives and related chemistry, 1611–8
Tungsten(VI) equilibria: a potentiometric and calorimetric investigation, 1701–6
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato-molybdenum and -tungsten nitrosyls. Part 4. The effects of ligating atom type on the reduction potentials of monometallic complexes, 1733–6
Preparation and characterisation of dicarboxylatobis(η-cyclopentadienyl)tungsten(IV) complexes: a novel and convenient synthesis from $[W(\eta\text{-}C_5H_5)_2H_2]$ using molecular oxygen as a hydrogen acceptor, 1857–60
Bis(organoimido)-complexes of tungsten(VI): the crystal and molecular structures of tetrachlorobis(t-butylamine)bis(t-butylimido)bis(μ-*p*-tolylimido)-ditungsten(VI) and tetrachlorobis(μ-phenylimido)-bis(t-butylamine)bis(t-butylimido)ditungsten(VI), 2059–66
Preparation and characterisation of mononuclear organoimido complexes of tungsten: *X*-ray crystal structures of $[WCl_4(NC_6H_4Me\text{-}p)(thf)]$, $[P(CH_2Ph)Ph_3][WCl_5(NC_6H_4Me\text{-}p)]$, and $[WCl_2(NPh)_2(bipy)]$, 2067–76
Reactions of main group homopolynuclear cations with transition-metal carbonyls. Stabilisation of μ-Se_2^{2+} and η³-Te_3^{2+} dicationic ligands and crystal structures of $[Mo(Te_3)(CO)_4][SbF_6]_2$ and $[FeW(Se_2)(CO)_8][SbF_6]_2$, 2111–6
Synthesis and reactions of the co-ordinatively unsaturated hexafluorobut-2-yne complex $[WBr_2(CO)(CF_3C{\equiv}CCF_3)_2]$. The crystal and molecular structure of $[\{W[C(CF_3){=}C(CF_3)P(OEt)_2O](\mu\text{-}Br)(CO)(CF_3C{\equiv}CCF_3)\}_2]$, 2165–70
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 62. Synthesis of penta-, hexa-, and hepta-heteronuclear metal cluster compounds involving tungsten or molybdenum with platinum or nickel: crystal structures of the chain compounds $[Pt_3W_3(\mu\text{-}CMe)(\mu_3\text{-}CMe)_2(CO)_6(cod)(\eta\text{-}C_5Me_5)_3]\cdot 0.5CH_2Cl_2$ and $[Pt_3W_4(\mu\text{-}CC_6H_4Me\text{-}4)_2(\mu_3\text{-}CC_6H_4Me\text{-}4)_2(CO)_8(\eta\text{-}C_5H_5)_4]$, 2177–88
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 63. Synthesis of eight-membered-ring metallacycles: *X*-ray crystal structures of $[Pt_4W_4(\mu\text{-}CR)(\mu_3\text{-}CR)_3(\mu\text{-}CO)(CO)_7(\eta\text{-}C_5H_5)_4]\cdot CH_2Cl_2$, $[Ni_2Pt_2W_4(\mu\text{-}CR)(\mu_3\text{-}CR)_3(\mu\text{-}CO)(CO)_7(\eta\text{-}C_5H_5)_4]\cdot CH_2Cl_2$ (R = C_6H_4Me-4), and $[Ni_2Pt_2W_4(\mu_3\text{-}CPh)_4(CO)_8(\eta\text{-}C_5H_5)_4]$, 2189–200
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 65. Reactions of the complexes $[FeW(\mu\text{-}CR)(CO)_5(\eta\text{-}C_5Me_5)]$ and $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5Me_5)]$ (R = C_6H_4Me-4) with alkynes R′C_2R′ (R′ = Me or Ph); crystal structures of $[FeW\{\mu\text{-}C(R)C(O)C(Me)C(Me)\}(CO)_5(\eta\text{-}C_5Me_5)]\cdot CH_2Cl_2$ and $[FeW\{\mu\text{-}C(R)C(Et)C(H)C(Me)C(Me)\}(\mu\text{-}CO)(CO)_3(\eta\text{-}C_5Me_5)]$, 2211–8
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 66. Carbaboranetungsten–platinum complexes. Polyhedral rearrangements of a 12-vertex cage system; crystal structures of $[PtW(CO)_2(PEt_3)_2\{\eta^6\text{-}C_2B_9H_8\text{-}(CH_2C_6H_4Me\text{-}4)Me_2\}]\cdot CH_2Cl_2$, $[PtW(\mu\text{-}H)\{\mu\text{-}\sigma{:}\eta^5\text{-}C_2B_9H_7(CH_2C_6H_4Me\text{-}4)Me\}(CO)_2(PMe_3)(PEt_3)_2]$, and related compounds, 2219–34
Preparation and properties of the tungsten(V) aqua dimer $[W_2O_4(H_2O)_6]^{2+}$: kinetic studies on the 1:1 reaction with NCS^-, 2293–8
Determination of the relative signs of $^2J(^{31}P\text{–}^{31}P)$ in complexes of tungsten(0) and molybdenum(0) using two-dimensional [^{31}P–^{31}P]-COSY-45 nuclear magnetic resonance chemical shift correlation, 2401–6
Dithiocarbamates of tungsten(III) and tungsten(IV), 2531–6
Monometallic, homo- and hetero-bimetallic complexes based on redox active tris(3,5-dimethylpyrazolyl)borato molybdenum and tungsten nitrosyls. Part 6. Synthesis and electrochemistry of complexes containing saturated heterocyclic amide ligands. *X*-Ray crystal and molecular structure of $[\{Mo(NO)[HB(Me_2pz)_3]Cl\}_2(NC_4H_8N)]$, 2653–60
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 67. Reactions of the salts $[X][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ [X = $P(CH_2Ph)Ph_3$, R = C_6H_4Me-4; X = NEt_4, R = Ph] with bis(cyclo-octa-1,5-diene)platinum, 2661–8
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η²-vinyl complexes of molybdenum and tungsten, $[M\{\eta^3\text{-}C(CF_3)C(CF_3)(L\text{–}L)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (L–L = $MeCO_2^-$, $Me_2PS_2^-$, $Et_2NCS_2^-$, or $Me_2NCS_2^-$), 2715–22
Heteronuclear complexes containing Group 6 transition metals; chemical, spectroscopic, and theoretical studies on some binuclear complexes and the *X*-ray crystal structure determination of $[WNi(CO)_3(PPh_3)_2(\eta^5\text{-}C_5H_5)]$, 2741–50
Carbene complexes. Part 21. Synthesis and characterisation of bis(carbene)molybdenum(II) complexes and dimetal(0) complexes of the Group 6 elements containing novel bridging bis(carbene) ligands; *X*-ray structures of $[Mo(CO)_2(L^{Et})_2(OSO_2CF_3)_2]$ [L^{Et} = ≡CN(Et)$(CH_2)_2$NEt] and $(W(CO)_5\{C(OEt)\text{-}CH_2C_6H_4CH_2C(OEt)\text{-}o\}W(CO)_5]$, 2843–52
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 68. Reactions between nonacarbonyldi-iron and the salts $[NEt_4][W(\equiv CR)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ (R = Me, Ph, or C_6H_4Me-4); crystal structures of $[NEt_4][FeW\{\mu\text{-}CH(C_6H_4Me\text{-}4)\}(\mu\text{-}\sigma{:}\eta^5\text{-}C_2B_9H_8Me_2)(\mu\text{-}CO)(CO)_5]$ and $[NEt_4][FeW(\mu_3\text{-}CPh)(\mu\text{-}\sigma{:}\sigma',\eta^5\text{-}C_2B_9H_7Me_2)(CO)_8]$, 2917–26
Instant ligands. Part 2. Preparation of some bidentate ligands containing PF_2O Groups from aromatic hydroxy compounds and their reactions to form molybdenum and tungsten complexes, 3023–8
The chemistry of cycloheptatrienyl complexes of molybdenum and tungsten: the synthesis and reactions of some vinylidene and alkynyl derivatives, 3049–54
TUNGSTEN-183
Triaquahenicosatungstodiphosphate(6−) heteropolyanion, $[P_2W_{21}O_{71}(OH_2)_3]^{6-}$: *X*-ray crystallographic and ^{183}W nuclear magnetic resonance structural studies (1986, 2237), 1283–4
UNDECABORATE
Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 58. Some reactions of the compounds $[MW(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_3(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$ (M = Mo or W); crystal structures of the complexes $[W_2(\mu\text{-}CC_6H_4Me\text{-}4)(CO)_2(L)(\eta^5\text{-}C_9H_7)(\eta^5\text{-}C_2B_9H_9Me_2)]$-[L = $N_2C(C_6H_4Me\text{-}4)_2$ or O], 1221–8
URANIUM
Reaction of monoborane with tris(cyclopentadienyl)uranium(IV) derivatives, 691–4
Uranyl(VI) complexes of 3-aminopropanoic acid and 4-aminobutanoic acid in aqueous solution: a potentiometric and calorimetric study, 695–8
Insertion of 2,6-dimethylphenyl isocyanide into the U–N bond of $[U(\eta\text{-}C_5H_5)_3(NEt_2)]$. Crystal structure of $[U(\eta\text{-}C_5H_5)_3\text{-}\{C(NEt_2){=}NC_6H_3Me_2\text{-}2,6\}]$, 2039–44
Infrared spectroscopic studies of uranyl(VI) species adsorbed from aqueous $[UO_2(CO_3)_3]^{4-}$ solutions on to a polymer bearing amidoxime groups, 2489–92
Ligand intra- and inter-molecular exchange processes in dioxopentakis(trimethylurea)uranium(VI). A proton nuclear magnetic resonance study, 2885–8
Surface characterisation of α-, β-, γ-, and δ-UO_3 using *X*-ray photoelectron spectroscopy, 3009–16
UREA
Ligand intra- and inter-molecular exchange processes in dioxopentakis(trimethylurea)uranium(VI). A proton nuclear magnetic resonance study, 2885–8
VALENCE
Ab initio study of stable bis(carbon dioxide)molybdenum complexes, 2373–8
VANADIUM
The single-crystal electron spin resonance spectrum of oxobis(2-methylquinolin-8-olato)vanadium(IV) as a pure compound and diluted in chlorobis(2-methylquinolin-8-olato)gallium(III), 111–8
Water-soluble Schiff-base complexes of vanadyl(IV) and vanadium(III), 1279–82
Reaction of elemental sulphur with bis(pentamethylcyclopentadienyl)vanadium derivatives: disulphido and μ-sulphido complexes. Crystal structures of $[V(\eta\text{-}C_5Me_5)_2(S_2)]$ and $[V_2(\eta\text{-}C_5Me_5)_2Cl_2(\mu\text{-}S)_2]$, 2099–104

VANADIUM (contd)
Electronic structure and redox reactions of vanadium(II) polypyridine, isocyanide, and phosphine complexes, 2881–4
VINYL
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η^2-vinyl complexes of tungsten, $[W\{\eta^3\text{-}C(CF_3)C(CF_3)(SR)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (R = Me, Et, Pr^n, Pr^i, or Bu^t), 895–906
The reaction of $[Mn_2\{\sigma{:}\eta^2\text{-}C(R){=}CHR\}(\mu\text{-}PPh_2)(CO)_7]$ (R = H or Ph) with nucleophiles: *X*-ray crystal structures of $[N(PPh_3)_2][Mn_2(\mu\text{-}H)\{\mu\text{-}C(O)C(Ph){=}CHPh\}(\mu\text{-}PPh_2)(CO)_6]$, $[Mn_2\{PPh_2C(Ph){=}CHPh\}(CO)_9]$, $[Mn_2(\mu\text{-}PPh_2\text{-}CH{=}CH_2)(CO)_8]$, $[Mn_2\{\mu\text{-}C(O)C(Ph){=}CHPh\}(\mu\text{-}PPh_2)(CO)_6(CNBu^t)_2]$, $[Mn_2\{\mu\text{-}\sigma{:}\eta^2\text{-}C(Ph){=}CHPh\}(\mu\text{-}PPh_2)(CO)_6(CNBu^t)]$, and $[Mn(\mu\text{-}PPh_2CH{=}CH_2)(CO)_7(PEt_3)]$, 2669–82
Synthesis, reactions, and dynamic nuclear magnetic resonance studies of η^2-vinyl complexes of molybdenum and tungsten, $[M\{\eta^3\text{-}C(CF_3)C(CF_3)(L\text{–}L)\}(CF_3C{\equiv}CCF_3)(\eta^5\text{-}C_5H_5)]$ (L–L = $MeCO_2^-$, $Me_2PS_2^-$, $Et_2NCS_2^-$, or $Me_2NCS_2^-$), 2715–22
VINYLIDENE
Reactions of co-ordinated ligands. Part 41. Synthesis and reactions of cationic functionalised alkyne molybdenum complexes; formation of carbyne and vinylidene complexes and crystal structure of $[Mo(C{=}CHPh)Br\{P(OMe)_3\}_2(\eta\text{-}C_5H_5)]$, 1319–28
Bimetallic systems. Part 21. Synthesis of vinylidene- and phenylene-bridged complexes of dinickel(II), nickel(II)–palladium(II), and nickel(II)–platinum(II) containing $Ph_2PCH_2PPh_2$ (dppm): crystal structures of $[(SCN)Ni(\mu\text{-}dppm)_2(\mu\text{-}C{=}CH_2)Ni(NCS)]$ and $[ClPd(\mu\text{-}dppm)_2(\mu\text{-}C{=}CCl_2)NiCl]$, 1501–8
The chemistry of cycloheptatrienyl complexes of molybdenum and tungsten: the synthesis and reactions of some vinylidene and alkynyl derivatives, 3049–54
VINYLIMIDAZOLE
Synthesis, characterisation, and photochemical properties of a series of ruthenium containing metallopolymers based on poly-*N*-vinylimidazole, 3073–8
VITAMIN
The chemistry of vitamin B_{12}. Part 27. The cage phosphite ester 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane and the anionic dimethyl phosphite as ligands for cobalt(III) corrinoids, 761–8
VITAMIN B_6
Ternary metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of copper(II)–2,2′-bipyridyl– pyridoxic acid and copper(II)–2,2′-bipyridyl–pyridoxic acid γ-lactone complexes, 1009–16
WATER
Water photolysis. Part 2. An investigation of the relative advantages of various components of the sensitiser–electron relay–metal colloid system for the photoproduction of hydrogen from water, and the use of these systems in the photohydrogenation of unsaturated organic substrates, 1299–304
***X*-RAY DIFFRACTION**
Studies on alkaline-earth metal sulphites. Part 9. Solid solutions of $Ca_3(SO_4)_2SO_3{\cdot}12H_2O$ in $Ca_3(SO_3)_2SO_4{\cdot}12H_2O$: structure, stability, and dehydration products, 235–40
Structure and solvation of mercury(II) iodide and bromide in tetrahydrothiophene solution, 2411–6
Surface characterisation of α-, β-, γ-, and δ-UO_3 using *X*-ray photoelectron spectroscopy, 3009–16
YLIDE
Synthesis and characterization of palladium(II)–η^3-allyl–ylide complexes. *X*-Ray crystal structure of $[PdCl(\eta^3\text{-}2\text{-}MeC_3H_4)\{Ph_3PC(H)COMe\}]$, 1381–8
Chemistry of 2,3-dihydro-2-metallaphenalenes. Part 2. Formation of $\overline{CHMMe_2C(MMe_3)C_{10}H_6}]^x$ *via* base decomposition of $[1,8\text{-}(Me_3MCH_2)_2C_{10}H_6]^y$ (M = Ge, $x = 2-$, $y = 0$; M = P, $x = 0$, $y = 2+$); crystal structures of the novel bis-ylide $\overline{CHPMe_2C(PMe_3)C_{10}H_6}$ and of a distorted *peri*-ylide, 1-Me-8-$(Me_3PCH)C_{10}H_6$, 1647–54
ZEOLITE
Synthesis and properties of gallium tetramethylammonium sodalite, $NMe_4GaSi_5O_{12}$, 623–6
Exchange of cadmium into the sodium and ammonium forms of the natural zeolites clinoptilolite, mordenite, and ferrierite, 1911–6
ZINC
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 5. Unsubstituted fifteen-membered-ring macrocycles containing 2,2′:6′,2″-terpyridine, 5–8
Investigation into aroylhydrazones as chelating agents. Part 8. Synthesis and spectroscopic characterization of complexes of Co, Ni, Cu, Zn, and Cd with 2,6-diacetylpyridine bis(salicyloylhydrazone); *X*-ray crystal structure of dichloro[2,6-diacetylpyridine bis(salicyloylhydrazone)]cadmium(II)–chloroform–methanol (1/1/1), 177–82
Proton and phosphorus-31 nuclear magnetic resonance study of zinc(II) *O*,*O*′-dialkyl dithiophosphates in solution, 807–14
Synthesis of pyridine-containing tetra-aza macrocycles: 3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), its 3,11-dibenzyl (L^2) and 3,7,11-tribenzyl (L^3) derivatives, and their nickel(II), copper(II), and zinc(II) complexes: crystal structures of $L^2{\cdot}HCl$ and $[Ni(L^2)Cl]ClO_4{\cdot}H_2O$, 889–94
Electron spin resonance and optical spectroscopic studies on Cu^{2+} and Co^{2+} in $Cs_2[Zn(N_3)_4]$, 939–42
Studies of the pyridine-containing tetra-azamacrocycle 3,7,11-trimethyl-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1) and its complexes with Ni^{2+}, Cu^{2+}, and Zn^{2+}. Crystal structures of the unsymmetric isomer of $[Ni(L^1)][ClO_4]_2$, and the five-co-ordinate complexes $[Ni(L^1)(dmso)][ClO_4]_2$ (dmso = dimethyl sulphoxide) and $[Ni(L^1)Cl]ClO_4$, 1107–14
Reactions of chromium(III) complexes of 1,10-phenanthroline, 2,2′-bipyridyl, and oxalate with the pulse radiolytically generated aquated electron, zinc(I), and cadmium(I), 1425–30
Concentrated aqueous solutions and liquid hydrates of zinc chloride: a study of their solvent behaviour using intershell, charge-transfer, and *d*–*d* adsorption spectra, 1485–8
Preparation and crystal structures of $[PPh_4]_2[M_2(SPh)_6]$ (M = Zn or Cd), 1577–80
Comments on the attempts by Wickens and Abrams to synthesize manganese(II) phosphine complexes: a comment, 1581–2
Ring-size and substituent effects in relatively rigid macrocyclic quinquedentate ligands. Structure determinations of $[ZnL^4(H_2O)_2][ClO_4]_2{\cdot}MeCN$ and $[ZnL^3(ClO_4)_2]$, 1631–6
Dithio-oxamides as ligands: crystal structures and vibrational analyses of bis(*N*,*N*′-dicyclohexyldithio-oxamidato)palladium(II), bis(*N*,*N*′-dibenzyldithio-oxamide)copper(II) diperchlorate, and dichloro(*N*,*N*′-dimethyldithio-oxamide)zinc(II) and the carbon-13 nuclear magnetic resonance spectra of related compounds, 1921–8
Thermal investigation and stereochemical studies of some cyclic diamine complexes of nickel(II), zinc(II), and cadmium(II) in the solid state, 2089–94
Polymeric thiolate complexes of Group 12 metals. Crystal and molecular structures of *catena*-[μ-(3-dimethylammonio-1-propanethiolate)]-dichlorocadmium(II) and bis[3-(dimethylammonio)propyl] disulphide tetrabromocadmate(II), 2391–6
Nickel(II), copper(II), and zinc (II) complexes of penta-azamacrocyclic ligands. Crystal and molecular structure of 3,6,9,12-tetramethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-trienezinc(II) perchlorate, 2643–8
Electrolytic behaviour of transition metal bromides in acetonitrile solution, 2863–6
Complexes formed between zinc(II) and diethylenetriamine: a carbon-13 nuclear magnetic resonance study, 2889–96
Carbon-13 nuclear magnetic resonance study of the zinc(II) complexes of 3,7-diazanonane-1,9-diamine and 1,2-diaminoethane, 2897–904
ZIRCONIUM
Metallocene derivatives of early transition elements. Part 4. Synthesis and crystal structures of a series of zirconocene(IV) halides $[Zr(\eta\text{-}C_5H_4SiMe_3)_2X_2]$ (X = Cl or Br) and $[Zr\{\eta\text{-}C_5H_3(SiMe_3)_2\text{-}1,3\}_2X_2]$ (X = F, Br, or I), 1463–72
Crystal and molecular structures of $[MCl(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf) and $[TiCl_{0.5}Br_{0.5}(CPh{=}CMe_2)(\eta\text{-}C_5H_5)_2]$: an unusual π-interaction with the metals and molecular orbital calculations on model compounds, 1641–6
Solid-state nuclear magnetic resonance spectroscopic study of γ-zirconium phosphate, 1877–82
Axially asymmetric metal alkyls. Part 5. Synthesis and reduction to Zr^{III} species of Group 4 metallepines $[ML(\eta\text{-}C_5H_5)_2][L = (2\text{-}CHRC_6H_4)_2^{2-}$, R = H, M = Ti, Zr, or Hf), isomers of $[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), and *meso*-$[M\{1,8\text{-}(CHSiMe_3)_2C_{10}H_6\}(\eta\text{-}C_5H_5)_2]$ (M = Zr or Hf); *X*-ray crystal structures of $[TiL(\eta\text{-}C_5H_5)_2]$ (R = H) and λ-and δ-*RS*-$[ZrL(\eta\text{-}C_5H_5)_2]$ (R = $SiMe_3$), 2347–58